

DISCONTINUOUS GAS EXCHANGE CYCLES IN *APHODIUS FOSSOR* (SCARABAEIDAE): A TEST OF HYPOTHESES CONCERNING ORIGINS AND MECHANISMS

STEVEN L. CHOWN^{1,*} AND PETER HOLTER²

¹Department of Zoology and Entomology, University of Pretoria, Pretoria 0001, South Africa and ²Department of Terrestrial Ecology, University of Copenhagen, Denmark

*e-mail: slchown@zoology.up.ac.za

Accepted 25 October; published on WWW 22 December 1999

Summary

It has been suggested that discontinuous gas exchange cycles (DGCs) in insects evolved in response to hypoxic and/or hypercapnic environments, and that the flutter phase of the DGC is predominantly diffusive. We tested these hypotheses by examining the rate of CO₂ release at a declining series of oxygen partial pressures in adults of *Aphodius fossor*, an inhabitant of moist to wet dung pats that can have low oxygen and high carbon dioxide concentrations. This mesic species showed a pronounced DGC, but progressively abandoned it when exposed to declining oxygen concentrations. That is, in response to a reduction in oxygen concentration, the closed phase declined in length and eventually disappeared, the flutter phase declined in length, while the flutter phase rate of CO₂ release increased, the open phase peak rate of CO₂ release declined and DGC frequency increased. Nonetheless,

overall CO₂ release rate remained virtually unchanged. Thus, it appears that the flutter phase in *A. fossor* has a significant convective component. In addition, in response to declining oxygen concentrations, the spiracles are opened for prolonged periods to increase gas exchange. This behaviour is unlikely to incur a significant water loss penalty in the moist habitats in which *A. fossor* lives. Given the switch to continuous ventilation by *A. fossor* under hypoxic conditions, we conclude that current adaptive explanations for the DGC are inadequate. Rather, the DGC may be the default state of a system regulated by two interacting feedback loops when demands are absent.

Key words: discontinuous gas exchange, ventilation, water loss, hypoxia, beetle, CO₂ release, *Aphodius fossor*.

Introduction

Discontinuous gas exchange cycles (DGCs) are known from a variety of insects and other arthropods. The DGC was first thought to have originated primarily as a means of reducing water loss because respiratory water loss rates would be zero during the closed (C) phase, and much reduced if the flutter (F) phase were solely or predominantly convective, as has been demonstrated in moth pupae, cockroaches and some ants (Levy and Schneiderman, 1966; Kestler, 1985; Lighton et al., 1993a). Such a restriction of respiratory transpiration rates would be particularly important where cuticular water loss rates had already evolved to low levels (for further discussion, see Hadley, 1994; Lighton, 1994, 1996). However, more recently, this idea has been questioned (see Hadley and Quinlan, 1993; Quinlan and Hadley, 1993; Hadley, 1994; Lighton, 1994, 1996; Lighton and Garrigan, 1995).

Lighton and Berrigan (1995) have proposed that the DGC may originally have evolved to enhance gas exchange efficacy, without the excessive water loss penalty associated with continuous gas exchange (for details, see Lighton, 1996), under hypoxic and hypercapnic circumstances. These are especially

likely to be encountered by psammophilous beetles and subterranean ants. Lighton (1996) argued that although the DGC could certainly be co-opted as a strategy to combat water loss (an exaptation; for a discussion, see Endler, 1986), especially where the F phase is predominantly convective (as in lepidopteran pupae), large fluctuations in an insect's internal environment associated with a DGC would impose physiological costs that species might seek to avoid. Experimental work by Lighton and Berrigan (1995) and a broad comparative analysis by Lighton (1996) provide some support for this chthonic origination hypothesis. Nonetheless, Lighton (1996) concluded that considerably more descriptive and experimental work, on a broader range of taxa, is required if the evolutionary genesis of the DGC is to be unravelled successfully.

Much of this work will undoubtedly have to focus on the factors that modulate the duration of the three phases of the DGC and the nature of the dominant gas exchange modality during the F phase (see above and Lighton, 1996). Lighton (1988, 1994, 1996), Lighton and Berrigan (1995) and Lighton

and Garrigan (1995) have concluded that at least in some tenebrionid beetles and in some ants, especially *Camponotus vicinus*, the F phase is primarily diffusive. They demonstrated this using simultaneous CO₂ and O₂ flow-through respirometry in the case of beetles (Lighton, 1988) and by exposing ants to various oxygen concentrations (Lighton and Garrigan, 1995). In the latter case, Lighton and Garrigan (1995) argued that because ambient O₂ levels determine the maximum molar concentration of oxygen available at the onset of the C phase, hypoxia should trigger an early switch to the F phase. Similarly, if convection predominates during the F phase, increasing levels of nitrogen will impede negative endotracheal pressure generation, thus restricting the inward movement of oxygen, which reduces F phase duration. In contrast, if the F phase is predominantly diffusion-based, then hypoxia must elevate CO₂ emission levels, causing a delay in the CO₂-mediated switch to the open (O) phase, thus increasing F phase duration. Lighton and Garrigan (1995) produced experimental support for the latter scenario and concluded that, in this case, the F phase is predominantly diffusive.

The aim of the present study is therefore to address both the evolutionary genesis and F phase gas exchange modality debates. To achieve this, we have focused our attention on *Aphodius fossor* (L.) (Scarabaeidae, Aphodiinae), an inhabitant of fresh cattle dung pats in the Holarctic (Dellacasa, 1983), a species ideally suited to address these questions. In Denmark, the adults of *A. fossor* are mostly active in the late spring/early summer (May–June). They fly mostly in the evening and at night, and occasionally during the day, and hence mostly at high ambient humidities (Landin, 1961, 1968; P. Holter, personal observations) (at Sjølsmark meteorological station approximately 12 km south-east of the site where *A. fossor* was collected for the present study, relative humidity ranges from 39 to 70 %, and mean temperatures from 9.0 to 17.6 °C; 10 year means, 1988–1997). Therefore, this species can be thought of as mesic. In addition, the adult beetles spend most of their lives in cattle dung pats that have water contents in excess of 80 % (Holter, 1982). Within the dung pat, they are not confined to well-defined zones (Holter, 1982), and in wet pats, where microbial activity is high, may consequently be exposed to oxygen concentrations as low as 2 % and CO₂ levels in excess of 10 % (Holter, 1991). Indeed, laboratory trials have indicated that adult *A. fossor* can maintain unvarying oxygen uptake rates from normoxic conditions down to 1 % O₂ (Holter and Spangenberg, 1997).

On these grounds, we set out to test two major predictions. First, if the DGC originated primarily in response to hypoxic and/or hypercapnic environments, rather than as a strategy to reduce water loss, *A. fossor* should show a pronounced DGC, especially under hypoxic conditions. It lives in profoundly hypoxic/hypercapnic environments, but is unlikely to encounter dry conditions and, even then, it has ready access to a water supply. Second, because it possesses flaccid, membranous abdominal tergites (as does the tenebrionid *Psammodes striatus*), modulation of the DGC in response to

hypoxia should reveal a largely diffusion-based F phase (see Lighton, 1988, 1996).

Materials and methods

Adults of *Aphodius fossor* (L.) were collected from dung pats in the Strødam Nature Reserve (55°58'N, 12°16'E), 35 km to the north west of Copenhagen, Denmark. Beetles were returned to the laboratory in Copenhagen within 4 h of collection and transferred individually to 185 ml plastic containers containing moist potting soil. The beetles were kept for a minimum of 3 days at 15±1 °C, the temperature at which all subsequent work was performed.

Following this period, CO₂ release was measured and gas exchange characteristics were examined using a Sable Systems (Henderson, Nevada, USA) respirometry system. Air was passed through Drierite and soda-lime columns to remove water vapour and CO₂. It was then passed through an automated baselining system, through the 10 ml cuvette containing the beetle, and finally through a Li-Cor CO₂/H₂O analyzer (model Li 6262; 1–2 p.p.m. accuracy for both channels at 350 p.p.m.) at a flow rate of 100 ml min⁻¹. Datacan V software was used for data capture and analysis. All measurements were corrected to standard temperature and pressure and are expressed as ml CO₂ g⁻¹ h⁻¹ or mg H₂O g⁻¹ h⁻¹. Respiration rate and gas exchange characteristics were first examined using ambient air. Thereafter, O₂ concentrations of approximately 15, 10 and 5 % in nitrogen (scrubbed of all water) were used. These concentrations were obtained by mixing O₂ and N₂ delivered by two Sierra Instruments mass-flow meters, both controlled by a Mikrolab Aarhus flow controller. Oxygen concentrations were subsequently assessed using a gas chromatograph (Mikrolab Aarhus) with thermal conductivity detector at 120 °C and a Molesieve 5A column (1.8 m×2 mm). The treatments were randomized by day, and five beetle specimens were examined per day. Within treatments, the order of examination of beetles was also randomized. Following this, beetles were finally examined at 2 % O₂ in nitrogen, again with the order of examination randomized within this treatment.

Individual beetles were weighed (using a Mettler AJ100 electronic microbalance sensitive to 0.1 mg), placed in the darkened cuvette 1 h before the treatment, after which each treatment/measurement was undertaken for 2 h in the absence of light, excluding the time required for automatic baseline determination (a further 30 min). On a variety of occasions, beetles were inspected for activity in the cuvette using a dim red light. In most instances, beetles were quiescent. In those cases in which they were active, this was also clearly visible on the DGC recording, and that particular measurement was discarded. After the measurements, beetles were reweighed. A final trial consisted of exposure of a further two beetles to a serial decline in O₂ concentration, over a 6 h period, from normoxic levels to approximately 2 %. Washout time, for each reduction in O₂ concentration, was determined prior to these

trials using the Li-Cor analyser with no beetle present in the cuvette, and was found to be approximately 6 min.

All calculations of CO₂ and H₂O release rates (on a mass-specific basis) and examinations of gas exchange characteristics were carried out using Sable Systems Datacan V software. For each individual, the mean CO₂ release rate (ml g⁻¹ h⁻¹), the frequency of the DGC, measured from O phase peak to O phase peak (mHz), and the mean height of these O phase peaks (ml g⁻¹ h⁻¹) were measured over the entire recording. Thereafter, mean CO₂ release rate and phase duration (s) were measured for each of the phases. Where a measure of emission volume (ml g⁻¹) was required, integration of the area under the curve against time (h) was undertaken. The resulting data were then used as input to Statistica for calculation of means and standard errors, for the determination of the percentage contribution of F phase CO₂ release rate to total CO₂ release rate and for investigating the relationships between phase characteristics and O₂ concentration using least-squares regression.

To compare our data on *A. fossor* with those obtained in a previous study by Holter and Spangenberg (1997), we used the rate of CO₂ release over the entire recording to estimate \dot{V}_{CO_2} , and converted this to μW (see Lighton and Fielden, 1995), assuming a respiratory quotient between 0.7 and 0.9.

To enable comparisons between water balance characteristics of these beetles and those determined previously for other Scarabaeidae (e.g. Zachariassen et al., 1987; Chown et al., 1995), the gravimetric protocol of Chown et al. (1995) was followed. Beetles were acclimated on moist filter paper for 2 days with no access to food. They were then placed individually into mesh-covered, open-ended plastic tubes (approximately 2 ml volume), which in turn were placed into 185 ml vials containing 100 ml of silica gel (providing a relative humidity of approximately 5%). Beetles were weighed at approximately 8 h intervals until they died. The mass recorded in the interval prior to death was used to calculate water loss rate, maximum water loss tolerated and time until death. No defecation was observed during these trials. Water loss rates obtained in this manner were compared with those calculated from the decrease in mass associated with 2 h exposures to dry air flowing over beetles at 100 ml min⁻¹ in the respirometry experiments. Because of the noise associated with infrared measurements, C phase and F




Fig. 1. Typical recording of discontinuous gas exchange in *Aphodius fossor* measured over 2 h at 15 °C under normoxia.

phase rates of H₂O release could not be adequately distinguished using data collected by the Li-Cor 6262. Hence, we were unable to characterize water loss during these phases. Nonetheless, we regarded the lowest value during the combined C/F phase as a measure of cuticular water loss, and subtracted this from O phase water loss to obtain an estimate of respiratory as opposed to cuticular water loss. This was performed for normoxic conditions only.

For all analyses, the significance level was set at $P < 0.05$, and values reported are means \pm S.E.M. throughout.

Results

Under normoxic conditions, gas exchange in *Aphodius fossor* (mean body mass 0.1213 ± 0.0034 g, $N=10$) was characterized by typical, discontinuous gas exchange cycles (Fig. 1). Under these conditions, standard metabolic rate, measured as mean CO₂ release rate over the entire recording was 0.1649 ± 0.0065 ml g⁻¹ h⁻¹. As ambient oxygen concentrations declined, mean CO₂ release rate remained constant, except at the lowest level (measured as 2.84% O₂), at which CO₂ release rate increased significantly and by approximately 23% over the mean value for \dot{V}_{CO_2} combined over all the higher O₂ concentrations (Table 1). Nonetheless, with declining ambient O₂ concentration, DGC frequency increased (duration declined) (DGC

Table 1. \dot{V}_{CO_2} , discontinuous gas exchange cycle (DGC) duration and frequency, and open phase peak height in *Aphodius fossor* under normoxic and declining ambient O₂ concentrations

O ₂ concentration (%)	<i>N</i>	\dot{V}_{CO_2} (ml g ⁻¹ h ⁻¹)	DGC period (s)	DGC frequency (mHz)	Open phase peak height (ml g ⁻¹ h ⁻¹)
20.47	10	0.165 \pm 0.007	999.4 \pm 115.4	1.146 \pm 0.144	0.673 \pm 0.028
16.55	9	0.203 \pm 0.015	514.6 \pm 69.9	2.271 \pm 0.312	0.659 \pm 0.041
11.28	10	0.195 \pm 0.014	285.0 \pm 16.7	3.615 \pm 0.208	0.468 \pm 0.032
6.13	10	0.203 \pm 0.016	218.4 \pm 21.9	5.080 \pm 0.568	0.406 \pm 0.031
2.84	10/3 ^a	0.236 \pm 0.013*	216.2 \pm 49.8	5.275 \pm 0.974	0.307 \pm 0.024

^aSample size at 2.84% was 10 for \dot{V}_{CO_2} and 3 for the other variables.

An asterisk denotes a significant difference from all other values at the 5% level using a Tukey HSD test ($F_{2,42}=4.171$, $P=0.006$).




Fig. 2. Changes in the characteristics of the discontinuous gas exchange cycle in *Aphodius fossor* during a sequential decline in oxygen concentration. Virtually instantaneous changes in O_2 concentration were made at the points indicated by the arrowheads. These are, from left to right: 20.47% to 16.55%, 16.55% to 11.28%, 11.28% to 6.13%, and 6.13% to 2.84%.

frequency = $-0.275[O_2] + 6.78$, $F_{2,39} = 99.63$, $P = 0.00001$, $r^2 = 0.714$) and O phase peak height declined (O phase peak CO_2 release = $0.022[O_2] + 0.255$, $F_{2,39} = 65.76$, $P = 0.00001$, $r^2 = 0.622$) significantly (Table 1). Because a randomized experimental design for collection of the data was used to eliminate ageing effects, this alteration in DGC characteristics is best illustrated using one of the experiments in which O_2 concentration was systematically reduced (Fig. 2).

The characteristics of the DGC phases also changed considerably with declining ambient oxygen concentration. C




Fig. 3. The effects of ambient oxygen concentration on closed (C) phase (filled circles) and flutter (F) phase (open circles) rates of CO_2 release. Values are means \pm S.E.M. ($N = 42$). The regression lines are based on the individual data points: C phase CO_2 release rate = $0.0008[O_2] - 0.006$ ($F_{2,39} = 63.9$, $P = 0.00001$, $r^2 = 0.615$); F phase CO_2 release rate = $-0.0072[O_2] + 0.151$ ($F_{2,39} = 86.24$, $P = 0.00001$, $r^2 = 0.683$).




Fig. 4. The effects of ambient oxygen concentration on closed (C) phase (filled circles) and flutter (F) phase (open circles) duration. Values are means \pm S.E.M. ($N = 42$). The regression lines are based on the individual data points: C phase duration = $8.29[O_2] - 60.81$ ($F_{2,39} = 63.67$, $P = 0.00001$, $r^2 = 0.614$); F phase duration = $21.04[O_2] - 117.9$ ($F_{2,39} = 47.16$, $P = 0.0001$, $r^2 = 0.541$).

phase CO_2 release rate and duration declined significantly with declining oxygen concentration, and it appeared that this phase was eliminated entirely at oxygen concentrations less than 16% (Figs 2–4). F phase duration also decreased significantly with declining O_2 concentration (Fig. 4), although the rate of F phase CO_2 release showed the opposite trend (Fig. 3), and the proportion of the total rate of CO_2 release attributed to F phase CO_2 release increased significantly with declining oxygen concentration (Fig. 5). With declining O_2 concentration, the rate of O phase CO_2 release declined (O phase CO_2 release rate = $0.0066[O_2] + 0.186$, $F_{2,39} = 30.08$, $P = 0.00001$, $r^2 = 0.429$), as did O phase duration (O phase duration = $17.96[O_2] + 58.76$, $F_{2,39} = 59.55$, $P = 0.00001$, $r^2 = 0.598$) and O phase CO_2 emission volume (O phase CO_2 emission volume = $0.0018[O_2] - 0.0013$, $F_{2,39} = 87.96$, $P = 0.00001$, $r^2 = 0.687$).

During normoxia, cuticular water loss prior to death (estimated as 0.319 ± 0.069 mg, but with a small F phase component) contributed approximately 95% to the total water loss, whereas respiratory water loss (0.013 ± 0.003 mg) accounted for 5%. Overall water loss rates in normoxic moving air (1.328 mg h^{-1}) were considerably higher than those obtained using gravimetric protocols (0.633 mg h^{-1}).

Discussion

Assuming a respiratory quotient (RQ) of between 0.7 and 0.9, a value that may be reasonable for animals feeding on dung and its microflora (Landin, 1961, 1968; Holter, 1982; P. Holter, unpublished results), the standard metabolic rate (SMR) of




Fig. 5. The effects of ambient oxygen concentration on the percentage contribution of the flutter (F) phase rate of CO₂ release to the total rate of CO₂ release. Values are means \pm S.E.M. ($N=42$). The regression line is based on the individual data points: percentage F phase $\dot{V}_{\text{CO}_2} = -3.249[\text{O}_2] + 70.76$ ($F_{2,39} = 134.06$, $P = 0.00001$, $r^2 = 0.770$).

Aphodius fossor, estimated from the CO₂ release rate, is 124–140 μW at 15 °C. This is approximately 65 % lower than the values obtained by Holter and Spangenberg (1997) for the same species using different methods. Those authors did not control for the activity of the beetles, as we did here using flow-through methods and observations, and they conducted their study at a temperature 5 °C higher than used in the present study, which may account for the differences in the two estimates of SMR (see also Lighton and Fielden, 1995).

Nonetheless, in terms of maintenance of SMR, we found almost the same response of *A. fossor* to declining oxygen concentrations as was found by Holter and Spangenberg (1997). CO₂ release rate did not differ significantly with decreasing [O₂] to approximately 6 % O₂, and the increase in metabolic rate observed at approximately 3 % O₂ was small relative to the overall mean CO₂ release rate at all higher O₂ concentrations. The absence of a short-term response of metabolic rate to declining oxygen concentrations is not unusual in insects, which can generally maintain constant oxygen uptake rates down to at least 5 % O₂ (Schneiderman and Williams, 1955; Keister and Buck, 1974; Herreid, 1980; Loudon, 1988). At lower O₂ molar fractions, insects can be characterized either as conformers, which generally respond with a decline in metabolic rate (e.g. Loudon, 1988), or as regulators, which often increase their ventilation rate in response to declining P_{O_2} (Herreid, 1980). The regulatory response was shown by *A. fossor*; the increase in rate of CO₂ release was probably due to a small increase in the cost of convective ventilation caused by active muscular contractions at the lowest ambient oxygen concentration (see below).




Fig. 6. Typical recording of discontinuous gas exchange in an individual *Aphodius fossor*, desiccated to approximately half its maximum tolerable water loss, at 15 °C (mean $\dot{V}_{\text{CO}_2} = 0.231 \text{ ml g}^{-1} \text{ h}^{-1}$).

In response to declining oxygen concentrations, the duration of the C phase declined, and it appeared to be entirely absent at ambient O₂ concentrations below 10–16 % (Figs 2, 4). This provides further evidence in support of declining O₂ tension acting as the major trigger for the termination of this phase (Levy and Schneiderman, 1966; Lighton, 1996). However, it is of considerable interest that this probably represents an endotracheal P_{O_2} setpoint of approximately 10–15 kPa, although this was not measured in the current study. This is considerably higher than the widely accepted value of 5 kPa reported by Levy and Schneiderman (1966) (see also Lighton, 1994, 1996).

In contrast to Lighton and Garrigan's (1995) findings for the ant *Camponotus vicinus*, F phase duration in *A. fossor* declined significantly with declining ambient O₂ concentration (Fig. 4), although F phase CO₂ release (Fig. 3) and the proportional contribution of F phase CO₂ release rate to the overall rate of CO₂ release (Fig. 5) increased with declining oxygen concentration. Following Lighton and Garrigan's (1995) arguments (see above; see also Snyder et al., 1995), we suggest that in *A. fossor* the F phase was at least partially convective. Thus, as ambient oxygen concentrations decline, the absence of a significant negative endotracheal pressure must force premature termination of this phase, hence causing the observed increase in DGC frequency. The apparent absence of air sacs in this species (S. L. Chown and P. Holter, personal observations) and the curious semi-telescoping movements of the sternites and tergites that can be observed at very low oxygen concentrations when the elytra and wings are removed lend additional support to the suggestion that a significant negative endotracheal pressure cannot be produced. Indeed, it appears that, at the lowest oxygen concentrations, active ventilation using muscular contractions is required to ensure that metabolic demands for O₂ can be met.

Nonetheless, the increase in F phase rate of CO₂ release suggests that there was also a significant diffusive component to this phase. This increase in spiracular conductance is

probably brought about by an increase in the effective diameter of the spiracles (i.e. the extent to which they open, see Lighton and Garrigan, 1995), an assumption supported by the disappearance of the C phase with declining O₂ concentration. That is, at the lower oxygen concentrations, the spiracles do not close at all.

Therefore, it appears that, in response to declining ambient oxygen concentrations, *A. fossor* switches from a pattern of discontinuous gas exchange to one of continuous diffusion/convection (see also Schneiderman and Williams, 1955). CO₂ release rate during the O phase declines in importance as a consequence of declining O phase volume and duration, while the rate of F phase CO₂ release increases despite a declining F phase duration. In the light of these dramatic changes, the existence of an F phase in the conventional sense may well be disputed (Fig. 2). However, Lighton and Berrigan's (1995) proposition that, in response to a hypoxic/hypercapnic environment, insects can either open their spiracles maximally for long periods or sequester CO₂ and deplete O₂ until an adequate partial pressure gradient can be established is intriguing in this context. Indeed, it appears that the former response is exactly the one shown by *A. fossor*. Although it is likely to be associated with elevated water loss rates (Lighton et al., 1993b; Lighton and Berrigan, 1995), the penalty imposed on *A. fossor* by continuous convection and diffusion is unlikely to be high for two reasons. First, *A. fossor* loses the vast majority of water through cuticular transpiration and, by the standards of Zachariassen et al. (1987) it is a typically mesic species in this regard. Second, such high water loss rates, even if compounded by an increase in water loss associated with continuous ventilation, are unlikely to be significant in a species that spends the greater part of its adult and larval existence in a microhabitat with an 80% water content and that is characterized by high humidities and relatively low temperatures, at least when the adults are active (see above; see also Landin, 1961, 1968; Holter, 1982).

If conservation of water is relatively unimportant to adult *A. fossor*, if the DGC is not used to enhance gas exchange under hypoxic conditions (and presumably hypercapnic conditions because the two are usually encountered simultaneously in this species, see Holter, 1991, 1994), and if the DGC is eliminated in water-stressed adults (Fig. 6), then why should this species exhibit a DGC at all? Lighton and Berrigan (1995) suggested that a DGC may not be necessary under normoxic conditions in insects because partial pressure gradients are sufficient to allow both diffusive and convective gas exchange, and Quinlan and Hadley (1993) have demonstrated that, even if partially water-stressed, insects are likely to abandon the DGC because of the reduced CO₂ buffering capacity of the haemolymph. One explanation for the presence of the DGC in *A. fossor* may be that, during diapause or quiescence in adults, water conservation is likely to be at a premium. Although a sound argument, there is no evidence for a period of quiescence in the summer activity period of this species, and it is the larval stage that overwinters (Landin, 1961). An alternative explanation might be that at some point in the evolutionary

history of this species water conservation was indeed at a premium, and that the DGC has been retained as a plesiotypic character state (but see Coope, 1995, for an argument that beetles, including those in the genus *Aphodius*, tracked their favoured climates). Both scenarios are plausible, and both would lend support to the argument that the DGC is of adaptive value to terrestrial insects. However, both call for a *deus ex machina* that the available data prevent us from employing.

Thus, neither of the adaptive scenarios proposed for the genesis of the DGC appears to be suitable in the case of *A. fossor*, and general dissatisfaction with the water conservation hypothesis is also widespread (e.g. Hadley, 1994; Lighton, 1996; Williams and Bradley, 1998). Interpreted in the light of suggestions that the distribution and characteristics of the DGC are somewhat idiosyncratic with regard to taxa from mesic and xeric environments (Lighton, 1996), and indications that the DGC is likely to be polyphyletic (Lighton and Fielden, 1996), our findings suggest that the DGC may not be adaptive in a strict sense. Rather, the regular periodic nature of the DGC may be an epiphenomenon of the interaction of two feedback systems regulating a single function faced with minimal demand. Recently, it has been found that, in the absence of any deterministic driving variables, interacting feedback systems can show a variety of behaviours ranging from a single steady state to ordered cyclic behaviour (Kauffman, 1993). The periodic nature of a variety of physiological phenomena is thought to be the consequence of such an interaction (Glass and Mackey, 1988) but, when demand is placed on such a system, the emergent behaviour may give way to more tightly regulated function. This may be the case with the DGC, although admittedly we have little positive evidence to support this idea. However, given the difficulty of confirming adaptive explanations for the DGC, we suggest that this possibility be given consideration.

Annette Spangenberg is thanked for assistance in the laboratory, and Søren Christensen for setting his experiments aside while we used the gas mixer. Nigel Bennett, Melodie McGeoch, Adrian Davis and two anonymous referees are thanked for critical comments on an earlier version of the manuscript. The Carlsberg Foundation generously funded the research visit of S.L.C. to Denmark. S.L.C. is supported by the Foundation for Research Development and the University of Pretoria; P.H. is supported by the University of Copenhagen.

References

- Chown, S. L., Scholtz, C. H., Klok, C. J., Joubert, F. J. and Coles, K. S. (1995). Ecophysiology, range contraction and survival of a geographically restricted African dung beetle (Coleoptera: Scarabaeidae). *Funct. Ecol.* **9**, 30–39.
- Coope, G. R. (1995). Insect faunas in ice age environments: why so little extinction? In *Extinction Rates* (ed. J. H. Lawton and R. M. May), pp. 55–74. Oxford: Oxford University Press.
- Dellacasa, G. (1983). Sistemica e nomenclatura degli Aphodiini

- italiani (Coleoptera Scarabaeidae: Aphodiinae). *Mus. Reg. Sci. Nat. Monografie I*, 1–463.
- Endler, J. A.** (1986). *Natural Selection in the Wild*. Princeton: Princeton University Press.
- Glass, L. C. and Mackey, M. C.** (1988). *From Clocks to Chaos. The Rhythms of Life*. Princeton: Princeton University Press.
- Hadley, N. F.** (1994). *Water Relations of Terrestrial Arthropods*. San Diego: Academic Press.
- Hadley, N. F. and Quinlan, M. C.** (1993). Discontinuous carbon dioxide release in the eastern lubber grasshopper *Romalea guttata* and its effect on respiratory transpiration. *J. Exp. Biol.* **177**, 169–180.
- Herreid, C. F.** (1980). Hypoxia in invertebrates. *Comp. Biochem. Physiol.* **67A**, 311–320.
- Holter, P.** (1982). Resource utilization and local coexistence in a guild of scarabaeid dung beetles (*Aphodius* spp.). *Oikos* **39**, 213–227.
- Holter, P.** (1991). Concentration of oxygen, carbon dioxide and methane in the air within dung pats. *Pedobiologia* **35**, 381–386.
- Holter, P.** (1994). Tolerance of dung insects to low oxygen and high carbon dioxide concentrations. *Eur. J. Soil Biol.* **30**, 187–193.
- Holter, P. and Spangenberg, A.** (1997). Oxygen uptake in coprophilous beetles (*Aphodius*, *Geotrupes*, *Sphaeridium*) at low oxygen and high carbon dioxide concentrations. *Physiol. Ent.* **22**, 339–343.
- Kauffman, S. A.** (1993). *The Origins of Order. Self Organization and Selection in Evolution*. Oxford: Oxford University Press.
- Keister, M. and Buck, J.** (1974). Respiration: some exogenous and endogenous effects on rate of respiration. In *Physiology of Insecta*, vol. VI, 2nd edition (ed. M. Rockstein), pp. 469–509. New York: Academic Press.
- Kestler, P.** (1985). Respiration and respiratory water loss. In *Environmental Physiology and Biochemistry of Insects* (ed. K. H. Hoffmann), pp. 137–186. Berlin: Springer.
- Landin, B.-O.** (1961). Ecological studies on dung-beetles (Col. Scarabaeidae). *Opusc. Ent. Suppl.* **XIX**, 1–228.
- Landin, B.-O.** (1968). The diel flight activity of dung-beetles (Coleoptera Scarabaeidae). A study of the influence of environmental factors, with particular reference to the light. *Opusc. Ent. Suppl.* **XXXII**, 1–172.
- Levy, R. I. and Schneiderman, H. A.** (1966). Discontinuous respiration in insects. II. The direct measurement and significance of changes in tracheal gas composition during the respiratory cycle of silkworm pupae. *J. Insect Physiol.* **12**, 83–104.
- Lighton, J. R. B.** (1988). Simultaneous measurement of oxygen uptake and carbon dioxide emission during discontinuous ventilation in the tok-tok beetle, *Psammodes striatus*. *J. Insect Physiol.* **34**, 361–367.
- Lighton, J. R. B.** (1994). Discontinuous ventilation in terrestrial insects. *Physiol. Zool.* **67**, 142–162.
- Lighton, J. R. B.** (1996). Discontinuous gas exchange in insects. *Annu. Rev. Ent.* **41**, 309–324.
- Lighton, J. R. B. and Berrigan, D.** (1995). Questioning paradigms: caste-specific ventilation in harvester ants, *Messor pergandei* and *M. julianus* (Hymenoptera: Formicidae). *J. Exp. Biol.* **198**, 521–530.
- Lighton, J. R. B. and Fielden, L. J.** (1995). Mass scaling of standard metabolism in ticks: a valid case of low metabolic rates in sit-and-wait strategists. *Physiol. Zool.* **68**, 43–62.
- Lighton, J. R. B. and Fielden, L. J.** (1996). Gas exchange in wind spiders (Arachnida, Solpugidae): independent evolution of convergent control strategies in solpugids and insects. *J. Insect Physiol.* **42**, 347–357.
- Lighton, J. R. B., Fukushi, T. and Wehner, R.** (1993a). Ventilation in *Cataglyphis bicolor*: regulation of CO₂ release from the thoracic and abdominal spiracles. *J. Insect Physiol.* **39**, 687–699.
- Lighton, J. R. B. and Garrigan, D.** (1995). Ant breathing: testing regulation and mechanism hypotheses with hypoxia. *J. Exp. Biol.* **198**, 1613–1620.
- Lighton, J. R. B., Garrigan, D., Duncan, F. D. and Johnson, R. A.** (1993b). Spiracular control of respiratory water loss in female alates of the harvester ant, *Pogonomyrmex rugosus*. *J. Exp. Biol.* **179**, 233–244.
- Loudon, C.** (1988). Development of *Tenebrio molitor* in low oxygen levels. *J. Insect Physiol.* **34**, 97–103.
- Quinlan, M. C. and Hadley, N. F.** (1993). Gas exchange, ventilatory patterns and water loss in two lubber grasshoppers: Quantifying cuticular and respiratory transpiration. *Physiol. Zool.* **66**, 628–642.
- Schneiderman, H. A. and Williams, C. M.** (1955). An experimental analysis of the discontinuous respiration of the *Cecropia* silkworm. *Biol. Bull.* **109**, 123–143.
- Snyder, G. K., Sheafor, B., Scholnick, D. and Farrelly, C.** (1995). Gas exchange in the insect tracheal system. *J. Theor. Biol.* **172**, 199–207.
- Williams, A. E. and Bradley, T. J.** (1998). The effect of respiratory pattern on water loss in desiccation-resistant *Drosophila melanogaster*. *J. Exp. Biol.* **201**, 2953–2959.
- Zachariassen, K. E., Andersen, J., Maloiy, G. M. O. and Kamau, J. M. Z.** (1987). Transpiratory water loss and metabolism of beetles from arid areas in East Africa. *Comp. Biochem. Physiol.* **86A**, 403–408.