

COMMENTARY

A radical shift in perspective: mitochondria as regulators of reactive oxygen species

Daniel Munro^{1,2} and Jason R. Treberg^{1,2,3,*}

ABSTRACT

Mitochondria are widely recognized as a source of reactive oxygen species (ROS) in animal cells, where it is assumed that overproduction of ROS leads to an overwhelmed antioxidant system and oxidative stress. In this Commentary, we describe a more nuanced model of mitochondrial ROS metabolism, where integration of ROS production with consumption by the mitochondrial antioxidant pathways may lead to the regulation of ROS levels. Superoxide and hydrogen peroxide (H₂O₂) are the main ROS formed by mitochondria. However, superoxide, a free radical, is converted to the non-radical, membrane-permeant H₂O₂; consequently, ROS may readily cross cellular compartments. By combining measurements of production and consumption of H₂O₂, it can be shown that isolated mitochondria can intrinsically approach a steady-state concentration of H₂O₂ in the medium. The central hypothesis here is that mitochondria regulate the concentration of H₂O₂ to a value set by the balance between production and consumption. In this context, the consumers of ROS are not simply a passive safeguard against oxidative stress; instead, they control the established steady-state concentration of H₂O₂. By considering the response of rat skeletal muscle mitochondria to high levels of ADP, we demonstrate that H₂O₂ production by mitochondria is far more sensitive to changes in mitochondrial energetics than is H₂O₂ consumption; this concept is further extended to evaluate how the muscle mitochondrial H₂O₂ balance should respond to changes in aerobic work load. We conclude by considering how differences in the ROS consumption pathways may lead to important distinctions amongst tissues, along with briefly examining implications for differing levels of activity, temperature change and metabolic depression.

KEY WORDS: Hydrogen peroxide, Energetics, Antioxidant, Skeletal muscle, Oxidative stress, Glutathione peroxidase, Thioredoxin, Peroxiredoxin

Introduction

Over 40 years ago (Jensen, 1966; Loschen et al., 1971; Boveris et al., 1972; Boveris and Chance, 1973) it was discovered that mitochondria can produce reactive oxygen species (ROS; see Glossary). Since then, the realization that mitochondria may be a source of potentially harmful ROS has profoundly influenced the perceived role(s) of mitochondria in cellular function. The proposed physiological effects of ROS include life-history and energetic trade-offs, cellular dysfunction and damage in response to intense or

prolonged activity, and even senescence and ageing (Trushina and McMurray, 2007; Costantini, 2008; Monaghan et al., 2009; Dai et al., 2014; Day, 2014; Mason and Wadley, 2014). For example, during migration, a trade-off may occur between individual performance and survival or reproductive fitness, owing to oxidative damage accrued during intense exercise. Publications in these areas, and many others, often describe mitochondria as the ‘major source’ of ROS in animal cells, despite the high antioxidant capacity of mitochondria (Zoccorato et al., 2004; Dreschel and Patel, 2010; Banh and Treberg, 2013). However, the rationale behind the notion of mitochondria as the major source of ROS in animal cells has been questioned (Brown and Borutaite, 2012), if not directly challenged, based on the capacity of isolated mitochondria to consume substantial amounts of ROS (Zoccorato et al., 2004). It is worth noting that it can be challenging to test hypotheses on mitochondrial function using cells, whole tissue or organism systems because of the additional control exerted by the plasma membrane. For this reason, isolated mitochondria are a valuable tool in understanding how mitochondria may respond to changing cellular conditions (which can be simulated by manipulation of the assay medium). Findings can then inform the generation of hypotheses applicable to higher levels of biological organization.

Recently, a different and more nuanced perspective on the role of mitochondria in ROS balance has been emerging: the contention that the ROS-producing and antioxidant-mediated ROS-consuming pathways of the mitochondrion can be integrated into a regulatory system. This perspective views ROS as specific regulatory molecules, consistent with their role in signalling, rather than simply as inevitable toxic by-products of aerobic metabolism, although excess ROS will cause oxidative stress. To our knowledge, the formalization of these ideas is rooted in work by Andreyev and colleagues (reviewed in Andreyev et al., 2005), initially laid out as a hypothesis by Starkov (2008). Recent experimental evidence in rodent brain and skeletal muscle mitochondria supports this hypothesis (Starkov et al., 2014; Treberg et al., 2015).

This Commentary will focus on the skeletal muscle system, and aims to illustrate how mitochondria can regulate ROS levels. After briefly addressing several concepts on the interplay between mitochondrial energetics (see Glossary) and the major pathways of mitochondrial ROS metabolism, we will illustrate how mitochondria can act as regulators of an important ROS, hydrogen peroxide (H₂O₂), *in vitro*. Using the proposed model of mitochondria as regulators of H₂O₂, the impact of high ADP availability on mitochondrial H₂O₂ metabolism will be explored, followed by some consideration of how differing degrees of physical activity should affect the regulation of H₂O₂ by skeletal muscle mitochondria. Finally, we consider how temperature may influence mitochondrial H₂O₂ metabolism, the impact our model could have on metabolic depression during torpor and how the relevant processes may vary across tissues.

¹Department of Biological Sciences, University of Manitoba, Winnipeg, MB, Canada R3T 2N2. ²Centre on Aging, University of Manitoba, Winnipeg, MB, Canada R3T 2N2. ³Department of Human Nutritional Sciences, University of Manitoba, Winnipeg, MB, Canada R3T 2N2.

*Author for correspondence (jason.treberg@umanitoba.ca)

 J.R.T., 0000-0001-5112-7325

List of symbols and abbreviations

GSH	reduced form of glutathione
GSSG	oxidized form of glutathione also known as glutathione disulphide
$[H_2O_2]_{ss}$	steady-state concentration of H_2O_2
k	rate constant
PMF	protonmotive force
%NAD(P)H	percent of combined NAD and NADP pools that are in the reduced form
V_c	rate of H_2O_2 consumption
$V_{c,app}$	apparent (measurable) rate of H_2O_2 consumption
V_p	rate of H_2O_2 production
$V_{p,app}$	apparent (measurable) rate of H_2O_2 production

Reactive oxygen species are metabolites too

Mitochondria are well known for their role in the aerobic supply of ATP by oxidative phosphorylation (Box 1), a process often purported to inevitably result in at least some ROS production. There are many sites for mitochondrial ROS production (simplified in Fig. 1A), which predominantly proceeds by the escape of an electron from a redox centre on an enzyme (or a subunit of an enzyme complex) onto oxygen (Fig. 1B). Thus, electrons indeed 'leak' from their usual route through the mitochondrial electron transport system (ETS) and Krebs cycle to form ROS. The superoxide radical is the main ROS formed at most sites of production. This free radical (see Glossary) is rapidly dismutated to the non-radical ROS H_2O_2 , which is membrane permeant and can thus diffuse between cellular compartments. The concentration of H_2O_2 in the mitochondrion (Cochemé et al., 2011) and in the cytosol (Arniaz et al., 1995; Palomero et al., 2008) appears to be maintained in the submicromolar range by antioxidants, which probably limits the formation of the membrane-permeant and highly reactive hydroxyl radical (Fig. 1B,C).

Recent studies have identified site-specific inhibitors of ROS production – at least for the outer ubiquinone-binding site in complex III – which have minimal effects on mitochondrial energetics (Orr et al., 2015). The potential for ROS production to cause damage to cells is nearly universal in aerobes – in light of the pharmacological effects discussed above, this raises the question of whether there might be evolutionary trade-offs between energy transformation in aerobic metabolism and ROS production. In other words, if ROS are universally damaging, then why has their production not been mitigated by natural selection to be effectively negligible?

The answer may lie in the now-recognized signalling role of ROS, which are often involved as second messengers in signalling cascades controlling, for example, apoptosis, cell differentiation, wound healing and cell-shape changes, and which also play a role in insulin signalling and redox signalling (see Glossary) in tumour cells (reviewed in Sies, 2014). Mitochondrial ROS in particular have been suggested to mediate feedback signalling to the nucleus and mitochondrial transcription machinery in order to adjust oxidative phosphorylation capacity through mitochondrial biogenesis and modulation of the activity of Krebs cycle enzymes (Moreno-Loshuertos et al., 2006). Mitochondrial ROS also act as regulators of mitophagy (Scherz-Shouval and Elazar, 2011). Recently, it has also been proposed that energy metabolism-linked mitochondrial redox signals may be indirectly mediated via the influence of H_2O_2 on glutathione (GSH) status – the ratio of GSH to its oxidized form, GSSG, is modulated in response to H_2O_2 concentration. This influence on the GSH pool alters the activity of proteins both inside and outside the mitochondrion through S-

Glossary**First-order kinetics**

Description of a reaction rate that is dependent on the concentration of one reactant and can be summarized by the following rate equation: rate of reaction= $k[A]$, where A is the reactant that on which the rate is dependent and k is the rate constant. First-order reactions follow an exponential decay curve for A over time, which can be described by $A_t=A_0 \cdot e^{-kt}$, where A_0 is the initial amount of A, k is the same reaction constant already described and t is time.

Free radical

An atom or molecule that has an unpaired electron, designated by the raised dot as seen with the superoxide anion ($O_2^{\cdot-}$), which is formed by the univalent (single electron) reduction of dioxygen (O_2).

Mitochondrial energetics

The interacting processes of electron and proton flux in the mitochondrion that link substrate oxidation to energy transformation. Major components include the membrane potential ($\Delta\Psi$) and the pH gradient (ΔpH), which make up the protonmotive force (PMF), as well as the rates of reactions and status of the involved metabolic intermediates.

Mitochondrial uncoupling

Process in which the protonmotive force established by the electron transport system is dissipated by proton leak across the mitochondrial inner membrane without being coupled to ATP synthesis.

%NAD(P)H

A measurement of the relative reduction status of the combined nicotinamide cofactor [NAD(H) and NADP(H)] pools. Generally measured by autofluorescence because the reduced form is strongly fluorescent while the oxidized form is not. For isolated mitochondria this is influenced by both cofactor pools, but is biased toward reflecting the NAD(H) pool.

Protonmotive force (PMF)

The potential energy established by the electrochemical disequilibrium created by proton translocation. It includes an electrical component, the membrane potential ($\Delta\Psi$) and a concentration gradient (ΔpH). The protonmotive force is used to couple oxidation of substrate to phosphorylation of ADP to ATP, which together are referred to as 'oxidative phosphorylation'.

Reactive oxygen species (ROS)

Molecules that contain oxygen and are chemically reactive, meaning they readily react with other molecules without the need for a catalyst. Examples include free radicals [superoxide anion ($O_2^{\cdot-}$) and hydroxyl radicals ($\cdot OH$)] and nonradicals [hydrogen peroxide (H_2O_2) and other peroxides].

Redox signal


Cellular signalling mechanism based on redox chemistry. Generally, reactive oxygen or nitrogen species or other chemical compounds are major messengers and the relative reduced to oxidized ratio of one or more major cellular redox couple(s) can be a central component (e.g. the NADH/NAD⁺) of the signal.

glutathionylation – a reversible post-translational modification compatible with a signalling role (Mailloux and Treberg, 2016). Therefore, ROS (or H_2O_2 at least) play roles in cell function that may require further processing by specific enzymes. Thus, H_2O_2 may be better viewed as a bona fide metabolite rather than simply an inevitable or unavoidable 'by-product' of aerobic metabolism.

Energetics and mitochondrial H_2O_2 metabolism

In this Commentary, our central thesis is that mitochondria can differentially regulate the concentration of H_2O_2 (through observed changes in its production and consumption) based on their current energetic state. Because electron flow through the mitochondrial ETS is coupled to proton translocation, the flux of electrons through to oxygen is impeded as the protonmotive force (PMF; see Glossary) increases. This impedance leads to a buildup of electrons in redox centres within the enzyme complexes that are

Box 1. The electron transport system of the mitochondrion



The electron transport system (ETS) of the mitochondrion plays a central role in oxidative phosphorylation by building the protonmotive force required for regenerating ADP into ATP. The ETS consists of four multisubunit enzyme complexes (CI to CIV) that span or are embedded in the surface of the inner mitochondrial membrane. In coordination with smaller mobile electron carrier proteins (ubiquinone, Q; and cytochrome c, C), these complexes channel the passage of substrate-derived electrons towards oxygen, the final acceptor. Electrons enter the ETS via complex I when originating from the oxidation of NADH-generating substrates, such as in the Krebs cycle. Electrons can also enter the ETS at the level of complex II, and other flavin adenine dinucleotide-containing enzymes (omitted for clarity). Inside the ETS, a net flux towards oxygen reduction is established by favourable electron transfers between redox couples of increasing electron affinity. This cascading of electrons towards a lower level of binding energy is harnessed at the level of complexes I, III and IV, and used to drive charges against their gradient and create the protonmotive force. More precisely, these three complexes couple the passage of electrons with the translocation of protons (H⁺) from the matrix side to the intermembrane space. A fifth complex (complex V, also known as F₁F₀ ATPase) is not involved directly with electron flux; instead, it harnesses the protonmotive force created by the ETS to drive the regeneration of ADP into ATP. Specifically, and under normal conditions, the channelling of H⁺ through complex V, with the direction of the electrochemical gradient (favourable direction), provides the energy for phosphorylation of ADP.

the source of superoxide or H₂O₂ (Fig. 1) and in the metabolic intermediates linking these enzymes. In other words, the mitochondrial redox centres and electron carriers will be in a more reduced state (higher electron availability) as the PMF increases. This increases ROS production and is central to the ‘uncoupling to survive’ hypothesis (Brand, 2000), which postulates that one of the selective advantages to mitochondrial uncoupling (see Glossary) is to prevent excessive ROS production, thereby limiting the oxidative damage that may lead to cellular senescence.

Animal mitochondria contain three major H₂O₂-consuming enzymes or enzymatic pathways. Catalase is an H₂O₂-consuming enzyme that does not require an external supply of electrons to convert H₂O₂ into H₂O and O₂, making it independent of the mitochondrial energetic state (Fig. 1C). Two respiration-dependent pathways also neutralize H₂O₂: the glutathione-dependent and the thioredoxin-dependent pathways (Fig. 1C). These pathways rely on a constant supply of electrons, which are supplied via NADPH. Oxidized NADP⁺ is reduced by the oxidation of respiratory substrates, leading to the maintained supply of reduced intermediates (GSH or thioredoxin), which are required for the activity of the peroxidases (e.g. peroxiredoxin-3 and glutathione peroxidase; Fig. 1C).

To summarize, oxidation of respiratory substrates contributes to the establishment of the PMF. The PMF will feed back on the flux of electrons through the ETS and the Krebs cycle that feeds into the ETS, leading to the accumulation of electrons in the enzymes and mitochondrial electron carriers, and thus influencing the reduction status of the NAD⁺ and NADP⁺ [i.e. NAD(P)⁺] pool in the mitochondrial matrix. Thus, substrate oxidation will prime both the ROS-producing sites and the respiration-dependent pathways of H₂O₂ consumption. Given that different respiratory substrates are not oxidized at equal rates, nor do they produce equal amounts of ROS, there can be a wide range of ROS production rates and NAD(P)H availability, depending on the substrate supply.

Fig. 1D shows the rate of H₂O₂ production and H₂O₂ consumption for a range of substrates. From these data, it is clear that H₂O₂ consumption capacity largely surpasses H₂O₂ production across a range of substrate conditions in muscle mitochondria. The figure also suggests that high and constant H₂O₂ consumption capacities are maintained across various respiratory substrates, and that any apparent decrease in H₂O₂ consumption is thus the result of an increase in production.

Regulation of metabolite concentration

In the simplest terms, the concentration of a metabolite is set by the interaction between the pathways for its formation and removal. Within a given system, the ability for regulation of a metabolite requires that the consumption or clearance capacity must be greater than the rate of production. This is, in fact, the case for H₂O₂ production for many respiratory substrates in rodent skeletal muscle (Fig. 1D; Treberg et al., 2015; Munro et al., 2016) and brain mitochondria (Starkov et al., 2014). An additional condition promoting metabolic regulation is that consumption of the metabolite should be able to respond to changes in production. Starkov et al. (2014) proposed that the consumption of H₂O₂ by mitochondria follows first-order kinetics (see Glossary), where the rate of disappearance is proportional to the [H₂O₂] and is described by a single rate constant (*k*), a simple means of satisfying this second requirement. Eqn 1 describes the relationship proposed by Starkov and colleagues, wherein for any rate of H₂O₂ production (*V_p*), the steady-state concentration of the metabolite ([H₂O₂]_{ss}) that will be achieved can be related to the first-order rate constant for consumption (*k*) as follows:

$$[\text{H}_2\text{O}_2]_{\text{ss}} = V_p/k. \quad (1)$$

This model predicts that when the concentration of the metabolite of interest starts at zero, then initially the rate of appearance or production (solid black line in Fig. 2A) will approximate the increase in metabolite concentration over time (dotted line in Fig. 2A). However, as the concentration of the metabolite increases, the rate of consumption (solid grey line in Fig. 2A) will also increase. As the system proceeds, the rate of consumption approaches the rate of production, and the metabolite concentration will approach a stable value ([H₂O₂]_{ss}). The system will tend towards this set-point regardless of whether the concentration of the metabolite starts at zero.

Using this simple model, we can explore what happens when parts of the system change. For example, if production (*V_p*) increases, but the first-order constant (*k*) for consumption is maintained, the resulting [H₂O₂]_{ss} will increase (Fig. 2B, left). Conversely, if *V_p* is maintained constant, but *k* is increased, then the [H₂O₂]_{ss} will decrease (Fig. 2B, centre). An important aspect of this model is that the speed at which the system approaches equilibrium increases with increasing values of *k*.






Fig. 1. Mitochondrial metabolism of H₂O₂. (A) Major sites of superoxide and H₂O₂ generation in enzyme complexes of the electron transport system (ETS) and the Krebs cycle (reviewed in Brand, 2010, 2016). CI–CIV, complexes I–IV; PDH, pyruvate dehydrogenase; αKGDH, alpha-ketoglutarate dehydrogenase; ETF:QOR, electron-transferring flavoprotein; ubiquinone-oxidoreductase; Q, ubiquinone; C, cytochrome c. Note: several additional sites and metabolic intermediates have been omitted for clarity. (B) Generalized process of electron leaks leading to reactive oxygen species (ROS) production. Superoxide is produced by a single electron reduction of dioxygen, and is detoxified into H₂O₂ by superoxide dismutase (SOD); H₂O₂ can form the hydroxyl radical (•OH) and the hydroxyl anion (OH⁻) via the non-enzymatic Fenton reaction (rxn) in the presence of certain transition metals. (C) Mitochondrial H₂O₂ consumption. Unlike catalase, the thioredoxin (Trx)- and glutathione (GSH)-dependent pathways for H₂O₂ consumption require reducing equivalents from NADPH. These respiration-dependent pathways rely on regeneration of Trx and GSH via conversion of an oxidized (S–S) and to reduced (–SH) form thiol intermediate between the NADPH-dependent reductase and the H₂O₂-consuming peroxidase. (D) Rates of H₂O₂ metabolism by isolated rat skeletal muscle mitochondria (data are expressed per mg of mitochondrial protein; from Munro et al., 2016). Note that the pathways for the consumption of H₂O₂ have been pharmacologically compromised in the production assay to minimize underestimation of the production rate. For the consumption assay, mitochondria are incubated with 2.5 μmol l⁻¹ H₂O₂ and allowed to simultaneously consume and produce H₂O₂ for 10 min. Very low consumption of H₂O₂ is observed in the absence of respiratory substrate, which suggests limited involvement of catalase. Mal, malate; Glu, glutamate; Succ, succinate; Rot, rotenone. Results are means ± s.e.m. (n=6).




Fig. 2. Illustration of the model for mitochondrial H₂O₂ regulation based on first-order kinetics of H₂O₂ consumption. (A) A simulation of increasing H₂O₂ consumption leading to stabilization of [H₂O₂] as time proceeds at a given production rate (V_p). The equation for calculating [H₂O₂] (dotted line) is given to the right, and cumulative (cumul.) H₂O₂ production and consumption are plotted as solid black and grey lines, respectively. The slope of the cumulative production and consumption lines indicates the rate of these processes. As the [H₂O₂] increases, the consumption increases and the [H₂O₂] approaches a steady-state ([H₂O₂]_{ss}, indicated by the dashed line). (B) Simulation of the effects of varying V_p and the consumption rate constant (k). See 'Regulation of metabolite concentration' for further details. (C) The fit of a first-order equation to the removal of H₂O₂ from the medium by respiring mitochondria. Data were obtained using a non-destructive probe for monitoring [H₂O₂]. The figure is from Treberg et al. (2015) (*Redox Biol.* 5, 216–224; copyright © 2015 the authors; published by Elsevier B.V.). Dark continuous lines indicate the measured [H₂O₂], with the s.e.m. shown by the surrounding light grey lines. The dotted lines indicate the average fit for the removal of H₂O₂, assuming a first-order reaction. Note that reducing k by addition of auranofin markedly alters the rate of H₂O₂ removal but that there is still a reasonable agreement between the modelled and measured data. (D) The apparent rate of H₂O₂ consumption is linearly related to the apparent rate of production across values obtained with multiple respiratory substrates (data are expressed per mg of mitochondrial protein). The actual rate of consumption can be approximated by extending the linear regression to the y-intercept where the influence of production will be negligible. See 'Differentiating between apparent and actual rates of H₂O₂ production and consumption' for the difference between apparent and true rates of production and consumption. Data are from Munro et al. (2016). Each symbol represents a different condition of respiratory substrate oxidation. Black symbols and dashed line, control mitochondria; grey symbols and dotted line, addition of auranofin. Results are means ± s.e.m. ($n=4-6$), and the linear regression is shown by the dashed and dotted lines.

This is illustrated in Fig. 2B (right panel) where both production and consumption have been doubled as compared with panel A. It can be seen that the [H₂O₂]_{ss} is set by V_p/k , which is the same as in panel A, but is reached much faster with the higher value of k .

First-order kinetics of respiration-dependent H₂O₂ consumption pathways?

Our general model (Fig. 2A,B) uses an assumption of first-order kinetics to describe the relationship between [H₂O₂] and the

capacity of mitochondria to consume this ROS, because the rate of consumption (V_c) is a function of [H₂O₂]. We have previously shown that the decay of extramitochondrial H₂O₂ fits reasonably with first-order kinetics when considering energized rat skeletal muscle mitochondria (Fig. 2C). However, the assumption of a simple first-order reaction describing the combined consumers of ROS allows for consumption to be infinitely high at infinite H₂O₂ concentration. An infinite value of V_c is, of course, unrealistic; the ROS consumers do appear to approach saturation (Banh

and Treberg, 2013; Munro et al., 2016). To clarify, all of the measurements of H_2O_2 consumption rate (V_c) used here so far (Figs 1D, 2D) have an initial, approximately linear phase at high $[\text{H}_2\text{O}_2]$ from which we could calculate the rate of H_2O_2 consumption in terms of nmol min^{-1} . Calculating linear rates provides a convenient means of comparing rates of H_2O_2 consumption (V_c) with rates of production (V_p), but the first-order kinetic approximation remains a useful means of describing the response of the consumers under conditions when $[\text{H}_2\text{O}_2]$ is low and more likely to be similar to physiological levels.

At the crossroads: producers and consumers interact with a common matrix pool of H_2O_2

Much of the experimental evidence for mitochondria as regulators of $[\text{H}_2\text{O}_2]$ comes from assays monitoring extramitochondrial H_2O_2 ; this is because components of the assay system that is used to quantify H_2O_2 cannot cross biological membranes and are confined to the medium. It is conventionally assumed that only minor amounts of H_2O_2 produced in the mitochondrial matrix are consumed before reaching the extramitochondrial detection system. However, using rodent skeletal muscle mitochondria, we recently demonstrated that this is not the case (Treberg et al., 2010, 2015; Munro et al., 2016). This means that mitochondrial ROS production rates are largely underestimated when based on extramitochondrial detection of H_2O_2 . This is an important additional complexity to consider when evaluating the model leading to Eqn 1.

Differentiating between apparent and actual rates of H_2O_2 production and consumption

The interaction between endogenous mitochondrial H_2O_2 metabolism and observed rates of change in extra-mitochondrial H_2O_2 is a clear demonstration that H_2O_2 is exchanged between the mitochondrial matrix and extramitochondrial pools. This means that there is a distinction between ‘actual’ rates of production and consumption of H_2O_2 (V_p and V_c , respectively), and their ‘apparent’ rates (those which are measured), $V_{p,\text{app}}$ and $V_{c,\text{app}}$. Appreciating this distinction led us to derive a means of estimating the actual values of V_p and V_c , by measuring their apparent values and extrapolating to where the rate of the competing intramitochondrial process would be zero. For example, Fig. 2D shows how $V_{c,\text{app}}$ relates to $V_{p,\text{app}}$. As $V_{c,\text{app}}$ declines, the rate of H_2O_2 production increases, which we interpret as competition between exogenous and endogenous supply of H_2O_2 to the consumption pathways. However, by extending the relationship to the y -intercept, an estimate of the actual rate of consumption of H_2O_2 can be made where the influence of H_2O_2 production will be zero (Fig. 2D). If the matrix H_2O_2 consumption capacity is inhibited with auranofin, which impairs the thioredoxin-dependent pathway (Munro et al., 2016), the slope of the relationship between $V_{p,\text{app}}$ and $V_{c,\text{app}}$ decreases, and the predicted rate of H_2O_2 consumption, in the absence of competition from endogenous production, also decreases.

Importantly, the strong correlation between $V_{p,\text{app}}$ and $V_{c,\text{app}}$ leads to the interpretation that the consumption rate (V_c) is maintained across different substrate conditions. In other words, the supply of NADPH to the reductases for consumption of H_2O_2 is not sufficiently low to limit the rate of H_2O_2 consumption by the peroxidases, at least for the set of substrate conditions that we tested. This satisfies the requirement of the model leading to Eqn 1 for greater overall capacity for H_2O_2 consumption than production across a large range of energetic conditions, with the added benefit that it is reasonable to assert that the capacity of the H_2O_2 -consuming pathways is consistent across these substrate conditions.

Direct evidence of mitochondrial H_2O_2 regulation

As we have shown above, skeletal muscle mitochondria have the requisite traits described in our general model of metabolite regulation that led to Eqn 1. For the interested reader, more detailed explanations can be found elsewhere (see Starkov et al., 2014; Treberg et al., 2015), but the requirements include $[\text{H}_2\text{O}_2]$ -dependent consumption that can act on both intramitochondrial and extramitochondrial H_2O_2 and outpace production. Combining these concepts, we will now discuss how isolated mitochondria can act as regulators of $[\text{H}_2\text{O}_2]$.

Our model predicts how the system can regulate towards a $[\text{H}_2\text{O}_2]_{\text{ss}}$. This $[\text{H}_2\text{O}_2]_{\text{ss}}$ should be a function of V_p , set by the substrates added, and V_c described by k , which does not change across substrate conditions (Fig. 2D). In order to determine whether H_2O_2 is accumulating in the medium or being consumed by the mitochondria, we used a modified version of the horseradish peroxidase-linked fluorometric Amplex UltraRed assay. Normally, using this assay to measure H_2O_2 that has escaped from the mitochondrion consumes all extramitochondrial H_2O_2 , thus maintaining an outward diffusion gradient (Fig. 3A). We modified the assay by adding respiratory substrate as normal but withholding the Amplex UltraRed for a period of time. This allows assessment of whether H_2O_2 has accumulated or been consumed – an accumulation of H_2O_2 is measured as a ‘jump’ in the fluorescence following addition of Amplex UltraRed (Fig. 3B). For example, while using glutamate and malate as the respiratory substrates, we see negligible accumulation of H_2O_2 in the presence of energized mitochondria, in accordance with low V_p but high and maintained V_c (Fig. 3C). In this example, it is clear that some H_2O_2 escapes the mitochondrion and can be detected when the detection system is completed by the addition of Amplex UltraRed at different time points for the four parallel reactions (Fig. 3). But there is no detectable accumulation of H_2O_2 in absence of a complete detection system, suggesting the consumers are maintaining H_2O_2 at very low concentrations. However, compromising V_c with the inhibitor auranofin, which decreases k (Treberg et al., 2015), and adding the same substrates leads to the appearance of H_2O_2 in the medium before the addition of Amplex UltraRed (Fig. 3D). If V_p is high enough for a given substrate, we would expect that some H_2O_2 should accumulate in the medium in absence of auranofin. Succinate leads to very high H_2O_2 production rates in well-coupled muscle mitochondria (Fig. 1D). When using succinate as a respiratory substrate, H_2O_2 accumulates (Fig. 3E) and even approaches a steady-state concentration over time (Fig. 3F), as predicted. These results are consistent with the general model in Fig. 2, and we conclude that for any condition where the consumption capacity does not change, any increase or decrease in the rate of H_2O_2 production should lead to a concomitant increase or decrease in the established $[\text{H}_2\text{O}_2]_{\text{ss}}$.

How should $[\text{H}_2\text{O}_2]_{\text{ss}}$ respond to changes in mitochondrial energetics? ADP as a test case

ADP activates the mitochondrial ATP synthase, thus allowing protons to return to the matrix side of the inner mitochondrial membrane, and partially collapsing the PMF (Box 1). This reduction in the PMF allows for increased flux through PMF-generating complexes in the ETS of the inner membrane, thus decreasing the concentration of the reduced form of NAD^+ and NADP^+ [NAD(P)H] in the matrix. Here, we illustrate these effects under two different substrate conditions: (1) malate (alone), which is a poorly oxidized respiratory substrate in muscle mitochondria, and (2) malate with glutamate, which allows for more rapid oxidation of




Fig. 3. Demonstrating the accumulation of H₂O₂ over time with isolated muscle mitochondria. (A) (Upper) Cartoon demonstrating the principle of the fluorometric detection of extramitochondrial H₂O₂ via the H₂O₂-consuming horseradish peroxidase (HPx)-linked enzymatic reaction. By eliminating all extramitochondrial H₂O₂, the assay maintains a strong outward diffusion gradient for H₂O₂ that can be followed over time as a measure of net H₂O₂ escape from the mitochondrion (lower). (B) (Upper) Illustration of conditions when the H₂O₂ detection system is incomplete because of absence of the fluorometric substrate. (Lower) Schematic showing how to use the jump in fluorescence, relative to a cuvette containing complete detection system at t=0, to determine extramitochondrial H₂O₂ accumulation in the medium. (C–E) All assay components, except the molecular fluorescent probe Amplex UltraRed (Amplex), were added before beginning the assay. Additions of Amplex UltraRed are indicated by vertical grey arrows. (C) There is negligible accumulation of H₂O₂ in the medium of mitochondria respiring on 5 mmol l⁻¹ glutamate and malate. (D) Addition of 2 μmol l⁻¹ auranofin to mitochondria respiring on 5 mmol l⁻¹ glutamate and malate results in H₂O₂ accumulation. (E) Mitochondria respiring on 5 mmol l⁻¹ succinate accumulate H₂O₂ in the medium in a time-dependent manner. Note that where the traces reach 1000 arbitrary units, the detector has reached its maximum for the sensitivity settings used. (F) Accumulation of H₂O₂ in the medium from mitochondria respiring on 5 mmol l⁻¹ succinate approaches a stable steady-state concentration as determined by Eqn 1 and based on experiments as illustrated in E. Data are means ± s.e.m. (n=3). k and V_p values are expressed per g of mitochondrial protein. C–E and legend are modified from Treberg et al. (2015) (*Redox Biol.* **5**, 216–224; copyright © 2015 the authors; published by Elsevier B.V.). For the interested reader, values for kinetic parameters and explanation of units can be found elsewhere (Treberg et al., 2015).




Fig. 4. The effects of mitochondrial energetics on apparent rates of H_2O_2 production ($V_{p,app}$) and consumption ($V_{c,app}$). (A) In isolated rat skeletal muscle mitochondria, the protonmotive force (PMF) and the matrix nicotinamide cofactor reduction state [%NAD(P)H] are correlated across different conditions of respiratory substrates. (B) Addition of ADP ($500 \mu\text{mol l}^{-1}$) decreases the PMF and $V_{p,app}$, but the effects on $V_{c,app}$ are limited ($V_{p,app}$ and $V_{c,app}$ values are expressed per mg of mitochondrial protein). (C) Addition of ADP decreases the %NAD(P)H and $V_{p,app}$, but the effects on $V_{c,app}$ are limited ($V_{p,app}$ and $V_{c,app}$ values are expressed per mg of mitochondrial protein). The upper right partial x-axis indicates the %NAD(P)H values associated with three levels of physical activity (rest, mild aerobic and intense aerobic exercise) in rat skeletal muscle mitochondria, according to Gonçalves et al. (2015). Data for A–C are from Quinlan et al. (2012) for PMF and %NAD(P)H or from unpublished observations (D.M. and J.R.T.; $n=4$; based on protocols described in Munro et al., 2016) for $V_{p,app}$ and $V_{c,app}$. All values are means \pm s.e.m. Mal, malate; Glu, glutamate.

respiratory substrates and a shift towards a more reduced state in the nicotinamide cofactor pools – that is, an increase in NADH/NAD⁺ and NADPH/NADP⁺ – as indicated by higher %NAD(P)H (see Glossary).

As expected, respiration with glutamate and malate leads to a higher PMF than with malate alone, and under both substrate conditions the presence of ADP markedly decreases the PMF (Fig. 4A) (data from Quinlan et al., 2012). Similarly, glutamate and malate in combination allow for a higher %NAD(P)H as compared with malate alone, and ADP markedly decreases %NAD(P)H for both substrate conditions. Note that measurement of PMF under the same experimental conditions demonstrates a strong relationship between PMF and the %NAD(P)H (Fig. 4A). The strength of the relationship allows us to exploit the effects of ADP on the mitochondrial energetic state in order to investigate the relationship between PMF and the production and consumption of H_2O_2 .

As the PMF decreases, in response to ADP, the capacity to consume H_2O_2 shows relatively little response compared with the rate of ROS production, which drops by a factor of three to four (Fig. 4B). Similarly, large decreases in %NAD(P)H in response to ADP addition largely affect H_2O_2 production rates while having relatively little impact on the capacity for consumption (Fig. 4C). This is typical, in our experience, of skeletal muscle mitochondria respiring under conditions that can readily reduce NAD⁺ to NADH either by matrix dehydrogenase reactions and buildup of Krebs cycle intermediates or by the reversal of complex I of the ETS (J.R.T., S. Banh, P. Zacharias, L. Wiens, D.M., unpublished observations).

Preferential activation of the consumption pathways over the production pathways at low %NAD(P)H

The respiration-dependent H_2O_2 consumption pathways characteristic of muscle mitochondria require a constant supply of NADPH to maintain flux and should theoretically fail under low %NAD(P)H (Fig. 1C). Above, we show that, under two different substrate conditions, and with or without ADP, the decrease in H_2O_2 production is much more pronounced than the decrease in H_2O_2 consumption when %NAD(P)H declines, indicating a more oxidized state of the matrix nicotinamide cofactor pool. In other words, the consumers appear to be primed and ready to quench H_2O_2 well in advance of reaching energetic states that are associated with high rates of ROS production. This has implications for the signalling role of H_2O_2 in transmitting information about the energetic status of the mitochondrion to its transcriptional machinery and to the nucleus (Moreno-Loshuertos et al., 2006). The resilience of the consumers in the face of declining %NAD(P)H, relative to the producers, leads to a system where – under most conditions – the $[\text{H}_2\text{O}_2]_{ss}$ will reflect the changes in H_2O_2 production (Eqn 1), which is largely a function of the mitochondrial energetic state.

How the mitochondrion interacts with other cellular elements in ROS handling is currently unknown. Some cytoplasmic components (homologs of mitochondrial H_2O_2 consumption and production pathways) also have the potential to regulate $[\text{H}_2\text{O}_2]_{ss}$ in a manner similar to what we propose for mitochondria. However, these cytosolic consumers and extramitochondrial sources of H_2O_2 could have different $[\text{H}_2\text{O}_2]_{ss}$ set-points compared with that of the mitochondrial matrix. Mitochondria may therefore represent a net sink or source of H_2O_2 , if the current set-point for matrix $[\text{H}_2\text{O}_2]_{ss}$ is lower or higher, respectively, than that of the cytosol.

Implications and future work

Here, we briefly elaborate on potential physiological consequences of our hypothesized role for mitochondria in H_2O_2 regulation on

working skeletal muscle. We also consider the influence of body temperature, inter-tissue variation and the possible role of post-translational protein modification.

Skeletal muscle mitochondria, oxidative stress and intense activity

Sustained intense physical activity, such as long-distance migration, increases levels of oxidative stress markers. This has been ascribed to a putative increase in mitochondrial ROS production during periods of intense oxygen consumption. However, physical activity increases the turnover rate of ATP, thus increasing ADP availability to the mitochondria. Here, we have shown that while H_2O_2 consumption is not significantly affected, H_2O_2 production decreases profoundly in the presence of high levels of ADP. Interestingly, a recent study provides some benchmarks for the physiological levels of %NAD(P)H in rat muscle mitochondria by comparing mixes of respiratory substrates and effectors mimicking the *in vivo* environment for three levels of physical activity: ‘rest’, ‘mild aerobic exercise’ and ‘intense aerobic exercise’ (Goncalves et al., 2015). The %NAD(P)H corresponding to these conditions are indicated in Fig. 4C. Comparing rest (the highest rate of H_2O_2 production) with intense exercise (the lowest rate of production) indicates an 85% decline in $V_{p,app}$ across this range of activity (Goncalves et al., 2015). In contrast, our results (Fig. 4C), would predict minimal effects on $V_{c,app}$ for the same decrease in %NAD(P)H. If this prediction and our model are correct, then – according to Eqn 1 – ‘intense’ aerobic physical activity should lead to a decrease in the $[H_2O_2]_{ss}$ as compared with ‘rest’ conditions. Additionally, we have previously argued that the acidification that occurs with intense muscle activity may shift mitochondria towards a more antioxidant state (Banh and Treberg, 2013).

In view of the combined effects of the expected decline in H_2O_2 production and maintained H_2O_2 consumption, which should reduce the $[H_2O_2]_{ss}$ set by muscle mitochondria during physical activity, it seems unlikely that mitochondrial function promotes oxidative stress during periods of high demand for ATP in the cell. Reconciling this with the observed increase in markers of oxidative stress after intense exercise (reviewed in Powers and Jackson, 2008) will require further investigation, but these arguments support a role for extramitochondrial sources of ROS, including NADPH oxidases, phospholipase A2 and lipoxygenases, as the cause of the observed oxidative stress.

Rapid changes in temperature

We recently demonstrated that H_2O_2 production by fish red muscle mitochondria is more sensitive to changes in assay temperature than are the reductases that supply NADPH to the respiration-dependent H_2O_2 consumers (Banh et al., 2016). We hypothesized that this ‘thermal mismatch’ could implicate mitochondria as a bona fide source of oxidative stress during acute heat stress in ectotherms. A mismatch in the temperature sensitivity of H_2O_2 production and H_2O_2 consumers may be particularly significant for animals that experience large and regular temperature fluctuations, such as intertidal species (Somero, 2002).

Mammalian torpor and hibernation

Many small endothermic hibernators go through repeated bouts of metabolic depression and torpor at low body temperature which are interrupted by acute and rapid rewarming by 20°C or more during the interbout euthermia phase (Geiser, 2004). Applying our thermal-mismatch observation (see above) to hibernating mammals predicts that decreasing body temperature could

passively contribute to decreasing $[H_2O_2]_{ss}$ during torpor. A previous study using isolated liver and skeletal muscle mitochondria of hibernating 13-lined ground squirrels found that mitochondrial H_2O_2 efflux often decreases as assay temperature declines from 37°C to 10°C (Brown et al., 2012). Although the interaction of physiological state (interbout euthermic, torpid or summer-active animals), tissue, assay and respiratory substrates added leads to a complex pattern to interpret (Brown et al., 2012), one consistency arises: efflux of H_2O_2 is lower for mitochondria isolated from torpid individuals when measured at a physiologically relevant temperature corresponding to torpor (10°C) compared with mitochondria isolated from individuals during interbout euthermia and measured at 37°C (Brown et al., 2012).

Factors that affect mitochondrial $[H_2O_2]_{ss}$ during torpor may, however, be much more diverse than simple temperature effects. Mitochondrial substrate oxidation capacity is markedly decreased, especially for succinate but also with NADH-linked substrates, during the torpor phase of hibernation (Staples, 2014, 2016). As discussed above, a decrease in mitochondrial capacity for substrate oxidation may affect the production of H_2O_2 more than the pathways for its consumption, thereby decreasing $[H_2O_2]_{ss}$. However, there are indications that proteomic and allosteric regulation of mitochondrial substrate oxidation may form part of the adaptation to torpor (reviewed in Staples, 2014, 2016). Post-translational modifications such as glutathionylation may decrease enzymatic flux and, at the same time, either decrease or increase ROS production as seen with complex I and II, respectively (reviewed in Mailloux and Treberg, 2016). Therefore, the declining substrate oxidation capacity may not necessarily lead to declining H_2O_2 production.

Very little research has been conducted on regulation (proteomic and/or allosteric) of the pathways for the consumption of H_2O_2 during hibernation and torpor. Levels of peroxiredoxins, including the mitochondrion-specific peroxiredoxin-3, increase in brown adipose tissue and heart of 13-lined ground squirrels during hibernation (Morin and Storey, 2007), which presumably should elevate the mitochondrial capacity for H_2O_2 consumption, thereby leading to lower $[H_2O_2]_{ss}$; however, this remains to be empirically tested.

In contrast with baseline torpor, the short period of rewarming from a torpor episode may lead to increased cell-level oxidative stress. Support for this stress includes increased antioxidant (ascorbate) uptake from plasma during rewarming (Tøien, 2001), and the recruitment of hypoxia-inducible factor 1 α (Ma et al., 2005) and elevated protein carbonyls and TBARS in late arousal phase (Orr et al., 2009). Oxidative stress during arousal from torpor has traditionally been attributed by some to the increased oxygen consumption by mitochondria. For instance, a previous attempt to estimate the elevated ROS formation was based on an assumed direct relationship between oxygen consumption and the fraction of oxygen diverted to ROS production by mitochondria (% free radical leak) during rewarming (Orr et al., 2009). However, as discussed above, our model shows that elevated oxygen consumption (either by demand for ATP or increased proton leak) and the associated decrease in PMF is expected to temporarily lower mitochondrial $[H_2O_2]_{ss}$. Clearly, the interaction between oxidative stress and rewarming is potentially more complex than the simple effect of oxygen consumption by mitochondria.

It will be important to reconcile how during hibernation, especially during the acute phase of rewarming leading into interbout euthermia, there is shifting metabolic capacity for mitochondrial substrate oxidation combined with alterations in mitochondrial H_2O_2 metabolism (both production and

consumption). This may be an important test of our hypothesized role for mitochondria as regulators of H_2O_2 .

Other organs and tissues

The nature and capacity of mitochondrial H_2O_2 consumers have only been explored for a limited number of tissues, so it is difficult to generalize the nature of H_2O_2 regulation across multiple tissues. Brain mitochondria are likely to regulate H_2O_2 similarly to muscle mitochondria, as they also rely on respiration-dependent pathways for the consumption of H_2O_2 , with minimal involvement of catalase. Brain and muscle mitochondria also have similar H_2O_2 consumption rates: ~ 11 and $6 \text{ nmol min}^{-1} \text{ mg}^{-1}$ protein, respectively (Dreschel and Patel, 2010; Munro et al., 2016).

It is difficult to estimate how qualitative and quantitative inter-tissue differences in mitochondrial H_2O_2 consumption translate into functional differences in regulation of matrix $[H_2O_2]_{ss}$. Although the effectiveness of catalase in maintaining very low levels of $[H_2O_2]$ is questionable owing to its low affinity [the K_m for H_2O_2 is typically found in the range of 80 to 100 mmol l^{-1} in mammals (Switala and Loewen, 2002)], catalase has an exceptionally high catalytic turnover (k_{cat}) and we have observed that liver mitochondria (which have a high catalase content) can readily consume a bolus of $2.5 \text{ } \mu\text{mol l}^{-1} H_2O_2$ to below detection capacity (i.e. below $0.1\text{--}0.2 \text{ } \mu\text{mol l}^{-1}$) in the absence of additional respiratory substrates (D.M. and J.R.T., unpublished observations). Hence, we cannot exclude the idea that catalase may well achieve respiration-independent regulation of H_2O_2 in tissues where it is highly expressed. In fact, liver mitochondria have rates of H_2O_2 consumption that are ~ 10 to 20 times higher than those of muscle and brain; their high catalase content may help to explain this (Dreschel et al., 2010; Lopert and Patel, 2014; D.M., unpublished observations). Thus, there may be some tissues where mitochondria would regulate H_2O_2 predominately in response to the current mitochondrial energetic state (skeletal muscle and brain), whereas in other tissues, such as liver, respiration-independent H_2O_2 consumption could be the dominant mitochondrial contribution. This will be important to consider when extending the arguments provided here to new tissues or other species.

Post-translational regulation

Except for brief mention in regards to hibernation and torpor, we have not considered the regulation of H_2O_2 at the level of specific enzymes; however, it is probable that allosteric or post-translational regulation may add additional levels of complexity. For example, macromolecular supercomplexes have been implicated in altering the production of superoxide (Genova and Lenaz, 2014), which – alongside modifications such as glutathionylation – could influence mitochondrial H_2O_2 production (Mailloux and Treberg, 2016). Likewise, post-translational modification could affect H_2O_2 consumption pathways, which would alter k in our model and thereby alter the $[H_2O_2]_{ss}$. Thus, our model fits a scenario where there should be a wide range of physiologically adjustable $[H_2O_2]_{ss}$ set-points possible in order to reflect and communicate mitochondrial status via redox signalling.

Conclusions

In this Commentary, we have shown that mitochondria have the attributes necessary to act as regulators of H_2O_2 concentration. The model described herein relies on the existence of a balance between H_2O_2 production and consumption by mitochondria. The existence of such a balance would indicate that mitochondrial antioxidant systems are not simply minimizing oxidative damage but, instead,

are an integral component of the ROS regulatory system. Moreover, the model discussed here would allow for mitochondria to alter the $[H_2O_2]_{ss}$ based on changes in the dynamics of ROS metabolism. Mitochondrial H_2O_2 production appears to be far more sensitive to changes in mitochondrial energetics than are the consumption pathways. Thus, mitochondrial H_2O_2 production could be an important link in redox signalling within the mitochondrion and between the mitochondria and the nucleus. If mitochondria are regulators of cellular H_2O_2 concentration, this will change our understanding of the role of mitochondrial ROS metabolism in environmental adaptation, oxidative stress in response to life-history or metabolic trade-offs, metabolic dysfunction and signalling. However, future work should expand these lines of investigation to other types of mitochondria (from different tissues and species) in order to further test this model. It will also be important to extend these experiments from isolated mitochondria to cell- and tissue-level work in order to test whether mitochondria also may act as significant regulators of H_2O_2 *in situ*.

Acknowledgements

The authors thank Joy Stacey for editorial comments and the feedback provided by three reviewers.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grant (418503-2012) and the Canada Research Chairs program (223744), as well as a postdoctoral fellowship to D.M. from the Fonds de Recherche du Québec – Nature et Technologies (FRQ-NT; grant 183703).

References

- Andreyev, A. Y., Kushnareva, Y. E. and Starkov, A. A. (2005). Mitochondrial metabolism of reactive oxygen species. *Biochemistry* **70**, 200–214.
- Arnaiz, S. L., Travacio, M., Llesuy, S. and Boveris, A. (1995). Hydrogen peroxide metabolism during peroxisome proliferation by fenofibrate. *Biochim. Biophys. Acta* **1272**, 175–180.
- Banh, S. and Treberg, J. R. (2013). The pH sensitivity of H_2O_2 metabolism in skeletal muscle mitochondria. *FEBS Lett.* **587**, 1799–1804.
- Banh, S., Wiens, L., Sotiri, E. and Treberg, J. R. (2016). Mitochondrial reactive oxygen species production by fish muscle mitochondria: potential role in acute heat-induced oxidative stress. *Comp. Biochem. Physiol.* **191**, 99–107.
- Boveris, A. and Chance, B. (1973). The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem. J.* **134**, 707–716.
- Boveris, A., Oshino, N. and Chance, B. (1972). The cellular production of hydrogen peroxide. *Biochem. J.* **128**, 617–630.
- Brand, M. D. (2000). Uncoupling to survive? The role of mitochondrial inefficiency in ageing. *Exp. Gerontol.* **35**, 811–820.
- Brand, M. D. (2010). The sites and topology of mitochondrial superoxide production. *Exp. Gerontol.* **45**, 466–472.
- Brand, M. D. (2016). Mitochondrial generation of superoxide and hydrogen peroxide as the source of mitochondrial redox signaling. *Free Radic. Biol. Med.* **100**, 14–31.
- Brown, G. C. and Borutaite, V. (2012). There is no evidence that mitochondria are the main source of reactive oxygen species in mammalian cells. *Mitochondrion* **12**, 1–4.
- Brown, J. C. L., Chung, D. J., Belgrave, K. R. and Staples, J. F. (2012). Mitochondrial metabolic suppression and reactive oxygen species production in liver and skeletal muscle of hibernating thirteen-lined ground squirrels. *Am. J. Physiol.: Reg. Integr. Comp. Physiol.* **302**, R15–R28.
- Cochemé, H. M., Quin, C., McQuaker, S. J., Cabreiro, F., Logan, A., Prime, T. A., Abakumova, I., Patel, J. V., Fearnley, I. M., James, A. M. et al. (2011). Measurement of H_2O_2 within living *Drosophila* during aging using a ratiometric mass spectrometry probe targeted to the mitochondrial matrix. *Cell Metab.* **13**, 340–350.
- Costantini, D. (2008). Oxidative stress in ecology and evolution: lessons from avian studies. *Ecol. Lett.* **11**, 1238–1251.
- Dai, D.-F., Chiao, Y. A., Marcinek, D. J., Szeto, H. H. and Rabinovitch, P. S. (2014). Mitochondrial oxidative stress in aging and healthspan. *Longev Healthspan* **3**, 6.
- Day, B. J. (2014). Antioxidant therapeutics: Pandora's box. *Free Rad. Biol. Med.* **66**, 58–64.

- Drechsel, D. A. and Patel, M.** (2010). Respiration-dependent H₂O₂ removal in brain mitochondria via the thioredoxin/peroxiredoxin system. *J. Biol. Chem.* **285**, 27850-27858.
- Geiser, F.** (2004). Metabolic rate and body temperature reduction during hibernation and daily torpor. *Annu. Rev. Physiol.* **66**, 239-274.
- Genova, M. L. and Lenaz, G.** (2014). Functional role of mitochondrial respiratory supercomplexes. *Biochim. Biophys. Acta - Bioenergetics* **1837**, 427-443.
- Goncalves, R. L. S., Quinlan, C. L., Perevoshchikova, I. V., Hey-Mogensen, M. and Brand, M. D.** (2015). Sites of superoxide and hydrogen peroxide production by muscle mitochondria assessed *ex vivo* under conditions mimicking rest and exercise. *J. Biol. Chem.* **290**, 209-227.
- Jensen, P. K.** (1966). Antimycin-insensitive oxidation of succinate and reduced nicotinamide-adenine dinucleotide in electron-transport particles I. pH dependency and hydrogen peroxide formation. *Biochim. Biophys. Acta* **122**, 157-166.
- Lopert, P. and Patel, M.** (2014). Nicotinamide nucleotide transhydrogenase (Nnt) links the substrate requirement in brain mitochondria for hydrogen peroxide removal to the thioredoxin/peroxiredoxin (Trx/Prx) system. *J. Biol. Chem.* **289**, 15611-15620.
- Loschen, G., Flohé, L. and Chance, B.** (1971). Respiratory chain linked H₂O₂ production in pigeon heart mitochondria. *FEBS Lett.* **18**, 261-264.
- Mailloux, R. J. and Treberg, J. R.** (2016). Protein S-glutathionylation links energy metabolism to redox signaling in mitochondria. *Redox Biol.* **8**, 110-118.
- Ma, Y. L., Zhu, X., Rivera, P. M., Tøien, O., Barnes, B. M., LaManna, J. C., Smith, M. A. and Drew, K. L.** (2005). Absence of cellular stress in brain after hypoxia induced by arousal from hibernation in Arctic ground squirrels. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **289**, R1297-R1306.
- Mason, S. and Wadley, G. D.** (2014). Skeletal muscle reactive oxygen species: a target of good cop/bad cop for exercise and disease. *Redox Rep.* **19**, 97-106.
- Monaghan, P., Metcalfe, N. B. and Torres, R.** (2009). Oxidative stress as a mediator of life history trade-offs: mechanisms, measurements and interpretation. *Ecol. Lett.* **12**, 75-92.
- Moreno-Loshuertos, R., Acín-Pérez, R., Fernández-Silva, P., Movilla, N., Pérez-Martos, A., Rodríguez de Córdoba, S., Gallardo, M. E., Enríquez, J. A., Acín-Pérez, R., Fernández-Silva, P. et al.** (2006). Differences in reactive oxygen species production explain the phenotypes associated with common mouse mitochondrial DNA variants. *Nat. Genet.* **38**, 1261-1268.
- Morin, P. Jr and Storey, K. B.** (2007). Antioxidant defense in hibernation: cloning and expression of peroxiredoxins from hibernating ground squirrels, *Spermophilus tridecemlineatus*. *Arch. Biochem. Biophys.* **461**, 59-65.
- Munro, D., Banh, S., Sotiri, E., Tamanna, N. and Treberg, J. R.** (2016). The thioredoxin and glutathione-dependent H₂O₂ consumption pathways in muscle mitochondria: Involvement in H₂O₂ metabolism and consequence to H₂O₂ efflux assays. *Free Rad. Biol. Med.* **96**, 334-346.
- Orr, A. L., Lohse, L. A., Drew, K. L. and Hermes-Lima, M.** (2009). Physiological oxidative stress after arousal from hibernation in Arctic ground squirrel. *Comp. Biochem. Physiol.* **153**, 213-221.
- Orr, A. L., Vargas, L., Turk, C. N., Baaten, J. E., Matzen, J. T., Dardov, Attle, S. J., Quackenbush, D. C., Goncalves, R. L. S., Perevoshchikova, I. V. et al.** (2015). Suppressors of superoxide production from mitochondrial complex III. *Nat. Chem. Biol.* **11**, 834-836.
- Palomero, J., Pye, D., Kabayo, T., Spiller, D. G. and Jackson, M. J.** (2008). In situ detection and measurement of intracellular reactive oxygen species in single isolated mature skeletal muscle fibers by real time fluorescence microscopy. *Antioxid. Redox Signal.* **10**, 1463-1474.
- Powers, S. K. and Jackson, M. J.** (2008). Exercise-induced oxidative stress: cellular mechanisms. and impact on muscle force production. *Physiol. Rev.* **88**, 1243-1276.
- Quinlan, C. L., Treberg, J. R., Perevoshchikova, I. V., Orr, A. L. and Brand, M. D.** (2012). Native rates of superoxide production from multiple sites in isolated mitochondria measured using endogenous reporters. *Free Rad. Biol. Med.* **53**, 1807-1817.
- Scherz-Shouval, R. and Elazar, Z.** (2011). Regulation of autophagy by ROS: physiology and pathology. *Trends in Biochem. Sci.* **36**, 30-38.
- Sies, H.** (2014). Role of metabolic H₂O₂ generation: redox signaling and oxidative stress. *J. Biol. Chem.* **289**, 8735-8741.
- Somero, G. N.** (2002). Thermal physiology and vertical zonation of intertidal animals: optima, limits, and costs of living. *Integr. Comp. Biol.* **42**, 780-789.
- Staples, J. F.** (2014). Metabolic suppression in mammalian hibernation: the role of mitochondria. *J. Exp. Biol.* **217**, 2032-2036.
- Staples, J. F.** (2016). Metabolic flexibility: hibernation, torpor, and estivation. *Comp. Physiol.* **6**, 737-771.
- Starkov, A. A.** (2008). The role of mitochondria in reactive oxygen species metabolism and signaling. *Annal. New York Acad. Sci.* **1147**, 37-52.
- Starkov, A. A., Andreyev, A., Zhang, S. F., Starkova, N., Korneeva, M., Syromyatnikov, M. and Popov, V. N.** (2014). Scavenging of H₂O₂ by mouse brain mitochondria. *J. Bioenerg. Biomembr.* **46**, 471-477.
- Switala, J. and Loewen, P. C.** (2002). Diversity of properties among catalases. *Arch. Biochem. Biophys.* **401**, 145-154.
- Tøien, Ø., Drew, K. L., Chao, M. L. and Rice, M. E.** (2001). Ascorbate dynamics and oxygen consumption during arousal from hibernation in Arctic ground squirrels. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **281**, R572-R583.
- Treberg, J. R., Quinlan, C. L. and Brand, M. D.** (2010). Hydrogen peroxide efflux from muscle mitochondria underestimates matrix superoxide production - a correction using glutathione depletion. *FEBS J.* **277**, 2766-2778.
- Treberg, J. R., Munro, D., Banh, S., Zacharias, P. and Sotiri, E.** (2015). Differentiating between apparent and actual rates of H₂O₂ metabolism by isolated rat muscle mitochondria to test a simple model of mitochondria as regulators of H₂O₂ concentration. *Redox Biol.* **5**, 216-224.
- Trushina, E. and McMurray, C. T.** (2007). Oxidative stress and mitochondrial dysfunction in neurodegenerative diseases. *Neuroscience* **145**, 1233-1248.
- Zoccarato, F., Cavallini, L. and Alexandre, A.** (2004). Respiration-dependent removal of exogenous H₂O₂ in brain mitochondria: inhibition by Ca²⁺. *J. Biol. Chem.* **279**, 4166-4174.