

## **Reduced non-bicarbonate skeletal muscle buffering capacity in mice with the mini-muscle phenotype**

Jarren C. Kay, Jocelyn Ramirez, Erick Contreras, and Theodore Garland, Jr.\*

Department of Evolution, Ecology, and Organismal Biology, University of California, Riverside,  
CA 92521, USA

\*Corresponding Author:

Phone: 951-827-3524

Fax: 951-827-4286

Email: [tg Garland@ucr.edu](mailto:tg Garland@ucr.edu)

## ABSTRACT

Muscle pH decreases during exercise, which may impair function. Endurance training typically reduces muscle buffering capacity due to changes in fiber type composition, but existing comparisons of species that vary in activity level are ambiguous. We hypothesized that High Runner (HR) lines of mice from an experiment that breeds for voluntary wheel running would have altered muscle buffering capacity as compared with their non-selected control counterparts. We also expected that six days of wheel access, as used in the selection protocol, would reduce buffering capacity, especially for HR mice. Finally, we expected a subset of HR mice with the "mini-muscle" phenotype to have relatively low buffering capacity due to fewer type IIb fibers. We tested non-bicarbonate buffering capacity of thigh muscles. Only HR mice expressing the mini-muscle phenotype had significantly reduced buffering capacity, females had lower buffering capacity than males, and wheel access had no significant effect.

**KEY WORDS:** Artificial selection, Buffering Capacity, Endurance, Voluntary Exercise, Skeletal muscle, Wheel running

## INTRODUCTION

During exercise, metabolic acidosis may occur from by-products of muscle contraction and ATP production. This acidosis may impair muscle function in various ways, including decreased rates of glycolysis and glycogenolysis due to inhibition of glycogen phosphorylase and phosphofructokinase activities, and through decreased maximum force (see Cairns, 2006 for review; Chase and Kushmerick, 1988; Chasiotis et al., 1983; Donaldson and Hermansen, 1978; Fitts, 1994; Trivedi and Danforth, 1966).

Organisms ameliorate decreases in muscle pH in at least two ways. One is to remove protons generated in the cytosolic compartment by transporting them to the mitochondria or out of the cell entirely. Transport of protons into the mitochondria using monocarboxylate transporter 1 (MCT1) also transports lactate, which can be used by the mitochondria and converted to pyruvate (intracellular lactate shuttle; Brooks, 1998; Brooks et al., 1999). Proton transport out of the cell is accomplished by numerous mechanisms, but MCT proteins are of particular interest because lactate may then be taken up by other muscle cells via MCT1 (usually by type I or type IIa fibers; Donovan and Pagliassotti, 2000) and converted into pyruvate via lactate dehydrogenase.

The other way to mitigate decreased pH is to buffer it, which is especially important in the blood and muscle of organisms that must undergo an oxygen deficit, such as during some types of exercise or diving. Organisms accomplish this in numerous ways, including the use of  $\text{HCO}_3^-$  and imidazole-containing (histidine-containing) compounds (carnosine, anserine, etc.), production of lactate, the hydrolysis of phosphocreatine (and subsequent production of  $\text{P}_i$ ), and behavioral changes (e.g., increased ventilation). As might be expected, marine mammals and pelagic fish have higher buffering capacity than terrestrial mammals or deep-sea/shallow water

fish, respectively (see Abe, 2000 for review; Castellini and Somero, 1981; Okuma and Abe, 1992). Greyhounds and thoroughbred horses, which regularly perform extended bouts of exercise, have increased buffering capacity as compared with humans of unspecified training status (Harris et al., 1990). In humans, buffering capacity changes in response to training, and differentially depending on the type of training (e.g., high-intensity interval training increases buffering capacity; Parkhouse et al., 1985; Sahlin and Henriksson, 1984; Weston et al., 1997). Possible mechanisms underlying the training responses in humans include increases in the amount of buffering compounds in the muscle (potentially by changing the number of fast glycolytic fibers), increases in the number of proton transporters (Juel et al., 2004a; Juel et al., 2004b; Juel et al., 2003; Pilegaard et al., 1999; Pilegaard et al., 1994), and/or increases in muscle capillarity (Jensen et al., 2004).

Given its association with athletic abilities, buffering capacity would be expected to evolve in concert (coadapt) when artificial selection targets forced or voluntary exercise (review in Swallow et al., 2009). The purpose of the present study was to examine muscle buffering capacity in replicate high-runner (HR) lines of mice that have experienced long-term breeding for high voluntary wheel running (Swallow et al., 1998). Several correlated responses in the HR lines seem to enhance capacity for endurance exercise, including more intermittent locomotion on wheels, altered stride characteristics during treadmill locomotion, increased maximal oxygen consumption ( $VO_2\text{max}$ ), increased heart ventricle mass, and larger femoral heads (Claghorn et al., 2017; Garland and Freeman, 2005; Girard et al., 2001; Kelly et al., 2017; Kelly et al., 2006; Rezende et al., 2006).

A subset of the HR mice, known as mini-muscle individuals, have a 50% reduction in hindlimb muscle mass when homozygous for a SNP mutation in the *Myh4* gene (Burniston et al., 2013; Kelly et al., 2014). This mutation leads to a severe reduction in IIB muscle fibers throughout the body (Talmadge et al., 2014), causes increased cost of transport, reduced maximal sprint speed, and differences in contractile properties of some muscles in the triceps surae complex (e.g., slower twitches; Dlugosz et al., 2009; Guderley et al., 2006; Syme et al., 2005). Mice with the mini-muscle phenotype also show increased capillarity in the medial gastrocnemius, increased VO<sub>2</sub>max during hypoxia, and decreased plasma [lactate] during peak wheel running, but not during exhaustive exercise (Meek et al., 2009; Rezende et al., 2006; Wong et al., 2009). Thus, mini-muscle individuals may have altered buffering capacity.

## **MATERIALS AND METHODS**

### **Experimental animals**

All procedures in this study were approved by and are in accordance with guidelines set forth by the Institutional Animal Care and Use Committee at the University of California, Riverside. Mice used in the present study are the same as in Thompson et al. (2017), which came from generation 74 of an ongoing artificial selection experiment in which mice are bred for high voluntary wheel-running behavior (Careau et al., 2013; Swallow et al., 1998). The founding population was 224 outbred Hsd:ICR mice (*Mus domesticus*). After two generations of random mating, individuals were randomly assigned to one of eight closed lines; four lines designated as non-selected control (C) and 4 designated as high runner (HR) lines. In the selection protocol, all mice were given access to Wahman-type activity wheels (1.12 m circumference, 10 cm wide,

35.7 cm diameter) attached to home-cages with *ad libitum* food and water. The HR mice were bred based on their mean amount of wheel running on days 5 and 6 of a 6-day trial, while C mice were bred without regard to their wheel running. In all cases, sibling mating was not allowed.

This study used 50 male and 50 female mice (evenly split between C and HR) that were not allowed access to wheels, along with another 50 male and 50 female mice that were given access to running wheels for six days (also evenly split between C and HR) immediately prior to dissection. A period of six days was chosen in order to gain insight regarding their running abilities in a way that matched the criterion used for selective breeding. As in Thompson et al. (2017), experimental animals were on a reverse photoperiod with lights on from 19:00 to 07:00. Mini-muscle status was determined by dissection of the triceps surae muscle complex (Garland et al., 2002). Because the mini-muscle phenotype includes altered fiber type of the hindlimb muscles (Bilodeau et al., 2009; Guderley et al., 2006; Guderley et al., 2008; Talmadge et al., 2014), it was used as an additional main effect in all statistical analyses (see below).

### **Wheel running**

For the half of the mice having access to wheels, revolutions were recorded for 23 hours/day by a computer that records revolutions in one-minute intervals (Careau et al., 2013; Swallow et al., 1998). Mice were removed from wheels during peak wheel running on day 6 and immediately sacrificed. Wheel freeness (measured by accelerating the wheel to a known velocity and recording the number of free-spinning revolutions) was used as a covariate in all analyses of wheel running (e.g., see Acosta et al., 2017; Copes et al., 2015; Kolb et al., 2010).

### **Home-cage activity**

Home-cage activity (HCA, an indicator of spontaneous physical activity; Garland et al., 2011) was measured using passive infrared motion sensors placed inside each cage. As previously described (Acosta et al., 2017; Copes et al., 2015; Thompson et al., 2017), HCA sensors were interfaced with a computer and record activity 3 times/second as a binary output (0 = no movement, 1 = movement) and then readings are averaged across every one-minute interval. As for wheel running, HCA was measured for 23h/day. Sensor sensitivity was used as an additional covariate in these analyses.

### **Buffering capacity**

Right thigh muscles from all mice were studied for buffering capacity. All mice were sacrificed via cardiac puncture (under anesthesia) at the time when peak wheel running would occur on the sixth day, dissected, and thigh muscles were frozen and stored at  $-80^{\circ}\text{C}$ . (Age at sampling was 71-91 days.) Muscle buffering capacity ( $\beta_{\text{m}}$  or Slyke; defined as the number of  $\mu\text{moles}$  of base needed to change the pH of muscle homogenate by one pH unit per gram of wet weight of muscle) was determined using methods modified from Castellini and Somero (1981). Muscles were homogenized in 0.9% NaCl solution on ice (1g of muscle in 19 ml of NaCl solution), and 2 ml were then transferred to a scintillation vial to be tested for buffering capacity at a constant temperature of  $37\pm 0.5^{\circ}\text{C}$ . Initial pH of homogenized muscle was taken, and 0.5N HCl was used to reduce the pH to 6, then, 0.0125N NaOH was used in 10  $\mu\text{l}$  increments to bring the pH up to 7. Preliminary studies showed no effect of total protein concentration on buffering capacity and therefore it was not measured in this experiment.

## Statistics

Dependent variables were transformed as needed to improve normality and homoscedasticity of residuals. Following previous studies of mice from this selection experiment (e.g., Belter et al., 2004; Meek et al., 2009; Swallow et al., 1998; Thompson et al., 2017), buffering capacity was analyzed by nested analysis of covariance (ANCOVA), with replicate line nested within linetype (HR vs C) using SAS Procedure Mixed. Analyses of wheel running and home-cage activity across 5 days were done by SAS Procedure Mixed repeated-measures ANCOVA. The main factors for all analyses were linetype (HR vs C), sex, and the mini-muscle phenotype (Thompson et al., 2017), and wheel access was also used when applicable. As noted above, a measure of wheel freeness and a measure of home-cage sensor sensitivity were used as covariates. For all analyses, age at sacrifice was used as a covariate. In preliminary analyses, body mass was never a significant predictor of home-cage activity, so it was removed for final analyses (following Thompson et al., 2017).

For analyses of buffering capacity (always log-transformed to improve normality of residuals), time at sacrifice was used as an additional covariate. Because previous studies have shown fiber type alteration within muscle due to as little as 1 week of training (Allen et al., 2001), we tested whether the amount of physical activity was a predictor of buffering capacity and if the level of activity would change the magnitude of the main effects or the interactions. Therefore, in some analyses of buffering capacity we also used the total amount of wheel running (assigning mice without wheels values of zero) and/or the total amount of home-cage activity as covariates (following Thompson et al., 2017). Note that the mini-muscle phenotype is



absent from all C lines, fixed (100% of individuals have it) in one of the HR lines, and polymorphic in another HR line, so mini-muscle status is confounded with linetype. Therefore, we performed additional analyses removing line and linetype from the model and performing an ANCOVA of buffering capacity for the four sex X linetype groups of mice. Statistical significance was judged at  $p < 0.05$ .

## RESULTS AND DISCUSSION

### Wheel running

As shown previously (e.g., see Belter et al., 2004), HR mice ran significantly more revolutions across days 1-5 than their C counterparts (repeated-measures ANCOVA,  $p = 0.0068$ ; see Fig. 1A and Table S1), with a strong effect of day ( $p < 0.0001$ ) and a linetype\*day interaction ( $p < 0.0001$ ). Figure 1A shows that HR mice had a dramatic increase from ~8,000 and ~8,500 revolutions/day for males and females, respectively, on day 1 to ~11,000 and ~14,000 revolutions/day on day 5. In contrast, both sexes of C mice had a smaller absolute and proportional increase from ~4,000 revolutions on day 1 to ~5,000 on day 5. Figure 1A suggests that females tended to run more than males in HR lines, but not in C lines, but the linetype\*sex interaction was non-significant ( $p = 0.2431$ ). The pattern for total revolutions per day was largely mirrored by the pattern for average running speed (Fig. 1B), with a strong effect of linetype ( $p = 0.0029$ ), day ( $p < 0.0001$ ), and their interaction ( $p < 0.0001$ ). The amount of time spent running was higher for HR than for C ( $p = 0.0157$ ; Fig. 1C) and higher for females than males ( $p = 0.0391$ ), with a strong effect of day ( $p < 0.0001$ ), where most groups ran the most on day 1, likely an effect of novelty (Table S1). ANCOVA indicated a significant positive effect of

mini-muscle phenotype on revolutions per day ( $p = 0.0119$ ; Fig. 1D) and average running speed ( $p = 0.0240$ ; Fig. 1E), but not running duration (Fig. 1F).

### **Home-cage activity**

Similar to results from previous studies of animals housed without wheels (Acosta et al., 2017; Copes et al., 2015; Thompson et al., 2017), HR mice had higher total activity, minutes per day of activity, average activity per minute, and maximal activity for any one-minute interval (Fig 2, left panels and Table S2: linetype  $p = 0.0015$ ,  $p = 0.0235$ ,  $p = 0.0007$ , and  $p = 0.0018$ , respectively). In addition, females without wheels had more total activity, more time spent active, more activity per minute, and higher maximal activity for any one-minute interval than male mice ( $p = 0.0133$ ,  $p = 0.0134$ ,  $p = 0.0410$ , and  $p = 0.0207$ , respectively; Fig. 2A-D and Table S2). Mini-muscle individuals without wheels had significantly lower total activity ( $p = 0.0061$ ; Fig. 2E), activity/minute ( $p = 0.0002$ ; Fig. 2G). Male mini-muscle mice without wheels had lower maximum activity in any 1-min interval than other mice (sex\*mini interaction  $p = 0.0114$ ; Fig. 2H). No other interaction terms were statistically significant (Table S2). All measures of activity tended to decline across days for all groups (all day  $p < 0.0001$ ).

With access to wheels, the only linetype effect was that HR mice were active in cages for significantly more time than C mice ( $p = 0.0402$ ; Fig 2B), while mini-muscle individuals were active for less time per day as compared with other mice ( $p = 0.0044$ ; Fig. 2F). Again, all measures of activity tended to decline across days (all day  $p < 0.0001$ ), at the same time that wheel running was increasing (Fig. 1), a phenomenon that has been reported previously for these mice (Acosta et al. 2017). No interaction terms were statistically significant for mice with access to wheels (Table S2).

## **Buffering capacity**

Our hypothesis that HR mice in general would have reduced buffering capacity in skeletal muscle was not supported by any statistical model (Fig. 3A and Table S3). However, the subset of HR mice with the mini-muscle phenotype had significantly lower buffering capacity than normal-muscled mice (Fig. 3 and Table S3). The mini-muscle phenotype has been favored (unintentionally) by the selective breeding protocol, and hence can be viewed as an adaptive response to selection, albeit one that occurs in only two of the four replicate HR lines because it was initially rare and the underlying Mendelian recessive allele was lost by random genetic drift in all other lines (Garland et al., 2002). Although it has been favored by selection, the mini-muscle phenotype includes several differences as compared with normal-muscled individuals (e.g., increased fatigue resistance in medial gastrocnemius: see Introduction), and it is not yet clear which one(s) of these provides the primary benefit(s) for endurance running. Indeed, reduced buffering capacity might even be maladaptive for endurance running. For example, a study of endurance capacity during forced treadmill exercise found that the HR line fixed for the mini-muscle phenotype has relatively low endurance compared with the other three HR lines (Meek et al., 2009). However, in the context of the prevailing selective regime, this "cost" appears to be outweighed by other benefits inherent to mini-muscle individuals.

Regardless of its adaptive significance, what might underlie the reduced buffering capacity of mini-muscles? Many previously mentioned comparative studies have shown that fiber type of the muscle and a few histidine-containing compounds are very important in the non-bicarbonate buffering capacity of muscles (carnosine, anserine, and balenine). These compounds

occur in most animals (Crush, 1970; Davey, 1960) but vary in their concentrations (among species; see Bate-Smith, 1938) and in different muscle fiber types (Dunnett and Harris, 1995). Much has been done on anserine and carnosine, which have pKa values of 6.83 and 7.04 (Bate-Smith, 1938; Davey, 1960), respectively, which would make them good buffering compounds. Dunnett and Harris (1995) showed that carnosine concentrations in the middle gluteal muscle of horses are increased in type IIa fibers, but highest in type IIb fibers. Mini-muscle mice have reduced numbers of type IIb fibers (Bilodeau et al., 2009; Guderley et al., 2008; Guderley et al., 2006; Talmadge et al., 2014), meaning that they should show reduced carnosine-mediated buffering capacity. Taurine (an organic acid) was shown to have marked effects on increasing buffering capacity as well, but its distribution is opposite of carnosine, being higher in type I fibers, but still present in lower concentrations in type IIa fibers and type IIb fibers (Dunnett and Harris, 1995). This evidence suggests that the differences in buffering capacity between mini-muscle and normal-muscled individuals are unlikely to be caused by differences in taurine concentrations.

Mini-muscle mice further differentiate themselves from normal-muscled mice by having lower blood lactate during peak wheel running (Meek et al., 2009), which could be caused by either reduced lactate production or increased usage within the mitochondria. If mini-muscle mice are generating less lactate, then they should have reduced ability to resist changes in pH, given that lactate acts as a buffering compound. Mini-muscle mice also have increased mitochondrial densities (Guderley et al., 2006), but lower lactate dehydrogenase activity per gram of muscle tissue (Houle-Leroy et al., 2003), which could prevent them from using the intracellular lactate shuttle to convert lactate to pyruvate in the mitochondria of type I and type IIa fibers. The increased capillarity of mini-muscle mice (Wong et al., 2009) may be more

effective at transporting H<sup>+</sup> away from the muscles (via MCT proteins), as suggested by Juel (2008), which could mean they would not require increased buffering capacity.

Female mice had a lower buffering capacity than males ( $p = 0.0051$ ; see Fig. 3A and Table S3). The differences in buffering between the sexes in these mice may reflect lower carnosine and/or anserine levels in muscles of females, as has been reported in some species of mammals (e.g., humans and mice; Mannion et al., 1992; Peñafiel et al., 2004), but not in others (i.e. rats; Peñafiel et al., 2004). The sex differences seem to be mediated by testosterone in mice, given that treatment of females with testosterone propionate increases their carnosine to levels similar to that seen in males, and hence may also equalize muscle buffering capacity (Peñafiel et al., 2004). Female mice also have lower levels of type IIb fibers in some hindlimb muscles (e.g., the tibialis anterior; see Figure 1 in Haizlip et al., 2015) which could also contribute to lowered buffering in those muscles.

Our results regarding the effects of sex and of mini-muscle phenotype were generally unaltered when we included measures of wheel running and/or of home-cage activity as covariates, and the activity metrics did not significantly predict buffering capacity in these statistical models (Table S3). However, in an analysis using only mice with wheel access, average running speed across 5 days negatively predicted buffering capacity ( $p = 0.0407$ ). Similarly, the sum of the wheel revolutions during the 8 minutes prior to sacrifice negatively predicted buffering capacity ( $p = 0.0440$ ) (preliminary analyses indicated that wheel running over this time interval had a higher predictive ability than over other intervals in the range of 1 through 120 min prior to sacrifice). Both of these results suggest that wheel-running activity may have some influence on muscle buffering capacity, i.e., cause training effects.

Training effects on muscle buffering capacity have been reported previously (Parkhouse et al., 1985; Sahlin and Henriksson, 1984). For example, endurance cyclists have lower muscle buffering capacity than other athletes that perform more short-term, high-intensity exercises (because of reduced numbers of fast glycolytic or type IIb fibers), but high-intensity, interval training can raise their muscle buffering capacity (Weston et al., 1997). This training effect is potentially caused by an increase in type IIb abundance after high-intensity training. [Type IIb fibers generate the most lactate due to their lower oxidative capacity (Baldwin et al., 1977).] In mice, Allen et al. (2001) showed that as little as 1 week of voluntary wheel running (commonly thought of as endurance training) is enough to induce a fiber-type shift in the tibialis anterior muscle, increasing the number of type IIa fibers and reducing the number of type IIb fibers. A change in fiber type toward more oxidative fibers (type I and type IIa) may lower buffering capacity due to either lower lactate or carnosine levels (Dunnett and Harris, 1995; Dunnett et al., 1997). Future studies could explore these possibilities by direct analysis of muscle fiber types at different time points after the initiation of wheel access in HR and C lines of mice.

In conclusion, we have shown that buffering capacity is reduced in mini-muscle mice (Garland et al., 2002) as well as female mice in general. We have also provided possible explanations for these differences (e.g., potentially lower levels of carnosine in mini-muscle mice); however, it is likely that more than one process is occurring simultaneously to affect the buffering capacity of these mice. Future studies should quantify the imidazole-containing peptides in the hindlimb muscles of mini- and normal-muscled mice, as well as the amount of MCT proteins (see Introduction) present in the same muscles. Based on our current knowledge of the fiber type composition in the mini-muscle mice (see references in Introduction), we would

expect lower levels of carnosine and lactate, but potentially increased concentrations of MCT1 proteins due to the lower numbers of type IIb fibers in their hindlimb muscles.

### **Acknowledgements**

We thank Dr. Zoe Thompson for help in obtaining the mice used here. Dr. Douglas A. Syme and Dr. Robert J. Talmadge offered helpful comments on the manuscript.

### **Competing interests**

Nothing to declare.

### **Author contributions**

JCK and TG designed the experiments, analyzed the data, and drafted the manuscript. JCK collected the data; JR and EC assisted with data collection and edited the manuscript.

### **Funding**

Supported by U.S. NSF grants IOS-1121273 and DEB-1655362 to T.G.

### **Supplementary information**

Supplementary information available online at xxt to be added by journal

## References

- Abe, H.** (2000). Role of histidine-related compounds as intracellular proton buffering constituents in vertebrate muscle. *Biochem. CC BIOKHIMIIA* **65**, 757–765.
- Acosta, W., Meek, T. H., Schutz, H., Dlugosz, E. M. and Garland Jr., T.** (2017). Preference for Western diet coadapts in High Runner mice and affects voluntary exercise and spontaneous physical activity in a genotype-dependent manner. *Behav. Processes* **135**, 56–65.
- Allen, D. L., Harrison, B. C., Maass, A., Bell, M. L., Byrnes, W. C. and Leinwand, L. A.** (2001). Cardiac and skeletal muscle adaptations to voluntary wheel running in the mouse. *J. Appl. Physiol.* 1900–1908.
- Baldwin, K. M., Campbell, P. J. and Cooke, D. A.** (1977). Glycogen, lactate, and alanine changes in muscle fiber types during graded exercise. *J. Appl. Physiol.* **43**, 288–291.
- Bate-Smith, E. C.** (1938). The buffering of muscle in rigor; protein, phosphate and carnosine. *J. Physiol.* **92**, 336–343.
- Belter, J. G., Carey, H. V. and Garland Jr., T.** (2004). Effects of voluntary exercise and genetic selection for high activity levels on HSP72 expression in house mice. *J. Appl. Physiol.* **96**, 1270–1276.
- Bilodeau, G. M., Guderley, H., Joanisse, D. R. and Garland Jr., T.** (2009). Reduction of type IIb myosin and IIB fibers in tibialis anterior muscle of mini-muscle mice from high-activity lines. *J. Exp. Zool. Part Ecol. Genet. Physiol.* **311A**, 189–198.
- Brooks, G. A.** (1998). Mammalian fuel utilization during sustained exercise. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **120**, 89–107.
- Brooks, G. A., Dubouchaud, H., Brown, M., Sicurello, J. P. and Butz, C. E.** (1999). Role of mitochondrial lactate dehydrogenase and lactate oxidation in the intracellular lactate shuttle. *Proc. Natl. Acad. Sci.* **96**, 1129–1134.
- Burniston, J. G., Meek, T. H., Pandey, S. N., Broitman-Maduro, G., Maduro, M. F., Bronikowski, A. M., Garland Jr., T. and Chen, Y.-W.** (2013). Gene expression profiling of gastrocnemius of “minimuscle” mice. *Physiol. Genomics* **45**, 228–236.
- Cairns, S. P.** (2006). Lactic acid and exercise performance. *Sports Med.* **36**, 279–291.
- Careau, V., Wolak, M. E., Carter, P. A. and Garland Jr., T.** (2013). Limits to behavioral evolution: the quantitative genetics of a complex trait under directional selection. *Evolution* **67**, 3102–3119.
- Castellini, M. A. and Somero, G. N.** (1981). Buffering capacity of vertebrate muscle: correlations with potentials for anaerobic function. *J. Comp. Physiol.* **143**, 191–198.
- Chase, P. B. and Kushmerick, M. J.** (1988). Effects of pH on contraction of rabbit fast and slow skeletal muscle fibers. *Biophys. J.* **53**, 935.




- Chasiotis, D., Hultman, E. and Sahlin, K.** (1983). Acidotic depression of cyclic AMP accumulation and phosphorylase b to a transformation in skeletal muscle of man. *J. Physiol.* **335**, 197–204.
- Claghorn, G. C., Thompson, Z., Kay, J. C., Ordonez, G., Hampton, T. G. and Garland Jr., T.** (2017). Selective breeding and short-term access to a running wheel alter stride characteristics in house mice. *Physiol. Biochem. Zool.* **90**, 533–545.
- Copes, L. E., Schutz, H., Dlugosz, E. M., Acosta, W., Chappell, M. A. and Garland Jr., T.** (2015). Effects of voluntary exercise on spontaneous physical activity and food consumption in mice: Results from an artificial selection experiment. *Physiol. Behav.* **149**, 86–94.
- Crush, K. G.** (1970). Carnosine and related substances in animal tissues. *Comp. Biochem. Physiol.* **34**, 3–30.
- Davey, C. L.** (1960). The significance of carnosine and anserine in striated skeletal muscle. *Arch. Biochem. Biophys.* **89**, 303–308.
- Dlugosz, E. M., Chappell, M. A., McGillivray, D. G., Syme, D. A. and Garland Jr., T.** (2009). Locomotor trade-offs in mice selectively bred for high voluntary wheel running. *J. Exp. Biol.* **212**, 2612–2618.
- Donaldson, S. K. B. and Hermansen, L.** (1978). Differential, direct effects of H<sup>+</sup> and Ca<sup>2+</sup>-activated force of skinned fibers from the soleus, cardiac and adductor magnus muscles of rabbits. *Pflug. Arch.* **376**, 55–65.
- Donovan, C. M. and Pagliassotti, M. J.** (2000). Quantitative assessment of pathways for lactate disposal in skeletal muscle fiber types: *Med. Sci. Sports Exerc.* **32**, 772–777.
- Dunnett, M. and Harris, R. C.** (1995). Carnosine and taurine contents of type I, IIA and IIB fibres in the middle gluteal muscle. *Equine Vet. J.* **27**, 214–217.
- Dunnett, M., Harris, R. C., Soliman, M. Z. and Suwar, A. A. S.** (1997). Carnosine, anserine and taurine contents in individual fibres from the middle gluteal muscle of the camel. *Res. Vet. Sci.* **62**, 213–216.
- Fitts, R. H.** (1994). Cellular mechanisms of muscle fatigue. *Physiol. Rev.* **74**, 49–94.
- Garland Jr., T. and Freeman, P. W.** (2005). Selective breeding for high endurance running increases hindlimb symmetry. *Evolution* **59**, 1851–1854.
- Garland Jr., T., Morgan, M. T., Swallow, J. G., Rhodes, J. S., Girard, I., Belter, J. G. and Carter, P. A.** (2002). Evolution of a small-muscle polymorphism in lines of house mice selected for high activity levels. *Evolution* **56**, 1267–1275.
- Garland Jr., T., Schutz, H., Chappell, M. A., Keeney, B. K., Meek, T. H., Copes, L. E., Acosta, W., Drenowatz, C., Maciel, R. C., van Dijk, G., et al.** (2011). The biological control of voluntary exercise, spontaneous physical activity and daily energy expenditure in relation to obesity: human and rodent perspectives. *J. Exp. Biol.* **214**, 206–229.

- Girard, I., McAleer, M. W., Rhodes, J. S. and Garland Jr., T.** (2001). Selection for high voluntary wheel-running increases speed and intermittency in house mice (*Mus domesticus*). *J. Exp. Biol.* **204**, 4311–4320.
- Guderley, H., Houle-Leroy, P., Diffie, G. M., Camp, D. M. and Garland Jr., T.** (2006). Morphometry, ultrastructure, myosin isoforms, and metabolic capacities of the “mini muscles” favoured by selection for high activity in house mice. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **144**, 271–282.
- Guderley, H., Joanisse, D. R., Mokas, S., Bilodeau, G. M. and Garland Jr., T.** (2008). Altered fibre types in gastrocnemius muscle of high wheel-running selected mice with mini-muscle phenotypes. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **149**, 490–500.
- Haizlip, K. M., Harrison, B. C. and Leinwand, L. A.** (2015). Sex-based differences in skeletal muscle kinetics and fiber-type composition. *Physiology* **30**, 30–39.
- Harris, R. C., Marlin, D. J., Dunnett, M., Snow, D. H. and Hultman, E.** (1990). Muscle buffering capacity and dipeptide content in the thoroughbred horse, greyhound dog and man. *Comp. Biochem. Physiol.* **97A**, 249–251.
- Houle-Leroy, P., Guderley, H., Swallow, J. G. and Garland Jr., T.** (2003). Artificial selection for high activity favors mighty mini-muscles in house mice. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **284**, R433–R443.
- Jensen, L., Bangsbo, J. and Hellsten, Y.** (2004). Effect of high intensity training on capillarization and presence of angiogenic factors in human skeletal muscle: capillarization in exercise-trained skeletal muscle. *J. Physiol.* **557**, 571–582.
- Juel, C.** (2008). Regulation of pH in human skeletal muscle: adaptations to physical activity. *Acta Physiol.* **193**, 17–24.
- Juel, C., Lundby, C., Sander, M., Calbet, J. A. L. and van Hall, G.** (2003). Human skeletal muscle and erythrocyte proteins involved in acid-base homeostasis: adaptations to chronic hypoxia. *J. Physiol.* **548**, 639–648.
- Juel, C., Holten, M. K. and Dela, F.** (2004a). Effects of strength training on muscle lactate release and MCT1 and MCT4 content in healthy and type 2 diabetic humans: MCT, type 2 diabetes and training. *J. Physiol.* **556**, 297–304.
- Juel, C., Klarskov, C., Nielsen, J. J., Krstrup, P., Mohr, M. and Bangsbo, J.** (2004b). Effect of high-intensity intermittent training on lactate and H<sup>+</sup> release from human skeletal muscle. *AJP Endocrinol. Metab.* **286**, 245E–251.
- Kelly, S. A., Czech, P. P., Wight, J. T., Blank, K. M. and Garland Jr., T.** (2006). Experimental evolution and phenotypic plasticity of hindlimb bones in high-activity house mice. *J. Morphol.* **267**, 360–374.
- Kelly, S. A., Nehrenberg, D. L., Hua, K., Garland Jr., T. and Pomp, D.** (2014). Quantitative genomics of voluntary exercise in mice: transcriptional analysis and mapping of expression QTL in muscle. *Physiol. Genomics* **46**, 593–601.

- Kelly, S. A., Gomes, F. R., Kolb, E. M., Malisch, J. L. and Garland Jr., T.** (2017). Effects of activity, genetic selection and their interaction on muscle metabolic capacities and organ masses in mice. *J. Exp. Biol.* **220**, 1038–1047.
- Kolb, E. M., Kelly, S. A., Middleton, K. M., Sermsakdi, L. S., Chappell, M. A. and Garland Jr., T.** (2010). Erythropoietin elevates but not voluntary wheel running in mice. *J. Exp. Biol.* **213**, 510–519.
- Mannion, A. F., Jakeman, P. M., Dunnett, M., Harris, R. C. and Willan, P. L. T.** (1992). Carnosine and anserine concentrations in the quadriceps femoris muscle of healthy humans. *Eur. J. Appl. Physiol.* **64**, 47–50.
- Meek, T. H., Lonquich, B. P., Hannon, R. M. and Garland Jr., T.** (2009). Endurance capacity of mice selectively bred for high voluntary wheel running. *J. Exp. Biol.* **212**, 2908–2917.
- Okuma, E. and Abe, H.** (1992). Major buffering constituents in animal muscle. *Comp. Biochem. Physiol.* **102A**, 37–41.
- Parkhouse, W. S., McKenzie, D. C., Hochachka, P. W. and Ovalle, W. K.** (1985). Buffering capacity of deproteinized human vastus lateralis muscle. *J. Appl. Physiol.* **58**, 14–17.
- Peñafiel, R., Ruzafa, C., Monserrat, F. and Cremades, A.** (2004). Gender-related differences in carnosine, anserine and lysine content of murine skeletal muscle. *Amino Acids* **26**, 53–58.
- Pilegaard, H., Bangsbo, J., Richter, E. A. and Juel, C.** (1994). Lactate transport studied in sarcolemmal giant vesicles from human muscle biopsies: relation to training status. *J. Appl. Physiol.* **77**, 1858–1862.
- Pilegaard, H., Domino, K., Noland, T., Juel, C., Hellsten, Y., Halestrap, A. P. and Bangsbo, J.** (1999). Effect of high-intensity exercise training on lactate/H<sup>+</sup> transport capacity in human skeletal muscle. *Am. J. Physiol.-Endocrinol. Metab.* **276**, E255–E261.
- Rezende, E. L., Gomes, F. R., Malisch, J. L., Chappell, M. A. and Garland Jr., T.** (2006). Maximal oxygen consumption in relation to subordinate traits in lines of house mice selectively bred for high voluntary wheel running. *J. Appl. Physiol.* **101**, 477–485.
- Sahlin, K. and Henriksson, J.** (1984). Buffer capacity and lactate accumulation in skeletal muscle of trained and untrained men. *Acta Physiol. Scand.* **122**, 331–339.
- Swallow, J. G., Carter, P. A. and Garland Jr., T.** (1998). Artificial selection for increased wheel-running behavior in house mice. *Behav. Genet.* **28**, 227–237.
- Swallow, J. G., Hayes, J. P., Koteja, P. and Garland Jr., T.** (2009). Selection experiments and experimental evolution of performance and physiology. *Exp. Evol. Concepts Methods Appl. Sel. Exp.* 301–351.
- Syme, D. A., Evashuk, K., Grintuch, B., Rezende, E. L. and Garland Jr., T.** (2005). Contractile abilities of normal and “mini” triceps surae muscles from mice (*Mus domesticus*) selectively bred for high voluntary wheel running. *J. Appl. Physiol.* **99**, 1308–1316.


- Talmadge, R. J., Acosta, W. and Garland Jr., T.** (2014). Myosin heavy chain isoform expression in adult and juvenile mini-muscle mice bred for high-voluntary wheel running. *Mech. Dev.* **134**, 16–30.
- Thompson, Z., Argueta, D., Garland Jr., T. and DiPatrizio, N.** (2017). Circulating levels of endocannabinoids respond acutely to voluntary exercise, are altered in mice selectively bred for high voluntary wheel running, and differ between the sexes. *Physiol. Behav.* **170**, 141–150.
- Trivedi, B. and Danforth, W. H.** (1966). Effect of pH on the kinetics of frog muscle phosphofructokinase. *J. Biol. Chem.* **241**, 4110–4114.
- Weston, A. R., Myburgh, K. H., Lindsay, F. H., Dennis, S. C., Noakes, T. D. and Hawley, J. A.** (1997). Skeletal muscle buffering capacity and endurance performance after high-intensity interval training by well-trained cyclists. *Eur. J. Appl. Physiol.* **75**, 7–13.
- Wong, L. E., Garland Jr., T., Rowan, S. L. and Hepple, R. T.** (2009). Anatomic capillarization is elevated in the medial gastrocnemius muscle of mighty mini mice. *J. Appl. Physiol.* **106**, 1660–1667.

# Figures



**Fig. 1. Average wheel running metrics across five days (n = 92).** Values are LS means  $\pm$  standard errors from SAS Procedure Mixed repeated-measures ANCOVAs (see Table S1 for full results). (A) total revolutions C vs. HR. (B) revolutions per minute C vs. HR (average speed). (C) number of one-minute intervals with at least one revolution C vs. HR (running duration). (D) total revolutions of normal vs. mini-muscle (E) revolutions per minute normal vs. mini-


muscle (average speed). (F) number of one-minute intervals with at least one revolution normal vs. mini-muscle (running duration). HR mice ran significantly more total revolutions per day, at higher average speeds, and for a longer duration than C mice. The day-to-day increase in distance run and average speed was significantly greater for HR than for C mice. In addition, mini-muscle individuals ran more distance and faster than normal-muscled individuals. The duration of activity varied significantly across days, such that all groups tended to run more on day 1, which probably represents a novelty effect.



**Fig. 2. Average home-cage activity metrics across five days for C vs. HR mice (left column) and Normal vs. Mini-muscle mice (right column).** Analyses of mice with and without wheel access were performed separately. Values are LS means  $\pm$  standard errors from SAS Procedure Mixed used to implement repeated-measures ANCOVAs. (A) Total HCA (n = 191). (B) Time spent active (n = 190). (C) Mean activity per minute (n = 191). (D) Mean maximum activity in any one-minute interval (n = 191). (E) Total HCA (n = 191). (F) Time spent active (n = 190). (G) Mean activity per minute (n = 191). (H) Mean maximum activity in any one-minute interval (n = 191). HR mice had higher total activity, mean activity per minute, and maximum activity in any one-minute interval compared to C mice when housed without access to wheels, and they were active for more minutes regardless of wheel access. Mini-muscle individuals had less total activity, less activity per minute, and less maximum activity in any one-minute interval than

normal-muscled mice when housed without access to wheels. Mini-muscle mice were also active for fewer minutes than normal-muscled mice when housed with wheels.





**Fig. 3. Buffering capacity ( $\beta_m$ ) of right thigh muscles.** Values are LS means  $\pm$  standard errors from SAS Procedure Mixed, based on  $\log_{10}$ -transformed data. (A) Analyses of all mice combined ( $n = 176$ ) showing male and female, Control and HR mice, with or without access to running wheels, as well as normal vs. mini-muscle. Female mice had significantly lower buffering capacity than males regardless of wheel access or linetype ( $p = 0.0051$ , Table S3).

Control and HR mice did not significantly differ in buffering capacity. Mini-muscle mice had lower buffering capacity than normal mice. Panels B-E represent separate analyses of one fourth of the total sample size. (B) Normal-muscled mice versus mini-muscle female mice without wheels ( $n = 43$ ,  $p = 0.0206$ ), (C) normal vs. mini-muscled male mice without wheels ( $n = 45$ ,  $p = 0.0413$ ), (D) normal vs. mini-muscled female mice with wheels ( $n = 46$ ,  $p = 0.0084$ ), and (E) normal vs. mini-muscled male mice with wheels ( $n = 42$ ,  $p = 0.0004$ ). Mini-muscle mice had significantly lower buffering capacity than normal-muscled mice regardless of sex or wheel access.

## Supplementary Table Abbreviations

<u>Abbreviation</u>	<u>Description</u>
RUN	Mean wheel revolutions across days 1-5
INT	Mean number of intervals with at least 1 wheel revolution across days 1-5
RPM	Mean speed across days 1-5
MAX	Mean maximal revolutions in any one-minute interval across days 1-5
HTOT15	Mean home-cage activity across days 1-5
HINT15	Mean active intervals across days 1-5
HMAX15	Mean maximal activity in any one-minute interval across days 1-5
HAPM15	Mean activity per minute across days 1-5
HPER15	Mean active intervals/total intervals across days 1-5
ECRUN15X	Mean wheel revolutions across days 1-5
ECINT15X	Mean number of intervals with at least 1 wheel revolution across days 1-5
ECRPM15X	Mean speed across days 1-5
ECMAX15X	Mean maximal revolutions in any one-minute interval across days 1-5
SUMOF8	Sum of wheel revolutions during the 8 minutes prior to sacrifice
SUMOF20	Sum of wheel revolutions during the 20 minutes prior to sacrifice
SUMOF30	Sum of wheel revolutions during the 30 minutes prior to sacrifice
SUMOF120	Sum of wheel revolutions during the 120 minutes prior to sacrifice

**Table S1.** Significance levels (p values) from repeated-measures ANCOVA analyzing wheel running metrics across 5 days.

Trait	N	PLinetype	PSex	PDay	PSex*Linetype	PDay*Linetype	PSex*Day	PSex*Linetype*Day	PMini	PMini*Day	PMini*Sex	PMini*Sex*Day	PAge	PFreeness
RUN	92	<b>0.0068+</b>	0.3580+	<.0001	0.2431	<.0001	0.3825	0.7147	<b>0.0119+</b>	<b>0.0047</b>	0.5560	0.2231	0.8848+	<b>0.0388+</b>
INT	92	<b>0.0157+</b>	<b>0.0391-</b>	<.0001	0.2440	0.4498	0.8332	0.9544	0.2273-	<b>0.0201</b>	0.1401	0.3871	0.1966-	0.2774+
RPM	92	<b>0.0029+</b>	0.7736+	<.0001	0.1249	<.0001	0.4908	0.0683	<b>0.0240+</b>	0.0514	0.1172	0.5371	0.3665+	0.5006+

Bold values indicate significant differences ( $p < 0.05$ ). Positive (+) indicates direction HR > C, Male > Female, and Mini > Normal.

**Table S2.** Significance levels (p values) from repeated-measures ANCOVAs analyzing home-cage activity (HCA) metrics across 5 days, performed separately for mice housed without or with wheels.

Trait	Wheels	N	Transform	P <sub>Linetype</sub>	P <sub>Sex</sub>	P <sub>Day</sub>	P <sub>Sex*Linetype</sub>	P <sub>Day*Linetype</sub>	P <sub>Sex*Day</sub>	P <sub>Sex*Linetype*Day</sub>	P <sub>Mini</sub>	P <sub>Mini*Day</sub>	P <sub>Mini*Sex</sub>	P <sub>Mini*Sex*Day</sub>	P <sub>Age</sub>	P <sub>Sensitivity</sub>
HTOT	0	95	log <sub>10</sub>	<b>0.0015+</b>	<b>0.0133-</b>	<b>&lt;.0001</b>	0.4590	0.1516	0.3095	0.8637	<b>0.0061-</b>	0.7419	0.0724	0.8083	0.6316-	0.0812-
	1	96	log <sub>10</sub>	0.3958+	0.2878-	<b>&lt;.0001</b>	0.7974	0.3667	0.5077	0.4791	0.7926-	0.0247	0.6725	0.6447	0.8199-	0.3092+
HINT	0	95	-	<b>0.0235+</b>	<b>0.0134-</b>	<b>&lt;.0001</b>	0.7105	0.6372	0.4304	0.9179	0.2489-	0.6405	0.0875	0.9186	0.0709+	0.2207-
	1	95	-	<b>0.0402+</b>	0.4089-	<b>&lt;.0001</b>	0.6007	0.5045	0.2511	0.9162	<b>0.0044-</b>	0.3701	0.0847	0.6688	0.5116+	0.5394-
HAPM	0	95	Rank	<b>0.0007+</b>	<b>0.0410-</b>	<b>&lt;.0001</b>	0.2823	0.2652	0.3310	0.8923	<b>0.0002-</b>	0.6581	0.1681	0.3558	<b>0.0492-</b>	0.0642-
	1	96	Rank	0.8296-	0.9096+	<b>&lt;.0001</b>	0.7555	0.7012	0.6064	0.4359	0.9954+	0.0563	0.8033	0.6064	0.3748-	0.0503+
HMAX	0	95	Rank	<b>0.0018+</b>	<b>0.0207-</b>	<b>0.0011</b>	0.3083	0.4637	0.7107	0.9928	<b>0.0007-</b>	0.5962	<b>0.0114</b>	0.9457	0.3745-	0.3008-
	1	96	Rank	0.8712-	0.9517-	<b>0.0001</b>	0.3336	0.5087	0.6800	0.4055	0.5120+	0.9149	0.5545	0.8103	0.5087-	0.3119+

Bold values indicate significant differences ( $p < 0.05$ ). Positive (+) indicates direction HR > C, Male > Female, and Mini > Normal. For Wheels, 0 = no access, 1 = access. Analyses for mice with and without wheel access were performed separately. Note that the measure of sensor sensitivity was never statistically significant.

**Table S3.** Significance levels (p values) from ANCOVA analyzing buffering capacity.

Trait	N	P <sub>WhlAcc</sub>	P <sub>Sex</sub>	P <sub>Linetype</sub>	P <sub>WhlAcc*Sex</sub>	P <sub>WhlAcc*Linetype</sub>	P <sub>Sex*Linetype</sub>	P <sub>WhlAcc*Sex*Linetype</sub>	P <sub>Mini</sub>	P <sub>HCA</sub>	P <sub>WheelRun</sub>
All mice (as shown in Fig 3A)	176	0.2217-	<b>0.0051+</b>	0.4864+	0.2860	0.3449	0.8344	0.5210	<b>&lt;.0001-</b>	-	-
All mice w HTOT15	176	0.1830-	<b>0.0085+</b>	0.4188+	0.3265	0.2932	0.7651	0.4772	<b>&lt;.0001-</b>	0.5017-	-
All mice w HINT15	176	0.2406-	<b>0.0067+</b>	0.5055+	0.2866	0.3493	0.8361	0.5278	<b>&lt;.0001-</b>	0.9092+	-
All mice w HMAX15	176	0.1348-	<b>0.0061+</b>	0.4542+	0.2797	0.2914	0.7374	0.5333	<b>&lt;.0001-</b>	0.3079-	-
All mice w HAPM15	176	0.1454-	<b>0.0074+</b>	0.3836+	0.3284	0.2486	0.7328	0.4655	<b>&lt;.0001-</b>	0.3155-	-
All mice w HPER15	176	0.2413-	<b>0.0067+</b>	0.5058+	0.2864	0.3495	0.8360	0.5276	<b>&lt;.0001-</b>	0.9081+	-
All mice w ECRUN15X	173	0.5045+	<b>0.0072+</b>	0.1576+	0.2087	0.7004	0.7324	0.6018	<b>&lt;.0001-</b>	-	0.1319-
All mice w ECINT15X	173	0.8644-	<b>0.0060+</b>	0.3998+	0.2797	0.4607	0.8875	0.4864	<b>&lt;.0001-</b>	-	0.7640-
All mice w ECRPM15X	173	0.2926+	<b>0.0062+</b>	0.1176+	0.2361	0.5494	0.6432	0.6992	<b>&lt;.0001-</b>	-	0.0762-
All mice w ECRUN15X and HTOT15	173	0.6783+	<b>0.0114+</b>	0.1437+	0.2421	0.8162	0.6755	0.5533	<b>&lt;.0001-</b>	0.5166-	0.1490-
All mice w ECINT15X and HINT15	173	0.9376-	<b>0.0068+</b>	0.4145+	0.2738	0.4956	0.8930	0.4960	<b>&lt;.0001-</b>	0.7991+	0.7184-
All mice w ECRPM15X and HAPM15	173	0.5030+	<b>0.0095+</b>	0.0873+	0.2750	0.7259	0.5372	0.6268	<b>&lt;.0001-</b>	0.2271-	0.0731-
All mice w ECRUN15X and HAPM15	173	0.8748+	<b>0.0107+</b>	0.1241+	0.2445	0.9145	0.6286	0.5354	<b>&lt;.0001-</b>	0.2597-	0.1425-
All mice w ECRUN15X and HINT15	173	0.4489+	<b>0.0079+</b>	0.1616+	0.2012	0.6517	0.7252	0.6345	<b>&lt;.0001-</b>	0.6557+	0.1160-
All mice w ECRPM15X and HTOT15	173	0.3746+	<b>0.0109+</b>	0.0989+	0.2758	0.6304	0.5692	0.6380	<b>&lt;.0001-</b>	0.4066-	0.0729-
All mice w ECINT15X and HTOT15	173	0.7218-	<b>0.0097+</b>	0.3413+	0.3263	0.3773	0.8064	0.4441	<b>&lt;.0001-</b>	0.4587-	0.8043-

**Table S3 continued.** Significance levels (p values) from ANCOVA analyzing buffering capacity

Trait	N	P <sub>WhlAcc</sub>	P <sub>Sex</sub>	P <sub>Linetype</sub>	P <sub>WhlAcc*Sex</sub>	P <sub>WhlAcc*Linetype</sub>	P <sub>Sex*Linetype</sub>	P <sub>WhlAcc*Sex*Linetype</sub>	P <sub>Mini</sub>	P <sub>HCA</sub>	P <sub>WheelRun</sub>
No Wheels	90	-	<b>0.0412+</b>	0.4207+	-	-	0.2782	-	<b>0.0177-</b>	-	-
No Wheels w HTOT15	90	-	<b>0.0424+</b>	0.6917+	-	-	0.3323	-	<b>0.0317-</b>	0.6306+	-
No Wheels w HINT15	90	-	<b>0.0263+</b>	0.7785+	-	-	0.3074	-	<b>0.0306-</b>	0.1795+	-
No Wheels w HMAX15	90	-	<b>0.0360+</b>	0.7165	-	-	0.3453	-	0.0520-	0.4268+	-
No Wheels w HAPM15	90	-	<b>0.0467+</b>	0.5898+	-	-	0.3021	-	<b>0.0281-</b>	0.8960+	-
No Wheels w HPER15	90	-	<b>0.0263+</b>	0.7764+	-	-	0.3058	-	<b>0.0295-</b>	0.1822+	-
Wheels	87	-	<b>0.0348+</b>	0.7093+	-	-	0.8390	-	<b>&lt;.0001-</b>	-	-
Wheels w HTOT15	87	-	0.0600+	0.6527+	-	-	0.8926	-	<b>0.0001-</b>	0.1004-	-
Wheels w HINT15	87	-	0.0788+	0.5693+	-	-	0.7840	-	<b>&lt;.0001-</b>	0.1558-	-
Wheels w HMAX15	87	-	<b>0.0410+</b>	0.7295+	-	-	0.9634	-	<b>&lt;.0001-</b>	0.1967-	-
Wheels w HAPM15	87	-	<b>0.0414+</b>	0.7292+	-	-	0.8959	-	<b>0.0001-</b>	0.2352-	-
Wheels w HPER15	87	-	0.0764+	0.5722+	-	-	0.7875	-	<b>&lt;.0001-</b>	0.1688-	-
Wheels w ECRUN15X	84	-	0.0613+	0.1582+	-	-	0.9744	-	<b>&lt;.0001-</b>	-	0.0894-
Wheels w ECINT15X	84	-	<b>0.0407+</b>	0.5857+	-	-	0.7514	-	<b>&lt;.0001-</b>	-	0.8159-
Wheels w ECRPM15X	84	-	<b>0.0441+</b>	0.1016+	-	-	0.8479	-	<b>&lt;.0001-</b>	-	<b>0.0407-</b>
Wheels w ECRUN15X and HTOT15	84	-	0.0915+	0.1882+	-	-	0.9997	-	<b>&lt;.0001-</b>	0.1302-	0.1535-
Wheels w ECINT15X and HINT15	84	-	0.0638+	0.5778+	-	-	0.7284	-	<b>&lt;.0001-</b>	0.8038+	0.1890-
Wheels w ECRPM15X and HAPM15	84	-	0.0562+	0.0956+	-	-	0.7624	-	<b>&lt;.0001-</b>	0.1388-	<b>0.0368-</b>
Wheels w ECRUN15X and HAPM15	84	-	0.0724+	0.1758+	-	-	0.9644	-	<b>&lt;.0001-</b>	0.1958-	0.1115-
Wheels w ECRUN15X and HINT15	84	-	0.0926+	0.1804+	-	-	0.8907	-	<b>&lt;.0001-</b>	0.3595-	0.1587-
Wheels w ECRPM15X and HTOT15	84	-	0.0828+	0.0830+	-	-	0.7641	-	<b>&lt;.0001-</b>	0.0593-	<b>0.0326-</b>
Wheels w ECINT15X and HTOT15	84	-	0.0629+	0.5826+	-	-	0.8391	-	<b>&lt;.0001-</b>	0.0833+	0.9897-
Wheels w SUMOF8**0.4	84	-	0.0624+	0.3280+	-	-	0.7500	-	<b>&lt;.0001-</b>	-	<b>0.0440-</b>
Wheels w TSUMOF20**0.4	84	-	<b>0.0397+</b>	0.3789+	-	-	0.9917	-	<b>&lt;.0001-</b>	-	0.2178-
Wheels w TSUMOF30**0.4	84	-	<b>0.0352+</b>	0.4207+	-	-	0.9266	-	<b>&lt;.0001-</b>	-	0.3229-
Wheels w TSUMOF120**0.4	84	-	<b>0.0370+</b>	0.3389+	-	-	0.9289	-	<b>&lt;.0001-</b>	-	0.2814-

Bold values indicate significant differences ( $p < 0.05$ ). Positive (+) indicates direction HR > C, Male > Female, and Mini > Normal. All analyses included age and time of day as covariates (results not shown).