

The role of parasitism in the energy management of a free-ranging bird

Olivia Hicks¹ *, Sarah J. Burthe², Francis Daunt², Mark Newell², Olivier Chastel³,
Charline Parenteau³, Jonathan A. Green¹

¹ School of Environmental Sciences, University of Liverpool, Liverpool L69 3GP, UK

² Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK

³ Centre d'Etudes Biologiques de Chizé, UMR 7372, CNRS-Université La Rochelle, France

*corresponding author och@liv.ac.uk

Keywords

Accelerometry, Daily energy expenditure, Immune costs, Resting metabolic rate, Thyroid hormones, Trade-off

Summary Statement

Demonstration of the sub-lethal costs of endo-parasites to maintenance costs in a wild population of seabirds

Abstract

Parasites often prompt sub-lethal costs to the host by eliciting immune responses. These costs can be hard to quantify but are crucial to our understanding of their host's ecology. Energy is a fundamental currency to quantify these costs, as energetic trade-offs often exist between key fitness-related processes. Daily energy expenditure (DEE) comprises of resting metabolic rate (RMR) and energy available for activity which are linked via the energy management strategy of an organism. Parasitism may play a role in the balance between self-maintenance and activity, as immune costs can be expressed in elevated RMR. Therefore, understanding energy use in the presence of parasitism enables mechanistic elucidation of potential parasite costs. Using a gradient of natural parasite load and proxies for RMR and DEE in a wild population of breeding European shags (*Phalacrocorax aristotelis*), we tested the effect of parasitism on maintenance costs as well as the relationship between proxies for RMR and DEE. We found a positive relationship between parasite load and our RMR proxy in females but not males and no relationship between proxies for RMR and DEE. This provides evidence for increased maintenance costs in individuals with higher parasite loads and suggests the use of an allocation energy management strategy, whereby an increase to RMR creates restrictions on energy allocation to other activities. This is likely to have fitness consequences as energy allocated to immunity is traded-off against reproduction. Our findings demonstrate that understanding energy management strategies alongside fitness drivers is central to understanding the mechanisms by which these drivers influence individual fitness.

Introduction

Parasites incur major fitness consequences to a huge array of taxa (Booth et al., 1993; Gooderham and Schulte-Hostedde, 2011; Reed et al., 2008). Often parasites prompt sub-lethal effects to the host by eliciting immune or stress responses (Lettini and Sukhdeo, 2010; Martin et al., 2003; Sheldon and Verhulst, 1996; Smyth and Drea, 2016). However, the costs of these sub-lethal effects can be hard to quantify and consequently their impact on individuals and populations is often neglected in ecological research (Binning et al., 2017). Additionally, the mechanisms underlying the fitness effects of parasitism and individual variation in these effects are not well understood. Ignoring these effects of parasites reduces our understanding of their host's ecology, as we know parasites can drive a large array of fitness-related traits (Boulinier et al., 2016; Hamilton and Zuk, 1982; Norris and Evans, 2000; Reed et al., 2008; Sheldon and Verhulst, 1996).

Energy can be a limited resource, and individuals are required to allocate energy to the demands of competing life-history traits to maximize fitness (Stearns, 1992). Therefore, trade-offs may exist in energetic terms between key fitness-related behaviours and processes and so energy is a central mechanism by which fitness responses to parasitism are manifested. A fundamental potential trade-off exists between basal or resting metabolic rate (RMR) and the amount of energy apportioned to activity, because ceilings can exist on the sum of these two components, the daily rate of energy expenditure (DEE) (Careau and Garland, 2012; Elliott et al., 2014; Mathot and Dingemanse, 2015; Welcker et al., 2010). For endotherms, RMR is usually defined as the minimum energetic cost of living during thermo-neutral rest in a free ranging animal (Mathot and Dingemanse, 2015; Welcker et al., 2015) and therefore largely represents the cost of self-maintenance, including immune activity (Burton et al., 2011). Remaining energy available to allocate to activity is variously referred to as metabolic scope or activity metabolism (Careau and Garland, 2012; Mathot and Dingemanse, 2015). DEE, RMR and

activity metabolism are linked in different ways depending on which energy management strategy the animal is operating under namely the performance, independent or allocation strategies (see Fig. 1. and appendix for details). A recent review of energy management in birds and mammals suggests that no single energy management strategy operates across all species (Portugal et al., 2016).

Parasitism may play a central role in the relationship between self-maintenance and activity, as a primary predicted cost of parasitism is to RMR, through immune associated costs (Svensson et al., 1998). Hosts should invest in immunity to reduce the impacts of parasitism, particularly in long-lived species, where in contrast to short-lived species, survival contributes more to fitness and individuals should be selected to allocate more resources to a rapid, strong immune response to combat cumulative damage from parasites and to protect future reproductive success (Lochmiller and Deerenberg, 2000; Moe et al., 2007). Under the allocation model of energy management, this investment would create a trade-off between immune activity and other essential fitness-related organismal functions (Råberg et al., 2000; Sheldon and Verhulst, 1996). Under the performance model, there would be no trade-off with increased RMR, but instead activity metabolism would also increase with a resultant relatively large increase in DEE. Likewise under the independent model, despite no relationship between RMR and activity metabolism, an increase in RMR results in an equivalent increase in DEE, since this is the sum of activity metabolism and RMR (Mathot and Dingemanse, 2015; Welcker et al., 2015). The lack of a trade-off would itself likely have implications for fitness, and animals might ultimately reach maximal or optimal limits of DEE (Careau and Garland, 2012; Mathot and Dingemanse, 2015).

Understanding the impacts of natural parasite burdens on individual metabolic rate is crucial, with experimental studies suggesting endo-parasites and elevated immune function incur significant costs to hosts (Albon et al., 2002; Newborn and Foster, 2002; Sheldon and Verhulst,

1996). However there has been little work on the role of parasitism in mediating the energy management of individuals despite this being fundamental to the understanding of the fitness effects of parasitism. This is in large part due to the difficulties in measuring natural parasite loads, particularly endo-parasites, in wild populations. Energetic rates are also hard to measure in free-ranging animals and so various proxies are used to evaluate energetic costs (Butler et al., 2004; Chastel et al., 2003; Halsey et al., 2011)

Here we measure proxies of both DEE and RMR in individuals with known natural endo-parasite loads in a free-living population of breeding European shags (*Phalacrocorax aristotelis*) to understand the effect of endo-parasites on maintenance costs as well as the relationship between maintenance and total energy expenditure during the breeding period under high energetic demands. Specifically, we investigate two questions: 1. Does parasite load relate to RMR? We expect that individuals with higher parasite loads will have higher immune costs and that this will be reflected in elevated RMR 2. Does DEE relate to RMR and what can this tell us about the allocation between energy to reproduction and self-maintenance of this species? We predict that individuals will be energetically constrained by rapidly growing chicks during the chick rearing period under the allocation model and therefore any increase in RMR will not result in an increase in DEE.

Materials and Methods

Study site and species

The study was carried out at the Isle of May National Nature Reserve, south-east Scotland (56°11'N, 2°33'W) during the breeding seasons of 2015-2017. All individuals were part of a long-term population study and are individually marked with a metal and a unique plastic ring for identification and accurately aging individuals that were first ringed as chicks.

Previous sampling of this population has shown a high prevalence of the nematode parasite *Contracaecum rudolphii* (Burthe et al., 2013; Granroth-Wilding et al., 2014; Reed et al., 2008) in European shags, though parasite loads vary markedly between individuals (Burthe et al., 2013; Granroth-Wilding et al., 2014; Reed et al., 2012). Shags become infected with third stage larvae via their fish diet. Larval worms moult to become sexually mature adults which attach to the lining of the proventriculus and lower oesophagus in the final seabird host (Abollo et al., 2001; Burthe et al., 2013).

Estimating resting metabolic rate

Measuring resting metabolic rate in the field is difficult (Wilson et al., 2006). Our previous work using calibrated estimates of resting metabolic rate (RMR) from accelerometry (Hicks et al 2017) suggested that RMR of European shags is linked to parasite burden (Hicks et al 2018). However, despite evidence that movement costs from accelerometry can be indeed be used to evaluate costs during relative inactivity (Green et al., 2009), we wanted to use a more sensitive proxy for RMR to investigate this effect further. Thyroid hormone plasma concentrations are increasingly being used in this context in free ranging animals. The role of thyroid hormones in energy metabolism in birds in particular is now well established (Chastel et al., 2003; Elliott et al., 2013; Welcker et al., 2013). Triiodothyronine hormone (T3) especially is considered one of the major controllers for the regulation of tissue oxygen consumption and metabolic activity

in endotherms (McNab, 1997). Studies carried out both in the laboratory and the field show close relationships for a range of species and life-history strategies (Chastel et al., 2003; Duriez et al., 2004; Elliott et al., 2013; Vézina et al., 2009; Welcker et al., 2013; Zheng et al., 2014). While a direct calibration of T3 hormone and RMR has yet to be conducted for our study species, we are unaware of any calibration studies during breeding which have failed to establish such a relationship. Thus we feel confident that T3 hormone can be used as a proxy for RMR at the individual level, without the confounding effect of stress associated with respirometry (Welcker et al., 2015).

Adult European shags were captured on the nest during chick rearing (when the chicks were between 5 and 36 days old) using a crook on the end of a long pole. At capture a 1 mL blood sample was collected from the brachial vein using a heparinized syringe and a 25-gauge needle. To avoid an effect of handling stress on hormone concentrations, sampling took place within 3 min of capture. Blood samples were taken between 03:30 and 07:30, before shags left for their first foraging trip of the day, and therefore birds were assumed to be post-absorptive, which was later confirmed by endoscopy. Throughout the fieldwork period, local air temperature was comfortably above the estimated lower critical temperature of 6°C for European shags (Enstipp et al., 2005), thus birds were in thermoneutrality at all times. Blood samples were stored on ice in the field. Whole blood was centrifuged and plasma and red blood cells were kept frozen at -20 °C until subsequent analyses.

Total 3, 3'-Triiodo-L-Thyronine (T3) hormone assays were performed at the Centre d'Etudes Biologiques de Chizé (CEBC), France. T3 analyses were performed using a single radioimmunoassay (RIA) and Total Thyroid hormone levels were assessed in duplicate without extraction as in Chastel et al., (2003) and Welcker et al., (2013). 25 µL of plasma was incubated with 10000 cpm of 125I-hormone (Perkin Elmer, US) and polyclonal rabbit antiserum (Sigma Aldrich, US). The bound fraction was separated from the free fraction by addition of a sheep

anti-rabbit antibody. After overnight incubation and centrifugation, the bound fraction activity was then counted on a Wizard 2 gamma counter (Perkin Elmer, US). For quality control, reference materials were used and were within the acceptable range determined by the laboratory. Furthermore, every plasma was run in duplicates, samples with a coefficient of variance (CV) higher than 12% were re-assayed. Inter- and intra-assay variations were respectively 14.34% and 7.75%. Cross-reactions of T3 antiserum were as follows: triiodoD-thyroacetic acid 6%, L-thyroxine 0.2%, diiodo-L-thyrosine <0.01%, monoiodo-L-thyrosine <0.01%. Cross-reactions of T4 antiserum were as follows: triiodothyronine 4%, diiodo-L-thyrosine <0.01%, monoiodo-L-thyrosine <0.01%.

Quantifying parasite load

Worms were counted visually using the endoscope video screen (for detailed endoscopy methods see Burthe et al. (2013)). Burdens higher than 40 worms were hard to quantify in the field accurately and these were counted retrospectively via endoscopy video footage. Quantification of parasite burdens was found to be repeatable within an individual across a season (Burthe et al. 2013). All endoscopy and blood sampling was performed by trained personnel (S. Burthe) holding a personal licence, and under a project licence issued by the UK Home Office.

Estimating daily energy expenditure

Tri-axial accelerometers (little Leonardo D3GT, AXY3 and Gulf Coast Data Concepts X8) were used to estimate daily energy expenditure of a subset of the birds used for the hormone assay. Accelerometers were set to record at 25 or 50 Hz and attached on the midline of the mid back of individuals (as close to the centre of gravity as possible) using Tesa tape. All birds were successfully recaptured and accelerometers were retrieved after three days of deployment. Energy expenditure was estimated for diving, flying and resting (the main activities of

European shags (Sakamoto et al., 2009) using behaviour specific calibrations derived from respirometry calibrations with heart rate and accelerometry (see Hicks et al., 2017 for behaviour specific calibrations and detailed methods). These data were then used to estimate total daily energy expenditure (DEE), the sum of the energetic costs of all behavioural bouts within 24-hour periods of activity which included both activity and RMR and any costs of thermoregulation in water (see Hicks et al 2018 for full details). Adult RMR and DEE have been found to vary with mass and reproductive stage and age in many seabirds (Elliott et al., 2014; Green et al., 2013; Grémillet, 1997; Weimerskirch et al., 1995). Thus, adult mass and age were recorded at endoscopy and chick age at time of sampling was later back-calculated from wing length at ringing (Granroth-Wilding et al., 2014).

Statistical Analysis

All models were fitted separately for males and females due to non-independence of nest pairs. To answer the first question of whether parasite load is related to elevated RMR, we modelled T3 concentration using linear mixed effects models (LMMs). Parasite load, brood size (number of chicks), brood age (age of the oldest chick in the brood at the time of sampling), year and adult age and body mass were explanatory variables and we controlled for variation between birds and repeated sampling by including individual as a random factor. We fitted models containing all combinations of the fixed effects, and interactions between parasite load and brood age and parasite load and brood size (see Table 1 for explanations of model terms).

Secondly to understand how variation in RMR is related to DEE, we modelled estimated DEE as a function of T3 concentration (a proxy for RMR) using linear mixed effects models (LMMs). Adult age, adult body mass, brood age (at the time of sampling) brood size and year were also included as explanatory variables (Table 1). We controlled for variation between birds and repeated sampling by including individual as a random factor.

All models were fitted using the *lme4* package in R (Bates et al., 2014; R Core Team, 2015) and model selection for both sets of models was based on Akaike's information criterion (AIC) (Burnham and Anderson, 2001).

Results

Relationship between parasite load and RMR

A total of 87 (48 males and 39 females, 2015:8, 2016:42, 2017:37) individuals were blood sampled and endoscoped over three years (of which 27 were sampled in more than one year). Whilst the sample size in 2015 is clearly smaller than the other years, removal of these data had no impact on the conclusions of the analyses therefore they were included. The best supported model for females showed a positive relationship between parasite load and T3 concentration, an effect of year, and an interaction between brood age and parasite load in that there was a more positive relationship between parasite load and T3 when broods were younger (See Fig 2 and Table 2). When year and brood age are accounted for, T3 concentration increased by 150% across the range in natural parasite load. The best supported model for males showed no effect of parasite load on T3 concentration. There was a negative effect of age as well as an effect of year on T3 concentration in males (see Fig 3 and Table 2).

Relationship between proxies for RMR and DEE

Daily energy expenditure was estimated in a subset of the birds (71 individuals; 2015=3, 2016=40, 2017=28) to gain insight into the relationship between RMR and DEE. The best supported model for females was the null model with no relationship found between T3 and estimated DEE (see Fig 4 and Table 3). For males, the best supported model indicated a

negative effect of body mass on estimated DEE and no effect of T3 (see Fig 4 and Table 3). However, the null model is also equally well supported (within $\Delta 2$ AIC of the top model) thus there is very little support for an effect of mass.

Discussion

Measuring both natural parasite burdens and metabolic rates in free-ranging animals is challenging. In particular, estimating metabolic rate requires the use of proxies, the potential limitations of which have been extensively documented (e.g. Butler et al., 2004; Halsey et al., 2011). Our study builds on previous work (Hicks et al 2018) to further explore links between RMR and parasite loads in European shags and how these might link to DEE. Despite the potential limitations, we found a positive relationship between natural parasite load and our hormonal proxy for resting metabolic rate (RMR) across multiple years in female shags, strongly suggesting increased maintenance costs in individuals with higher parasite loads. Host maintenance costs are likely to be increased in response to parasitism due to increased cell repair and cost of dealing with the parasite infection as well as the immune response itself (Lochmiller and Deerenberg, 2000). The limitations of the approaches used would have made it difficult to distinguish between the independent and performance strategies, since our proxies for RMR and daily energy expenditure (DEE) use different units of measurement. However the lack of relationship between these two proxies provides strong evidence that female European shags operate under an allocation energy management strategy. Therefore a heightened immune response to parasitism creates restrictions on the allocation of energy to other activities and is likely associated with subsequent fitness consequences.

We found no relationship between our proxy for RMR and parasitism in males despite the strong relationship found in females. Males are hypothesised to experience the immunosuppressive qualities of testosterone (Grossman, 1985), thus the lack of relationship may be due to suppression of their parasite-induced immune response. Additionally, trade-offs between reproduction and self-maintenance are especially strong in female shags (Hicks et al., 2018; Reed et al., 2008), who have higher estimated DEE than males on average (see Fig 3. and Hicks et al., 2018) and can be more adversely affected by environmental conditions than males (Lewis et al., 2015). Females across many taxa often invest more in reproduction than males (Clutton-Brock, 1991) meaning that females may be closer to their energy expenditure limits than males. Thus the greater energetic constraint of females to extrinsic and intrinsic conditions could explain these sex differences in response to parasitism (Hicks et al., 2018; Lewis et al., 2015).

This study is the first to use T3 concentration as a proxy for RMR in the context of parasitism effects. This enables us to gain greater insight into the energy use of free-ranging animals and complements our previous findings and other approaches. Hicks et al. (2018) show that for female European shags, flight costs are positively associated with parasite burden but that the proportion of time spent in flight per day reduced with higher parasite loads. However this decrease in flight duration was more than expected based on the increased cost of flight alone. The results from this study suggest that the reduction in flight duration observed in heavily parasitised birds may be due to compensation for increased maintenance costs as well as the increased cost of flight (Hicks et al., 2018). For European shags it is not possible to quantify the extent of this compensation as we have yet to calibrate T3 concentration against RMR using respirometry. However T3 plasma concentration is increasingly being used as a proxy for RMR due to strong correlations found in multiple species and the reduction in the confounding effect of stress associated with respirometry (Blackmer et al., 2005; Bouwhuis et al., 2011; Chastel

et al., 2003; Elliott et al., 2013; Moe et al., 2007; Welcker et al., 2013; Welcker et al., 2015). In an experimental study, a 50% increase in free T3 corresponded to a 50% increase in RMR in kittiwakes (Welcker et al., 2015). In this study, we found a 150% change in T3 concentration across the natural range of parasite load, meaning that the corresponding change in RMR to parasitism could be even greater than in previous experimental work.

It is likely that energy management is not fixed across time but varies according to particular conditions as seen in Hicks et al., (2018) where estimated DEE increased with brood age but not in relation to parasite load. An adult's cumulative energetic investment in its brood increases with brood age (Drent and Daan, 1980). Thus the trade-off between allocation to reproduction and self-maintenance may also shift with brood age. In a young brood, when cumulative investment is lower than an older brood energy allocation to immune response may be beneficial to ensure that the next reproductive event is reached (Ilmonen et al., 2000), but may reduce parental effort (Råberg et al., 2000) and negatively affect offspring success. However when cumulative investment in a brood is higher, investment to current reproduction may be prioritised. Consistent with this concept, we found the positive relationship between RMR and parasite load is greater when the adult's chicks are younger suggesting that the shags are investing more in parasite resistance at this time.

At the scale of a lifetime, life-history theory predicts trade-offs between reproduction and self-maintenance change with age. As the probability of future reproduction declines with age, it is predicted that resources will be increasingly allocated to current reproduction rather than maintenance (Herborn et al., 2015; van Noordwijk and de Jong, 1986). There is good evidence for this in short-lived species such as great tits and zebra finch (Bouwhuis et al., 2011; Moe et al., 2009), however in longer lived birds evidence is equivocal e.g. (Blackmer et al., 2005; Elliott et al., 2015; Moe et al., 2007). In this study we found a negative relationship between age and RMR in males, suggesting older individuals invest less energy in maintenance as in

thick billed murre (Elliott et al., 2015). However, we found no effect of age on RMR in females. Sex-specific senescence occurs in a number of wild vertebrate populations, possibly due to different trade-offs existing with age in males and females (Clay et al., 2018; Froy et al., 2013; Froy et al., 2017; Murgatroyd et al., 2018).

We found no evidence for a relationship between proxies for RMR and DEE indicating that DEE does not vary with maintenance costs. Despite parasite induced behavioural costs, European shag estimated DEE does not correlate with parasite load (Hicks et al., 2018) suggesting that there is a level of energy expenditure for this population that they are unwilling or unable to exceed. Drent and Daan (1980) first introduced the concept of an ‘Optimal Working Capacity’ as a rate beyond which parents would increasingly suffer from any kind of risks or dangers with the consequence that their lifetime reproductive success would be reduced (Green et al 2009). Evidence for such an optimum, occasionally described as an ‘energetic ceiling’, have been described previously (e.g. Welcker et al., (2010) and provide evidence for the existence of allocation management strategies. Our results also provide evidence for both an optimum to energy expenditure and the allocation management strategy in European shags (Drent and Daan, 1980; Mathot and Dingemans, 2015). As such, an increase in RMR will cause a reduction in activity metabolism, which may have consequences for fitness-related behaviours such as chick provisioning or foraging. Our findings are in agreement with studies providing evidence for the allocation management strategy in three seabird species (Blackmer et al., 2005; Elliott et al., 2014; Welcker et al., 2015) but are contrary to other studies (including those of other seabird species) providing evidence for both performance and independent strategies (Chastel et al., 2003; Portugal et al., 2016).

It is unclear why species vary in their energy management strategy, though during energetically demanding periods, the capacity to increase expenditure can impose energy limits, in terms of foraging constraints, making an allocation strategy more likely (Elliott et al., 2014). Under

limited energy it can be difficult to buffer the additional costs of parasites. Therefore, considering the role of parasitism in mediating energy allocation between reproduction and self-maintenance is crucial in understanding the mechanism of its fitness effects. The allocation strategy creates energetic trade-offs under an overarching constraint, yet before now we had little knowledge about what drives these trade-offs. Changes in energy allocation can influence reproduction in the short term, but in the long term there may be negative consequences of reducing allocation to self-maintenance (Blackmer et al., 2005). Therefore, quantifying energy management strategies alongside potential fitness drivers is crucial to understand the mechanisms by which they act and influence energy allocation to optimise individual success.

Authors' contributions

OH, SB, FD and JG designed the study. OH, SB, FD and MN collected the data. OC and CP performed the hormone assays and helped with the interpretation of the data. OH processed the accelerometry data, conducted the statistical analyses and wrote the manuscript. All authors (OH, SB, FD, MN, OC, CP and JG) contributed to interpreting results and improvement of this paper.

Acknowledgements

We are grateful to everyone involved in the data collection in particular Carrie Gunn for her invaluable help in the endoscopy procedure and Scottish Natural Heritage for allowing us to work on the Isle of May. We thank Colette Trouvé at CEBC, for her help in the hormone assays and Alice Carravieri for helpful comments to improve the manuscript

Data accessibility

Data and code are uploaded to figshare and will be made available at time of publication

Competing interests

We declare we have no competing interest

Funding

OH was funded by a PhD studentship from the Natural Environment Research Council (NERC). Accelerometry data collection was made possible by funding for data loggers by the Japanese Society for the Promotion of Science, UTBLS: Bio-Logging Science, University of Tokyo. Endoscopy work was funded by the NERC National Capability and undertaken under Home Office licence PPL70/8107 snD SNH research licence.

Appendix

Energy management strategies

Performance model: Variation in RMR reflects variation in the size of organs that mobilize energy (e.g., digestive organs, muscle). Therefore, individuals with higher maintenance MR are able to maintain higher levels of energy output (i.e., have higher DEE). This strategy predicts higher RMR to be associated with both greater activity metabolism and therefore when summed, a higher DEE. (Careau et al., 2008).

Allocation model: Variation in RMR does not reflect variation in the size of organs that mobilize energy; differences in RMR are therefore not associated with differences in DEE. Individuals with higher maintenance MR have less energy available to allocate to energetically costly behaviours and so are predicted to have lower expression of such traits (Mathot and Dingemans, 2015).

Independent Model: Regulation of RMR and DEE are independent but an increase in RMR causes an increase in total energy expenditure as activity metabolism is also independent of RMR. Therefore, higher RMR is associated with higher DEE because while activity metabolism is unaffected it sums with an increasing RMR to drive an increased DEE.

References

- Abollo, E., Gestal, C. and Pascual, S.** (2001). Anisakid infection in the European shag *Phalacrocorax aristotelis aristotelis*. *J. Helminthol.* **75**, 209–214.
- Albon, S. D., Stien, A., Irvine, R. J., Langvatn, R., Ropstad, E. and Halvorsen, O.** (2002). The role of parasites in the dynamics of a reindeer population. *Proc. R. Soc. B Biol. Sci.* **269**, 1625–1632.
- Bates, D., Mächler, M., Bolker, B. and Walker, S.** (2014). Fitting Linear Mixed-Effects Models using lme4. *J. Stat. Softw.* **67**,.
- Binning, S. A., Shaw, A. K. and Roche, D. G.** (2017). Parasites and Host Performance: Incorporating Infection into Our Understanding of Animal Movement. *Integr. Comp. Biol.* **57**, 1–14.
- Blackmer, A. L., Mauck, R. A., Ackerman, J. T., Huntington, C. E., Nevitt, G. A. and Williams, J. B.** (2005). Exploring individual quality: Basal metabolic rate and reproductive performance in storm-petrels. *Behav. Ecol.* **16**, 906–913.
- Booth, D. T., Clayton, D. H. and Block, B. A.** (1993). Experimental Demonstration of the Energetic Cost of Parasitism in Free-Ranging Hosts. *Proc. R. Soc. B Biol. Sci.* **253**, 125–129.
- Boulinier, T., Kada, S., Ponchon, A., Dupraz, M., Dietrich, M., Gamble, A., Bourret, V., Duriez, O., Bazire, R., Tornos, J., et al.** (2016). Migration, Prospecting, Dispersal? What Host Movement Matters for Infectious Agent Circulation? *Integr. Comp. Biol.* **56**, 330–342.
- Bouwhuis, S., Sheldon, B. C. and Verhulst, S.** (2011). Basal metabolic rate and the rate of senescence in the great tit. *Funct. Ecol.* **25**, 829–838.

- Burnham, K. P. and Anderson, D. R.** (2001). Kullback-Leiber information as a basis for strong inference in ecological studies. *Wildl. Res.* **28**, 111–119.
- Burthe, S. J., Newell, M. A., Goodman, G., Butler, A., Bregnballe, T., Harris, E., Wanless, S., Cunningham, E. J. A. and Daunt, F.** (2013). Endoscopy as a novel method for assessing endoparasite burdens in free-ranging European shags (*Phalacrocorax aristotelis*). *Methods Ecol. Evol.* **4**, 207–216.
- Burton, T., Killen, S. S., Armstrong, J. D. and Metcalfe, N. B.** (2011). What causes intraspecific variation in resting metabolic rate and what are its ecological consequences? *Proc. R. Soc. B Biol. Sci.* **278**, 3465–73.
- Butler, P. J., Green, J. A., Boyd, I. L. and Speakman, J. R.** (2004). Measuring metabolic rate in the field: the pros and cons of the doubly labelled water and heart rate methods. *Funct. Ecol.* **18**, 168–183.
- Careau, V. and Garland, T.** (2012). Performance, Personality, and Energetics: Correlation, Causation, and Mechanism. *Physiol. Biochem. Zool. Ecol. Evol. Approaches* **85**, 543–571.
- Careau, V., Thomas, D., Humphries, M. M. and Réale, D.** (2008). Energy metabolism and animal personality. *Oikos* **117**, 641–653.
- Chastel, O., Lacroix, A. and Kersten, M.** (2003). Pre-breeding energy requirements: thyroid hormone, metabolism and the timing of reproduction in house sparrows *Passer domesticus*. *J. Avian Biol.* **34**, 298–306.
- Clay, T. A., Pearmain, E. J., McGill, R. A. R., Manica, A. and Phillips, R. A.** (2018). Age-related variation in non-breeding foraging behaviour and carry-over effects on fitness in an extremely long-lived bird. *Funct. Ecol.* **50**, 700.
- Clutton-Brock, T. H.** (1991). *The evolution of parental care*. Princeton University Press.
- Drent, R. and Daan, S.** (1980). The prudent parent: energetic adjustments in avian breeding. *Ardea* **68**, 225–252.
- Duriez, O., PastoutLucchini, L., Boos, M., Chastel, O., Fritz, H., Ferrand, Y. and Clobert, J.** (2004). Low levels of energy expenditure in a nocturnal, forest-dwelling wader, the Eurasian Woodcock *Scolopax rusticola*. *Ardea* **92**, 31–42.
- Elliott, K. H., Welcker, J., Gaston, A. J., Hatch, S. A., Palace, V., Hare, J. F., Speakman, J. R. and Anderson, W. G.** (2013). Thyroid hormones correlate with resting metabolic rate, not daily energy expenditure, in two charadriiform seabirds. *Biol. Open* **2**, 580–586.
- Elliott, K. H., Le Vaillant, M., Kato, A., Gaston, A. J., Ropert-Coudert, Y., Hare, J. F., Speakman, J. R. and Croll, D.** (2014). Age-related variation in energy expenditure in a long-lived bird within the envelope of an energy ceiling. *J. Anim. Ecol.* **83**, 136–46.
- Elliott, K. H., Hare, J. F., Le Vaillant, M., Gaston, A. J., Ropert-Coudert, Y. and Anderson, W. G.** (2015). Ageing gracefully: Physiology but not behaviour declines with age in a diving seabird. *Funct. Ecol.* **29**, 219–228.
- Enstipp, M. R., Grémillet, D. and Lorentsen, S.-H.** (2005). Energetic costs of diving and thermal status in European shags (*Phalacrocorax aristotelis*). *J. Exp. Biol.* **208**, 3451–3461.

- Froy, H., Phillips, R. A., Wood, A. G., Nussey, D. H. and Lewis, S.** (2013). Age-related variation in reproductive traits in the wandering albatross: Evidence for terminal improvement following senescence. *Ecol. Lett.* **16**, 642–649.
- Froy, H., Lewis, S., Nussey, D. H., Wood, A. G. and Phillips, R. A.** (2017). Contrasting drivers of reproductive ageing in albatrosses. *J. Anim. Ecol.* **86**, 1022–1032.
- Gooderham, K. and Schulte-Hostedde, A.** (2011). Macroparasitism influences reproductive success in red squirrels (*Tamiasciurus hudsonicus*). *Behav. Ecol.* **22**, 1195–1200.
- Granroth-Wilding, H. M. V., Burthe, S. J., Lewis, S., Reed, T. E., Herborn, K. A., Newell, M. A., Takahashi, E. A., Daunt, F. and Cunningham, E. J. A.** (2014). Parasitism in early life: Environmental conditions shape within-brood variation in responses to infection. *Ecol. Evol.* **4**, 3408–3419.
- Green, J. A., Halsey, L. G., Wilson, R. P. and Frappell, P. B.** (2009). Estimating energy expenditure of animals using the accelerometry technique: activity, inactivity and comparison with the heart-rate technique. *J. Exp. Biol.* **212**, 471–482.
- Green, J. A., Aitken-Simpson, E. J., White, C. R., Bunce, A., Butler, P. J. and Frappell, P. B.** (2013). An increase in minimum metabolic rate and not activity explains field metabolic rate changes in a breeding seabird. *J. Exp. Biol.* **216**, 1726–1735.
- Grémillet, D.** (1997). Catch per unit effort, foraging efficiency, and parental investment in breeding great cormorants (*Phalacrocorax carbo carbo*). *ICES J. Mar. Sci.* **54**, 635–644.
- Grossman, C. J.** (1985). Interactions between the Gonadal Steroids and the Immune System. *Science* (80-.). **227**, 257–261.
- Halsey, L. G., Shepard, E. L. C. and Wilson, R. P.** (2011). Assessing the development and application of the accelerometry technique for estimating energy expenditure. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **158**, 305–314.
- Hamilton, W. D. and Zuk, M.** (1982). Heritable true fitness and bright birds: A role for parasites? *Science* (80-.). **218**, 384–387.
- Herborn, K. A., Daunt, F., Heidinger, B. J., Granroth-Wilding, H. M. V., Burthe, S. J., Newell, M. A. and Monaghan, P.** (2015). Age, oxidative stress exposure and fitness in a long-lived seabird. *Funct. Ecol.* 1–9.
- Hicks, O., Burthe, S., Daunt, F., Butler, A., Bishop, C. and Green, J. A.** (2017). Validating accelerometry estimates of energy expenditure across behaviours using heart rate data in a free-living seabird. *J. Exp. Biol.* **220**, 1875–1881.
- Hicks, O., Burthe, S. J., Daunt, F., Newell, M., Butler, A., Ito, M., Sato, K. and Green, J. A.** (2018). The energetic cost of parasitism in a wild population. *Proc. R. Soc. B Biol. Sci.* **285**, 20180489.
- Ilmonen, P., Taarna, T. and Hasselquist, D.** (2000). Experimentally activated immune defence in female pied flycatchers results in reduced breeding success. *Proc. R. Soc. B Biol. Sci.* **267**, 665–670.
- Lettini, S. E. and Sukhdeo, M. V. K.** (2010). The energetic cost of parasitism in isopods. *Écoscience* **17**, 1–8.
- Lewis, S., Phillips, R. A., Burthe, S. J., Wanless, S. and Daunt, F.** (2015). Contrasting

- responses of male and female foraging effort to year-round wind conditions. *J. Anim. Ecol.* **84**, 1490–1496.
- Lochmiller, R. L. and Deerenberg, C.** (2000). Trade-Offs in Evolutionary Immunology: Just What Is the Cost of Immunity? *Oikos* **88**, 87–98.
- Martin, L. B., Scheuerlein, A. and Wikelski, M.** (2003). Immune activity elevates energy expenditure of house sparrows: a link between direct and indirect costs? *Proc. R. Soc. B Biol. Sci.* **270**, 153–158.
- Mathot, K. J. and Dingemans, N. J.** (2015). Energetics and behavior: Unrequited needs and new directions. *Trends Ecol. Evol.* **30**, 199–206.
- McNab, B. K.** (1997). On the Utility of Uniformity in the Definition of Basal Rate of Metabolism. *Physiol. Zool.* **70**, 718–720.
- Moe, B., Angelier, F., Bech, C. and Chastel, O.** (2007). Is basal metabolic rate influenced by age in a long-lived seabird, the snow petrel? *J. Exp. Biol.* **210**, 3407–14.
- Moe, B., Ronning, B., Verhulst, S. and Bech, C.** (2009). Metabolic ageing in individual zebra finches. *Biol. Lett.* **5**, 86–89.
- Murgatroyd, M., Roos, S., Evans, R., Sansom, A., Whitfield, D. P., Sexton, D., Reidb, R., Grant, J. and Amar, A.** (2018). Sex-specific patterns of reproductive senescence in a long-lived reintroduced raptor. *J. Anim. Ecol.*
- Newborn, D. and Foster, R.** (2002). Control of parasite burdens in wild red grouse *Lagopus lagopus scoticus* through the indirect application of anthelmintics. *J. Appl. Ecol.* **39**, 909–914.
- Norris, K. and Evans, M. R.** (2000). Ecological immunology: life history trade-offs and immune defense in birds. *Behav. Ecol.* **11**, 19–26.
- Portugal, S. J., Green, J. A., Halsey, L. G., Arnold, W., Careau, V., Dann, P., Frappell, P. B., Grémillet, D., Handrich, Y., Martin, G. R., et al.** (2016). Associations between Resting, Activity, and Daily Metabolic Rate in Free-Living Endotherms: No Universal Rule in Birds and Mammals. *Physiol. Biochem. Zool.* 000–000.
- R Core Team** (2015). R: A language and environment for statistical computing.
- Råberg, L., Nilsson, J., Imonen, P., Stjernman, M. and Hasselquist, D.** (2000). The cost of an immune response: Vaccination reduces parental effort. *Ecol. Lett.* **3**, 382–386.
- Reed, T. E., Daunt, F., Hall, M. E., Phillips, R. A., Wanless, S. and Cunningham, E. J. A.** (2008). Parasite Treatment Affects Maternal Investment in Sons. *Science (80-)*. **321**, 1681–1682.
- Reed, T. E., Daunt, F., Kiploks, A. J., Burthe, S. J., Granroth-Wilding, H. M. V., Takahashi, E. A., Newell, M., Wanless, S. and Cunningham, E. J. A.** (2012). Impacts of parasites in early life: Contrasting effects on juvenile growth for different family members. *PLoS One* **7**,.
- Sakamoto, K. Q., Sato, K., Ishizuka, M., Watanuki, Y., Takahashi, A., Daunt, F. and Wanless, S.** (2009). Can ethograms be automatically generated using body acceleration data from free-ranging birds? *PLoS One* **4**, e5379.
- Sheldon, B. C. and Verhulst, S.** (1996). Ecological immunology: costly parasite defences

and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* **11**, 317–321.

- Smyth, K. N. and Drea, C. M.** (2016). Patterns of parasitism in the cooperatively breeding meerkat: A cost of dominance for females. *Behav. Ecol.* **27**, 148–157.
- Stearns, S. C.** (1992). *The evolution of life histories*. (ed. Stearns, S. C.) OUP Oxford.
- Svensson, E., Råberg, L., Koch, C. and Hasselquist, D.** (1998). Energetic stress, immunosuppression and the costs of an antibody response. *Funct. Ecol.* **12**, 912–919.
- van Noordwijk, A. J. and de Jong, G.** (1986). Acquisition and Allocation of Resources : Their Influence on Variation in Life History Tactics. *Am. Nat.* **128**, 137–142.
- Vézina, F., Gustowska, A., Jalvingh, K. M., Chastel, O. and Piersma, T.** (2009). Hormonal Correlates and Thermoregulatory Consequences of Molting on Metabolic Rate in a Northerly Wintering Shorebird. *Physiol. Biochem. Zool.* **82**, 129–142.
- Weimerskirch, H., Chastel, O. and Ackermann, L.** (1995). Adjustment of parental effort to manipulated foraging ability in a pelagic seabird, the thin-billed prion *Pachyptila belcheri*. *Behav. Ecol. Sociobiol.* **36**, 11–16.
- Welcker, J., Moe, B., Bech, C., Fyhn, M., Schultner, J., Speakman, J. R. and Gabrielsen, G. W.** (2010). Evidence for an intrinsic energetic ceiling in free-ranging kittiwakes *Rissa tridactyla*. *J. Anim. Ecol.* **79**, 205–213.
- Welcker, J., Chastel, O., Gabrielsen, G. W., Guillaumin, J., Kitaysky, A. S., Speakman, J. R., Tremblay, Y. and Bech, C.** (2013). Thyroid Hormones Correlate with Basal Metabolic Rate but Not Field Metabolic Rate in a Wild Bird Species. *PLoS One* **8**,.
- Welcker, J., Speakman, J. R., Elliott, K. H., Hatch, S. A. and Kitaysky, A. S.** (2015). Resting and daily energy expenditures during reproduction are adjusted in opposite directions in free-living birds. *Funct. Ecol.* 250–258.
- Wilson, R. P., White, C. R., Quintana, F., Halsey, L. G., Liebsch, N., Martin, G. R. and Butler, P. J.** (2006). Moving towards acceleration for estimates of activity-specific metabolic rate in free-living animals: the case of the cormorant. *J. Anim. Ecol.* **75**, 1081–1090.
- Zheng, W.-H., Liu, J.-S. and Swanson, D. L.** (2014). Seasonal Phenotypic Flexibility of Body Mass, Organ Masses, and Tissue Oxidative Capacity and Their Relationship to Resting Metabolic Rate in Chinese Bulbuls. *Physiol. Biochem. Zool.* **87**, 432–444.

Tables

Table 1. Response and explanatory variables in models used. Explanation of each variable as well as units included

	Variable	Units	Code	Explanation
Response	Thyroid hormone concentration	ng/mL	T3	Concentration of total Triiodothyronine hormone in blood samples from adult shags. Hormonal proxy for RMR
	Daily Energy Expenditure	L/day	DEE	Total energy expenditure per 24 hour estimated using accelerometry and converted to rate of oxygen consumption via published calibrations for this species
Explanatory	Parasite load	worms	Para	Number of nematode worms in individual quantified with endoscope
	Brood Size	chicks	B.size	Number of chicks in nest may affect the energetic constraint adults are under in terms of provisioning
	Brood Age	days	B.age	Age of the oldest chick in the brood in days when the logger was deployed, may affect the energetic constraint adults are under as larger chicks need more food.
	Body mass	g	mass	Body mass of the adult sampled; body mass is found to relate to Thyroid hormone levels in some studies
	Age	years	Age	Age of the adult in years, this may have effects on thyroid hormone levels The study incorporated a full range of ages (Males 3-20, Females 4-14)
	Year		Year	Year in which the data was collected may have effects on thyroid hormone levels. Year is included as a factor to account for this variation

Table 2. The top ten best supported modes selected by AIC of a linear mixed effect model to explain the effect of Parasite load and other variables on RMR (T3 concentration) for females and males. Asterisks between terms indicate an interaction between terms and the top best supported model is indicated in bold.

	(Int)	Age	mass	Para	B. Size	B. Age	Year	Para * B.size	Para * B. Age	df	log link	AIC	Δ AIC	Weight
Female	-0.11			0.53		-0.29	+		-0.35	8	-40.23	96.50	0.00	0.23
	-0.11	-0.17		0.52		-0.29	+		-0.34	9	-39.66	97.30	0.87	0.15
	-0.26		-0.23	0.56		-0.37	+		-0.46	9	-39.74	97.50	1.02	0.14
	-0.27	-0.18	-0.24	0.54		-0.37	+		-0.46	10	-39.11	98.20	1.77	0.09
	-0.10			0.53	-0.05	-0.29	+		-0.37	9	-40.16	98.30	1.87	0.09
	0.24			0.64		-0.17	+			7	-42.44	98.90	2.43	0.07
	0.39			0.68			+			6	-43.45	98.90	2.45	0.07
	-0.11	-0.17		0.51	-0.05	-0.28	+		-0.36	10	-39.59	99.20	2.72	0.06
	-0.26		-0.24	0.55	-0.06	-0.37	+		-0.49	10	-39.66	99.30	2.86	0.06
	-0.09			0.59	-0.08	-0.33	+	-0.20	-0.37	10	-39.68	99.40	2.91	0.05
Male	-0.12	-0.40					+			6	-56.57	125.10	0.00	0.18
	-0.32	-0.37	0.33	-0.17			+			8	-54.66	125.30	0.17	0.16
	-0.31	-0.39	0.26				+			7	-55.75	125.50	0.37	0.15
	-0.10	-0.38		-0.13			+			7	-55.92	125.80	0.71	0.13
	-0.39	-0.39	0.38	-0.20		0.11	+			9	-54.43	126.90	1.72	0.08
	-0.11	-0.39				-0.04	+			7	-56.54	127.10	1.94	0.07
	-0.13	-0.40			-0.02		+			7	-56.56	127.10	1.98	0.07
	-0.33	-0.37	0.34	-0.16	-0.02		+			9	-54.65	127.30	2.15	0.06
	-0.35	-0.39	0.28		-0.06		+			8	-55.66	127.30	2.18	0.06
	-0.31	-0.39	0.27			0.01	+			8	-55.75	127.50	2.37	0.06

Table 3. The top ten best supported modes selected by AIC from a mixed effect linear model to explain the relationship between daily energy expenditure estimated from accelerometry and RMR (assessed by a proxy; plasma concentration of T3 hormone) for female and male European shags. The best supported models are indicated in bold.

	(Int)	Age	mass	T3	B. Size	B. Age	Year	df	log link	AIC	Δ AIC	Weight
Female	0.58							3	-38.18	82.40	0.00	0.20
	0.41						+	4	-37.29	82.60	0.22	0.18
	0.10		-0.52					4	-37.39	82.80	0.41	0.16
	-0.12		-0.55				+	5	-36.40	82.80	0.43	0.16
	0.57					0.20		4	-38.14	84.30	1.91	0.08
	0.42					0.17	+	5	-37.61	85.20	2.85	0.05
	0.19		-0.41			0.17		5	-37.68	85.40	2.99	0.05
	0.57	-0.07						4	-38.71	85.40	3.06	0.04
	0.39	-0.08					+	5	-37.79	85.60	3.21	0.04
	0.59				-0.14			4	-38.80	85.60	3.23	0.04
Male	-0.18		-0.50					4	-49.03	106.10	0.00	0.31
	-0.54							3	-50.27	106.50	0.49	0.24
	-0.15		-0.59		0.39		+	7	-47.20	108.40	2.34	0.10
	-0.15		-0.51		0.17			5	-49.37	108.70	2.68	0.08
	-0.53				0.15			4	-50.78	109.60	3.50	0.05
	-0.20		-0.47			0.09		5	-49.90	109.80	3.73	0.05
	-0.24		-0.48				+	6	-48.97	109.90	3.89	0.04
	-0.54					0.12		4	-50.98	110.00	3.90	0.04
	-0.17	-0.02	-0.50					5	-49.99	110.00	3.92	0.04
	-0.18		-0.48	-0.05				5	-50.05	110.10	4.03	0.04

Figures




Fig. 1. Graphical representation of energy budgets and possible energy management strategies. **a.** Daily energy expenditure (DEE) is the sum of resting metabolic rate (RMR) and activity metabolism. **b.** Three theorised energy management strategies in graphical form adapted from Mathot & Dingemans, (2015) and Careau & Garland, (2012) showing the relationship between RMR and DEE and activity metabolism under the three scenarios, the dashed 1-to-1 line is presented to aid interpretation of the slope of the lines. It represents a theoretical situation under which animals are completely inactive and thus DEE is equal to RMR .




Fig 2. Relationship between parasite load and plasma total 3, 3'-Triiodo-L-Thyronine (T3) concentration (a proxy for resting metabolic rate) in female European shags. Data points represent individual T3 concentrations, the predicted lines from the best supported model (solid lines) and their 95% confidence intervals (dashed). **A** shows the relationship between parasite load and T3 concentration when year and brood age are accounted for, **B** shows the interaction between parasite load and brood age in relation to T3 concentration, the explanatory terms in the best supported model. Solid lines represent predicted lines from the best supported model under different brood age scenarios -1 Standard deviation of the mean brood age (light blue) Mean brood age (mid blue), and +1 Standard deviation of the mean (dark blue).




Fig 3. Significant relationship between adult age and plasma total 3, 3'-Triiodo-L-Thyronine (T3) concentration (a proxy for resting metabolic rate) in male European shags. Data points represent individual T3 concentrations, the predicted lines from the best supported model (solid lines) and their 95% confidence intervals (dashed).




Fig 4. Relationship between plasma total 3, 3'-Triiodo-L-Thyronine (T3) concentration (a proxy for resting metabolic rate) and estimated DEE in European shags. Dots represent individual T3 concentrations, the predicted lines from the best supported model (solid lines) and their 95% confidence intervals (dashed) are presented for males (blue) and females (Orange).