

CONTROL OF INSECT ECDYSIS BY A POSITIVE-FEEDBACK ENDOCRINE SYSTEM: ROLES OF ECLOSION HORMONE AND ECDYSIS TRIGGERING HORMONE

JOHN EWER, STEPHEN C. GAMMIE AND JAMES W. TRUMAN*

Department of Zoology, University of Washington, Box 351800, Seattle, WA 98195-1800, USA

Accepted 5 December 1996

Summary

A successful ecdysis in insects requires the precise coordination of behaviour with the developmental changes that occur late in a moult. This coordination involves two sets of endocrine cells: the peripherally located Inka cells, which release ecdysis triggering hormone (ETH), and the centrally located neurosecretory neurones, the VM neurones, which release eclosion hormone (EH). These two sets of endocrine cells mutually excite one another: EH acts on the Inka cells to cause the release of ETH. ETH, in turn, acts on the VM neurones to cause the release of EH. This positive-feedback relationship allows the Inka cells and the VM neurones to be the peripheral and central halves, respectively, of a decision-making circuit. Once conditions

for both halves have been satisfied, their reciprocal excitation results in a massive EH/ETH surge in the blood as well as a release of EH within the central nervous system. This phasic signal then causes the tonic activation of a distributed network of peptidergic neurones that contain crustacean cardioactive peptide. The relationship of the latter cells to the subsequent maintenance of the ecdysis motor programme is discussed.

Key words: eclosion hormone, ecdysis triggering hormone, crustacean cardioactive peptide, neuromodulation, feedback, *Manduca sexta*.

Introduction

Positive-feedback systems are rare in biology but are utilized in situations that require a phasic, unambiguous signal. In endocrinology, the classic example of such a system is the control of ovulation in mammals in which the hypothalamus and ovary interact positively to generate the luteinizing hormone surge that coordinates ovulatory events. In the case of insect physiology, the process of ecdysis likewise requires an unambiguous signal. Its success is dependent on the precise coordination of behaviour with ongoing events of the moult and, once initiated, these events cannot be reversed.

The insect goes through a sequence of behavioural phases as the time for ecdysis approaches (Carlson, 1977; Hughes, 1980). Late in a moult, as the synthesis of the new cuticle and the digestion of the old one near completion, an insect typically becomes restless and seeks a suitable site for ecdysis. As this preparatory phase progresses, the animal anchors itself to the substratum and begins movements that loosen the connections between the old and new cuticles. The preparatory phase is variable in duration, but it eventually culminates in the ecdysis attempt. The ecdysis phase starts with abdominal peristaltic waves that begin to move the old cuticle posteriorly. These are coordinated with other behavioural subroutines to extract the appendages from the cuticle as it is shed and with changes in

blood distribution and with air-swallowing that expand the anterior end of the animal and aid in rupturing the old cuticle (Reynolds, 1980). The initiation of ecdysis is typically associated with irreversible changes in the new cuticle so that the failure to accomplish ecdysis in a timely manner results in crippling deformities or an animal permanently trapped within its old skin.

Most of the studies on the hormonal control of ecdysis have been carried out on various stages of the tobacco hornworm *Manduca sexta*. Two endocrine systems have been implicated in initiating ecdysis (Fig. 1). One is in the brain and includes two pairs of neurones (the VM neurones) whose axons project through the length of the central nervous system (CNS) as well as to peripheral neurohaemal sites (Truman and Copenhaver, 1989). These cells are the source of the neuropeptide eclosion hormone (EH). The second endocrine system is a peripheral one consisting of a set of three-celled glands, the epitracheal glands, one of which is located on the trachea near each of the 18 spiracles (Fig. 1D,E; Zitnan *et al.* 1996). The Inka cell component of each gland produces a peptide hormone, ecdysis triggering hormone (ETH; Zitnan *et al.* 1996). Injection of either ETH (Zitnan *et al.* 1996) or EH (Copenhaver and Truman, 1982; Truman *et al.* 1983)

*e-mail: trumanj@zoology.washington.edu.




Fig. 1. Drawing of a moulting second-instar larva of *Manduca sexta* showing the cells that are involved in ecdysis control. Photographs are confocal images showing: (A,B) eclosion hormone immunostaining of (A) the cell bodies of the two pairs of VM neurones near the midline of the brain and (B) the VM cell axons in the proctodeal nerves extending on either side of the hindgut (hg); (C) CCAP immunoreactivity in paired cells 27 and 704 in ganglion A2; (D,E) a pharate pupal epitrichal gland attached to tracheal epidermis (t) showing (D) the arrangement of the three cells in the gland as revealed by their autofluorescence and (E) the ETH-like immunostaining present in the Inka cell (ic).

induces ecdysis, but ETH triggers the behaviour more rapidly than does EH and behavioural sensitivity to ETH occurs earlier in the moult. Zitnan *et al.* (1996) proposed that EH caused the release of ETH which, in turn, caused ecdysis behaviour. Although the evidence for this endocrine pathway is compelling, this relationship by itself fails to account for some known aspects of ecdysis control (Hesterlee and Morton, 1996). For example, debrained larvae injected with EH show good pre-ecdysis behaviour but not ecdysis (A. Novicki and J. C. Weeks, unpublished results, cited in Hesterlee and Morton, 1996), and back-firing of the VM neurones that contain EH can lead to the rapid onset of ecdysis behaviour (Hewes and Truman, 1991).

Besides the VM neurones and Inka cells, the neurones of the cell 27/704 group are another set of peptidergic cells that are associated with ecdysis (Fig. 1C). These segmentally reiterated neurones include both interneurones and neurosecretory cells. In most insects, the neurones of this group contain crustacean cardioactive peptide (CCAP) (Davis *et al.* 1993; Dirksen, 1994). In *Manduca sexta* (Ewer *et al.* 1994) and a wide variety of other insects (Ewer and Truman, 1996), these neurones show a striking accumulation of the second messenger cyclic 3',5'-guanosine monophosphate (cGMP) just prior to the onset of ecdysis. The rise in intracellular cGMP level enhances the

excitability of these cells (Gammie and Truman, 1977), presumably facilitating the release of their neuropeptides. Since some of these neurones innervate the foregut and the alary muscles of the heart (Braüning, 1990; Dirksen *et al.* 1991; Ewer *et al.* 1994), they are the likely candidates for causing the air-swallowing and circulatory changes that accompany ecdysis (Reynolds, 1980).

This paper reports that the VM neurones, the Inka cells and the CCAP neurones constitute a neuroendocrine circuit that initiates and maintains ecdysis behaviour. The VM neurones and the Inka cells are two halves of a positive-feedback circuit that ensures that peripheral and central conditions are appropriate for ecdysis. A key outcome of their interaction is the activation of the neurones of the cell 27/704 group whose peptides then shift the nervous system into the ecdysis phase itself.

Materials and methods

Experimental animals

Larvae of *Manduca sexta* (Johansson) were raised on an artificial diet at 26 °C under a 17h:7h L:D photoperiod. Some experiments were performed on larvae during the moult from the first to the second instar. Fifth-instar larvae were also used. Larvae were selected at the time of head-capsule slippage (HCS, when the old head capsule moves down over the forming mandibles of the next larval stage) and were maintained at room temperature (18–20 °C) until ecdysis approximately 16h later. The larvae showed obvious tanning of the new mandibles (brown mandible stage) by 10h after HCS. The time course of events in moulting fifth-stage larvae is described in Truman (1972) and Curtis *et al.* (1984).

Immunocytochemistry

For immunocytochemistry of moulting second-instar larvae, staged animals were cut along the dorsal midline, pinned out under 4% buffered formaldehyde, and fixed for 1–2h. Tissues were then pre-incubated in 2.5% normal goat serum (NGS) and 2.5% normal donkey serum (NDS) for 1h in phosphate-buffered saline with 0.3% Triton X-100 (PBS-TX), and then overnight at 5 °C in a cocktail of primary antibodies [a monoclonal antibody against the molluscan small cardioactive peptide (SCP; Masinovsky *et al.* 1988), which recognizes the C terminus of ETH (Zitnan *et al.* 1996), was used at a dilution of 1:500, a rabbit anti-cGMP antiserum (De Vente *et al.* 1987) was used at a dilution of 1:4000, and a rabbit anti-EH antiserum (Truman and Copenhaver, 1989) was used at a dilution of 1:250] in PBS-TX with 1% NGS and NDS. After repeated washes, the tissues were incubated overnight at 5 °C in 1:1000 dilutions of a FITC-conjugated donkey anti-rabbit IgG and a Texas-Red-conjugated goat anti-mouse IgG (both Jackson Laboratories) in PBS-TX with 1% NGS and NDS. After repeated washes, the tissues were dehydrated, mounted in DPX (Fluka) and examined using a BioRad 600 confocal microscope. EH and cGMP immunoreactivity were distinguishable from one another because they occur in distinct

cell populations. In a limited number of preparations, the spatial distinctness of the EH and cGMP signals was confirmed using an anti-cGMP antiserum that had been raised in sheep rather than rabbit.

For analysis of fifth-stage larvae, individual tissues were dissected, fixed overnight, rinsed in PBS-TX, treated in 0.5 mg ml^{-1} collagenase (Sigma Chemical Co.) for 30 min at room temperature to aid in antibody penetration, and processed for immunocytochemistry as described above.

Immunoassays for ETH

The amount of ETH in the bathing medium was determined by an antibody-capture immunoassay (Harlow and Lane, 1988). $5 \mu\text{l}$ samples of the medium were bound to the bottom of wells of polystyrene 96-well plates (Corning, no. 25802). Wells were incubated overnight with 1:1000 anti-SCP monoclonal antibody, followed by 1:1000 peroxidase-conjugated goat anti-mouse IgG. The amount of bound antibody-enzyme complex was determined using a Peroxidase Substrate kit (Bio-Rad), and the intensity of product was measured using a Microplate Autoreader (BioTek Instruments). Each sample was run in duplicate. Absorbance from each well was compared with a standard dilution series of Inka cell extract.

Hormonal materials

The eclosion hormone used in these studies were from two sources. Some studies utilized recombinant EH produced from baculovirus that carried a copy of the *Manduca* EH gene (Eldridge *et al.* 1991). We also used EH that had been purified from corpora cardiaca-corpora allata complexes from pharate adult *Manduca sexta* (Terzi *et al.* 1988).

For preparation of epitracheal gland extracts, freshly dissected glands from prepupae together with some adhering tracheal tissue were homogenized in acidified methanol (90:9:1 methanol:water:acetic acid). After centrifugation for 1 min in a PicoFuge (Stratagene), the supernatant was blown to dryness under a stream of nitrogen and resuspended in saline at a concentration of one epitracheal gland equivalent per $10 \mu\text{l}$. Synthetic ETH was prepared by the Howard Hughes Macromolecular Synthesis Facility at the University of Washington.

Pharate second-stage larvae were injected using glass microneedles that were tip-filled with the appropriate solution. The needles were then fitted into an oil-filled injection system that delivered $1 \mu\text{l}$ into the larva. Pharate fifth-instar larvae were injected using 50 or $100 \mu\text{l}$ syringes.

Electrophysiological techniques

Nervous systems were dissected from pharate fifth-instar larvae that were approximately 5 h before ecdysis. The CNS was briefly exposed to 4% collagenase/dispase (Boehringer Mannheim) to aid in penetration of the sheath. The bathing solution contained (in mmol l^{-1}): 140 NaCl, 5 KCl, 4 CaCl_2 , 28 D-glucose, and 5 Hepes (pH 7.4). Glass microelectrodes were filled with 2 mol l^{-1} potassium acetate and 10 mmol l^{-1} KCl (pH 7.4). VM cell bodies could be identified visually and

are the only neurones in that region of the brain with overshooting action potentials in their somata (Hewes and Truman, 1994).

The signals were amplified using a Getting microelectrode amplifier (Getting Instruments, Iowa City, IA, USA), stored on a VHS recorder (Vetter Co., Rebersburg, PA, USA) and analysed using SuperScope (GW Instruments, Somerville, MA, USA). Any compensation errors made during bridge-balance recordings were corrected during off-line analysis by subtracting the near instantaneous voltage record attributable to the potential drop across the electrode (Hewes and Truman, 1994). Threshold was determined by injecting positive current pulses (600–1000 ms) into the somata and calculating the minimum voltage deflection from the resting potential needed to fire an action potential.

Results

The timing of behavioural and endocrine events at ecdysis

Fig. 2 examines the timing of the endocrine events that occur during ecdysis of second-instar larvae of *Manduca sexta*. In *M. sexta* larvae, the preparatory phase consists of the pre-ecdysis behaviour, a set of rhythmic thoracic and abdominal contractions that loosen connections between the old and new cuticles (Copenhaver and Truman, 1982; Miles and Weeks, 1991; Novicki and Weeks, 1993). Prior to the onset of pre-ecdysis behaviour, the proctodeal nerves, the peripheral release site for EH in larvae (Truman and Copenhaver, 1989), were well stocked with EH, showing an average of approximately 50 EH-immunoreactive varicosities per $150 \mu\text{m}$ of nerve (Fig. 2B). EH staining in the proctodeal nerve remained unchanged through the first 25 min of the pre-ecdysis behaviour, but there was then a marked reduction in staining between 25 and 35 min. In larvae that were in this transition period, we occasionally found proctodeal nerves that showed diffuse EH staining in the extracellular stroma surrounding the terminal varicosities (Fig. 2A, 32 min). We interpret this extracellular staining as being due to newly released EH that was fixed before it could diffuse through the stroma and into the circulation. The loss of EH immunoreactivity from peripheral release sites was also accompanied by the disappearance of EH immunoreactivity from the VM cell axons within the CNS (Fig. 2A). Thus, for ecdysis of the second instar, as for pupal ecdysis (Hewes and Truman, 1991), EH is released within the CNS coincident with its liberation into the circulation.

The neurones of the cell 27/704 group showed an abrupt rise in intracellular cGMP levels during the latter phases of the pre-ecdysis behaviour (Fig. 2). These cGMP-immunoreactive neurones were found only in those larvae that had at least begun the process of EH release from the VM cells. These neurones then maintained elevated cGMP levels throughout ecdysis (see also Ewer *et al.* 1994).

M. sexta larvae have 18 epitracheal glands, one associated with each spiracle (Zitnan *et al.* 1996). The Inka cell component of these glands showed intense ETH staining up to




Fig. 2. Timing of the endocrine events during the period leading up to ecdysis of second-instar larvae of *Manduca sexta*. (A) Confocal images from larvae that were early (15 min) or late (32 min) in pre-ecdysis behaviour and at ecdysis (50 min): a, EH immunoreactivity in a segment of proctodeal nerve; b, EH immunoreactivity in the axons of the VM neurones in the terminal ganglion (arrowhead: EH immunoreactivity still present at 15 and 32 min); c, cGMP immunoreactivity in the paired cells 27 and 704 in ganglion T1; and double labels showing: d, ETH-like immunoreactivity and e, cGMP immunoreactivity in Inka cells. Scale bars, 25 μm . (B) Top, changes in the number of EH-staining (EH-IR) varicosities observed in a standard length (150 μm) of proctodeal nerve late in the moult: each point is the average of counts from the right and left nerves at the level of the ileum for a single individual; filled circles are larvae which also showed cGMP immunoreactivity in the cell 27/704 group. Bottom, the number of Inka cells per larva that stained positively for ETH (ETH-IR); filled circles are larvae whose Inka cells also exhibited cGMP immunoreactivity. BM, brown mandible stage, approximately 5–6 h before ecdysis; Air, air appearing in the old head capsule; Ecd, ecdysis.

and through the start of pre-ecdysis behaviour (Fig. 2B). ETH immunoreactivity began to disappear abruptly from these cells in larvae that were undergoing EH release at 25–30 min into the pre-ecdysis behaviour. As they were losing their ETH immunoreactivity, these cells also showed elevated intracellular cGMP levels (Fig. 2A, 32 min). Unlike that in the neurones of the cell 27/704 group, however, the increase in cGMP level in the Inka cells was transient and their cGMP levels had declined to below the level of detectability by ecdysis (Fig. 2).

The relationship between EH and ETH release

Experiments involving the incubation of isolated nervous systems with epitracheal glands and EH suggested that EH acts on these glands to cause ETH release (Zitnan *et al.* 1996). We tested this hypothesis using isolated epitracheal glands attached to a small segment of trachea. Individual epitracheal glands, taken from prepupae at the 'anterior shrink' stage (approximately 3–4 h before ecdysis), were incubated with agitation in 50 μl drops of *Manduca* saline (Riddiford *et al.* 1979) containing 0.02% BSA and 1 nmol l⁻¹ purified EH (Terzi *et al.* 1988). After the incubations, the bathing medium was collected and frozen until assayed. As seen in Fig. 3A, the presence of EH resulted in a time-dependent release of ETH

immunoreactivity into the medium. By contrast, control incubations with BSA alone resulted in a low, basal level of ETH secretion.

The time course of the response of the Inka cells to EH exposure *in vitro* was followed using immunocytochemistry. Inka cells were exposed to EH *in vitro* and then, at various times thereafter, fixed and double-stained for ETH and for cGMP immunoreactivity. As seen in Fig. 3C, detectable cGMP immunoreactivity became evident in the Inka cells by approximately 5 min after EH addition and was prominent by 10 min. The ETH staining in the resting cell was evident in small granules concentrated in the cortical region of the cell, but by 10–20 min following EH addition these granules began to increase in size and to disappear from the cell cortex (Fig. 3C). We do not know whether the increase in granule size is due to osmotic swelling or to fusion of granules with each other as well as with the cell membrane. ETH staining was severely depleted in these cells by 40–60 min after EH addition. These results clearly support the proposal by Zitnan *et al.* (1996) that EH acts on the epitracheal cells to cause the release of ETH. A direct demonstration of this action of EH on the Inka cells has also been made (T. Kingen, J. Hermesman, D. Zitnan and M. Adams, in preparation).

Zitnan *et al.* (1996) suggested that injected EH causes its




Fig. 3. Secretory response of isolated epitracheal glands to eclosion hormone (EH) *in vitro*. (A) Mean + s.e.m. ETH-like immunoreactivity (ETH-IR) released into the medium by single epitracheal glands cultured for 30 and 60 min with 0.02 % bovine serum albumin (BSA) or with BSA with 1 nmol l^{-1} EH. Numbers of cultures are indicated. (B) Dual-channel, optical sections through the Inka cell component of epitracheal glands. Examples show cGMP- and ETH-like immunostaining for glands that were fixed before (0 min) or 40 or 60 min after exposure to EH. EH causes the appearance of cGMP immunoreactivity and the loss of ETH immunoreactivity. The boxed area in the 40 min cGMP panel shows the approximate amount of the cell represented in C. (C) cGMP and ETH immunoreactivity changes in the cortical region of Inka cells at various times after exposure to EH. Five glands were exposed in each group; the example shown is the most advanced gland in each group. C, control glands incubated without EH for 60 min. Scale bars $10 \mu\text{m}$ for B and C.

behavioural effects by stimulating ETH release and that the latter, in turn, acts on the CNS to cause ecdysis. Injections of EH are effective in triggering ecdysis behaviour only during the final hours prior to ecdysis (Truman *et al.* 1983; Copenhaver and Truman, 1982). For example, in pharate second-instar larvae, the onset of EH sensitivity occurred approximately 2.5 h before normal ecdysis (Fig. 4A). As seen in Fig. 4D,E, this time of behavioural sensitivity coincided with the time when EH injection was first able to induce the loss of ETH immunoreactivity from the Inka cells. The ability of animals to respond to injected EH requires a decline in the circulating titres of the ecdysteroids that drive the moulting process (Truman *et al.* 1983). We had previously interpreted this time as reflecting the time when the CNS became responsive to circulating EH. It now appears that this timing reflects the competence of the Inka cells rather than that of the CNS. Whether these cells become competent through a direct action of ecdysteroids on them or indirectly through the steroid-induced changes in the epidermis on which they reside is not yet known.

Effects of ETH on the activity of VM neurones

Although EH causes ETH release, the situation is reciprocal in that we find that ETH acts on the CNS to excite the VM neurones, resulting in EH release. Outside their normal time of EH release, the VM cells are relatively inexcitable and sustained depolarization results in single action potentials rather than trains of spikes (Hewes and Truman, 1994). For example, intracellular recordings from VM cells in isolated, pharate fifth-instar nervous systems showed neurones with a spike threshold of $27 \pm 3 \text{ mV}$ (mean \pm s.e.m.; $N=6$) over rest. Addition of epitracheal gland extracts resulted in the onset of tonic firing of these cells within 4–20 min (mean time 10 min; $N=6$) (Fig. 5A). Once activated by these extracts, the VM neurones then maintained tonic firing at 1–2 Hz for at least 20 min.

The epitracheal gland extracts also caused a dramatic broadening of the soma action potential of the VM neurones. Prior to addition of the extracts, the duration of the action potential was $34 \pm 3 \text{ ms}$, but it then increased almost threefold to $84 \pm 10 \text{ ms}$ by 10–15 min after extract addition ($N=6$;




Fig. 5B). Two observations show that this dramatic spike-broadening was not due simply to the repetitive firing of the VM cell. First, driving these neurones at 1–2 Hz in the absence of gland extracts did not result in significant spike-broadening (S. C. Gammie and J. W. Truman, unpublished observations). Second, neurones depolarized after extract addition but prior to the start of the spontaneous spike train already show a markedly broadened soma spike (Fig. 5B, spike b).

Fig. 4. Development of sensitivity of endocrine centres to injections of EH at various times during the moult to the second larval stage. (A) Percentage of larvae showing ecdysis behaviour within 60 min after injection. (B–E) Immunological assessment of the amount of EH and ETH still stored in the proctodeal nerve and Inka cells, respectively, at 60 min after the injection. (B) Number of EH-positive (EH-IR) varicosities in a 150 μm length of proctodeal nerve. (C) Confocal images showing examples of the EH immunoreactivity still present in the proctodeal nerve following EH injection at the indicated times before ecdysis (in h). (D) the number of Inka cells per larva that stained positively for ETH (ETH-IR); filled circles are larvae whose Inka cells also exhibited cGMP immunoreactivity at the time of dissection. (E) Confocal images showing the typical level of ETH-like immunoreactivity present in the Inka cells following EH injection at the indicated times. The two confocal examples given for –2.5 h show the proctodeal nerve and corresponding Inka cell from a representative of the group that had released EH and one from the group that had not. Data sets in B and D are represented as in Fig. 2B, except the cGMP status of the cell 27/704 group is not presented. The –6 h time point coincided with the start of the brown mandible stage.

The epitracheal glands apparently contain a number of biologically active peptides (Zitnan *et al.* 1996). Preliminary experiments using synthetic ETH ($1\text{--}5\ \mu\text{mol l}^{-1}$) show that this peptide is sufficient to induce both spike-broadening and tonic firing of the VM neurones ($N=3$).

Effects of Inka cell extracts on EH release *in vivo*

Inka cell peptides promote the release of EH *in vivo* as well as *in vitro*. Their effects on EH release were first tested in pharate fifth-instar larvae that were selected 2 h after the head capsule became filled with air (approximately 3 h before ecdysis). Injection of two Inka-cell-equivalents of extract was followed by an abrupt loss of EH immunoreactivity from the proctodeal nerve within 20 min (Fig. 6F) followed by ecdysis at 35 ± 1 min (mean \pm S.E.M.; $N=12$). Injection of similar larvae with about 1.5 ng of recombinant EH also caused the release of endogenous stores of EH but, interestingly, the latency to EH release was longer (30–35 min; Fig. 6E) and ecdysis occurred later (49 ± 2 min; mean \pm S.E.M.; $N=17$) than that seen after Inka cell extract treatment. The longer latency to EH release in the latter case presumably reflects the time required for the injected EH to cause ETH release which, in turn, causes release of endogenous EH stores.

In the case of both EH and Inka cell extract injections, the release of endogenous EH stores coincided with the appearance of cGMP in the cell 27/704 group (Fig. 6A–D). This was then followed by ecdysis 15–20 min later. Thus, irrespective of which peptide was injected, the onset of ecdysis had a constant temporal relationship with the release of endogenous EH and the accompanying increase in cGMP levels in the cell 27/704 group.

The window of development during which *M. sexta* will show ecdysis behaviour in response to injections of peptide is strikingly different for EH *versus* ETH. There is only a very narrow window, late in the moult, during which the animal will




Fig. 5. Intracellular recording from a VM neurone showing its response to an epitracheal gland extract (EGX) containing approximately 5 pmol of ETH. (A) Induction of spontaneous firing after addition of the extract. (B) Records showing the change in duration of the VM cell action potential (a) before, and (b) 6 and (c) 15 min after addition of the gland extract. In a and b, the spike was elicited by depolarization of the neurone.

respond to injections of EH (e.g. Reynolds *et al.* 1979; Copenhagen and Truman, 1982), whereas the window of sensitivity to ETH begins many hours earlier (Zitnan *et al.* 1996). For example, moulting fifth-stage larvae begin to show premature ecdysis behaviour in response to Inka cell extract injections starting shortly after head-capsule slippage (HCS, approximately 28–34 h before ecdysis). By contrast, these larvae respond to EH injections only much later, at approximately 4–5 h before normal ecdysis (Copenhagen and Truman, 1982).

As seen in Fig. 7A, moulting second-instar larvae showed premature ecdysis behaviour in response to injections of Inka cell extracts starting approximately 13 h before normal ecdysis (approximately 3 h after HCS), while EH injections were not effective until approximately 2.5 h before ecdysis (Fig. 4A). These larvae were then all examined 60–90 min after peptide injection to determine their state of EH release. Importantly, all of the animals that showed premature ecdysis in response to either Inka cell extract or EH injection also showed the premature release of their store of EH (Figs 4B,C, 7B,C). These larvae also consistently showed cGMP in the neurones of the cell 27/704 group (data not shown).

Although injections of epitracheal gland extracts were capable of causing EH release starting approximately 13 h before expected ecdysis, we saw no evidence of release of endogenous stores of ETH from the Inka cells until approximately 2.5 h before ecdysis (Fig. 7D,E). This latter time corresponds to when Inka cells first start to show ETH release after injections of EH (Fig. 4D,E). This temporal relationship suggests that the release of ETH after injection of gland extracts is probably a secondary response that follows the release of EH into the circulation. However, it only occurs at late times when the Inka cells finally become responsive to EH.

Relationship of EH and ETH to the activation of the neurones in the cell 27/704 group

Neurones of the cell 27/704 group invariably showed an increase in cGMP levels in association with EH release, whether during a normal, spontaneous ecdysis (Fig. 2) or during an early ecdysis provoked by injection of Inka cell

peptides or EH (Fig. 6E,F). Direct evidence that the VM cells were responsible for this cGMP increase was obtained by repetitively firing the VM neurones with intracellular electrodes (1–2 Hz) or by ‘back-firing’ the VM cell axons through stimulation of the proctodeal nerves as in Hewes and Truman (1991). Both treatments resulted in the appearance of cGMP in the neurones of the cell 27/704 group (in 4 of 10 cells for intracellular stimulation of single VM cells and in 12 of 15 cells for back-firing of the proctodeal nerves). The intracellular stimulation was carried out on isolated nervous systems that lacked tracheae and, hence, epitracheal glands. Consequently, these glands and the ETH that they release are not required for the appearance of cGMP in the cell 27/704 group.

The role of the VM neurones in inducing the cGMP increase in this cell group is also consistent with *in vivo* experiments in which the ventral nerve cord was transected prior to larvae being challenged with either EH ($N=14$) or Inka cell peptides ($N=9$). After peptide challenge, the CNS anterior to the transection subsequently showed EH depletion from the VM axons, the appearance of cGMP in the cell 27/704 homologues and the onset of ecdysis movements. Posterior to the transection, by contrast, there was no EH release, no appearance of a robust cGMP response and no performance of ecdysis behaviour. Thus, rather than acting directly, the signal from peptides circulating in the blood needs to be relayed through the descending signal provided by the activity of the VM neurones.

Discussion

The endocrine trigger for ecdysis behaviour

During the past 25 years it has been thought that a single peptide, eclosion hormone, was responsible for triggering ecdysis behaviour (Truman, 1992). The recent report of a second ecdysis triggering peptide, ETH (Zitnan *et al.* 1996), showed that the regulation of ecdysis is more complex than originally thought and has sparked a re-evaluation of the role of EH (e.g. Hesterlee and Morton, 1996). In assessing the relationship between EH and ETH, a number of previously held assumptions about EH require re-examination. One assumption has been that EH release in *M. sexta* occurs prior




Fig. 6. Time course of EH depletion from the proctodeal nerves and the appearance of cGMP immunoreactivity in the cell 27/704 group after injection of pharate fifth-instar larvae with EH or Inka cell peptides. (A–D) Confocal images showing cGMP immunoreactivity in ganglion A4 (left panels) and EH immunoreactivity in the proctodeal nerve (right panels) from the same animal for larvae at 0 (A) and 30 min (B–D) after injection with EH. (B,C) A partial cGMP response in cells 27/704 (arrowheads), accompanied by a reduction (B) or loss (C) of EH immunoreactivity in the VM axon (arrow in B). (E,F) Percentage of animals showing EH immunoreactivity in the proctodeal nerves (triangles) and cGMP immunoreactivity in the cell 27/704 group (squares) at various times after injection of EH (E) or Inka cell peptides (F). In both cases, the appearance of cGMP immunoreactivity coincided with the loss of EH immunoreactivity. Numbers indicate the number of preparations examined at each time. Note that the events triggered by Inka cell peptides (F) occur approximately 15 min earlier and with a steeper time course than those triggered by EH (E). Scale bars, 100 μ m.

to the start of pre-ecdysis behaviour. Since the injection of EH is always followed by the performance of the pre-ecdysis behaviour (Copenhaver and Truman, 1982; Miles and Weeks, 1991), the normal sequence of events was thought to be: EH release, pre-ecdysis behaviour, then ecdysis behaviour. The data in Fig. 2, as well as results from Novicki and Weeks (1996), are incompatible with this ordering of events, although we cannot exclude the possibility that a small release of EH precedes pre-ecdysis. Nevertheless, the major surge of EH

clearly occurs well into pre-ecdysis behaviour, approximately 15–20 min before the start of ecdysis. This timing is consistent with old data from pupal ecdysis in *M. sexta* showing EH activity appearing in the haemolymph approximately 20–25 min prior to the start of ecdysis (Truman *et al.* 1980). The shortness of this interval was surprising at the time since the normal latency between EH injection and pupal ecdysis was 60–65 min.

The isolated abdominal CNS of *Hyalophora cecropia* was




Fig. 7. Development of sensitivity of endocrine centres to injections with extracts of Inka cells (0.1 prepupal cell equivalent per larva) at various times during the moult to the second larval stage. (A) Percentage of larvae showing ecdysis behaviour within 60 min of injection with the extract. (B–E) Immunological assessment of the amounts of EH and ETH still present in the proctodeal nerves and Inka cells, respectively, at 60–90 min after the injection. (B) Number of EH-immunoreactive (EH-IR) varicosities in a 150 μ m length of proctodeal nerve. (C) Confocal images showing examples of the EH immunoreactivity present in the proctodeal nerve following extract injection at the indicated times before ecdysis (in h). (D) The number of Inka cells per larva that stained positively for ETH (ETH-IR). (E) Confocal images showing the typical level of ETH-like immunoreactivity present in the Inka cells following extract injection at the indicated times. The two confocal examples given for –2.5 h animals in E are from larvae that did and did not release ETH. Data sets in B and D are represented as in Fig. 2B, except that the cGMP status of the cell 27/704 group is not presented.

used to demonstrate that EH acts on the CNS to induce the ecdysis motor programme (Truman, 1978). The success of these experiments, however, required that the tracheal supply to the CNS be included, presumably to ensure an adequate delivery of oxygen to the tissue (Truman, 1978). The results from Zitnan *et al.* (1996), however, suggest a new

interpretation for the need for the tracheae since these tracheal trunks are the sites on which the epitracheal glands reside. Since ETH can induce an ecdysis motor programme from a *M. sexta* CNS that lacks tracheae, these authors suggested that the EH added *in vitro* acts on the epitracheal glands to cause ETH release which, in turn, acts on the CNS to cause ecdysis. The

fulfilled before ecdysis is attempted. Once an attempt has begun, however, it is a 'one-chance' affair that cannot be interrupted or reinitiated. As summarized in Fig. 8, the endocrine components of this ecdysis-triggering system mirror these features of its behavioural control. The Inka cells and the VM neurones appear to be the peripheral and central halves, respectively, of a decision-making circuit. Once conditions for both halves have been satisfied, their mutually excitatory relationship ensures that ecdysis is initiated by the EH/ETH surge, but the surge can only occur once for a given moult.

Larval ecdysis in *M. sexta* is an example of a *developmentally controlled* ecdysis since pre-ecdysis and ecdysis occur at a fixed time during the developmental progression of the moult. Their onset can be delayed in a graded fashion by injecting increasing dosages of ecdysteroid that delay the terminal events of the moult (Truman *et al.* 1983). We had postulated that the declining ecdysteroid titre determined the time of ecdysis through an action on the VM neurones and EH release (Hewes and Truman, 1994). Now, it is more likely that at least some of the effects of ecdysteroids are indirect and mediated through the release of ETH. Presently, ETH secretion is the best candidate for the initiator of pre-ecdysis behaviour (Fig. 8A). Novicki and Weeks (1996) showed that abdomens isolated late in a larval moult, nevertheless, subsequently initiated pre-ecdysis behaviour and did so at the correct time. Hence, the brain and thoracic ganglia are not necessary for the proper initiation of pre-ecdysis behaviour, although they are essential for the subsequent shift to ecdysis (data above and Novicki and Weeks, 1996). Since low levels of ETH rapidly induce pre-ecdysis behaviour (Zitnan *et al.* 1996), the beginning of a low-level spontaneous release of ETH, perhaps triggered directly or indirectly by declining ecdysteroid titres, probably provides the starting signal. The levels of ETH needed to cause ecdysis are higher than those required for pre-ecdysis (Zitnan *et al.* 1996). Consequently, a rising titre of ETH, as pre-ecdysis progresses, may then begin to excite the VM neurones. In the case of larval *M. sexta*, the excitation provided by ETH is sufficient to initiate EH release (Fig. 5), but in insects such as grasshoppers, that need an appropriate ecdysis site (Hughes, 1980), the excitatory effects of ETH on the VM cells may be balanced by inhibitory inputs if the animal lacks a proper ecdysis site. With the attainment of an appropriate site, these postulated inhibitory inputs would disappear, allowing the initiation of EH release. The positive-feedback cycle between EH and ETH would then rapidly lead to the massive release of both peptides (Fig. 8B). The rapidity and completeness of peptide release from the VM cells is unprecedented for neurosecretory systems. This rapid depletion probably results from the dramatic spike-broadening seen after ETH exposure (Fig. 5B). If similar changes occur at the terminals of the VM neurones, the prolonged depolarization during the spike may provide the enhanced Ca^{2+} entry needed for the massive mobilization and release of EH. Once begun, the ETH/EH surge is apparently completed within a few minutes (Fig. 2) and is terminated

when the Inka cells and the VM neurones become exhausted of their respective peptides.

Besides developmentally controlled ecdyses, other ecdyses, such as adult ecdysis of many insects, are under a *circadian-gated control*. Under these conditions, the time of ecdysis is partially divorced from the steroid-regulated events of the moult (Truman *et al.* 1983), and brain transplantation experiments show that the timing of these ecdyses is controlled by the brain (Truman and Riddiford, 1970). Circadian-gated ecdyses are especially intriguing since the circadian clock can either advance or delay the behaviour depending on the insect's developmental state relative to the circadian cycle (Skopik and Pittendrigh, 1967). Assuming, in these cases, that ETH release is similarly linked to the developmental progression of the moult, as postulated for a larval moult, an ecdysis delay would arise from the circadian oscillator inhibiting the VM neurones if the Inka cells become ready for ecdysis during an 'intergate' period. By contrast, circadian excitation of the VM cells due to the arrival of an ecdysis gate could lead to an early peptide surge if the Inka cells had acquired their sensitivity to EH and, hence, could reinforce the excitatory effects of the circadian input. The premature arrival of the gate, however, when the Inka cells were not yet responsive, would not provide this reinforcing feedback, and the insect would have to wait for the opening of the gate on the next day before it could generate a full-scale surge.

Relationship between the triggering events and ecdysis behaviour

Both EH and ETH are completely released 10–15 min before ecdysis begins and, once begun, the ecdysis behaviour may then carry on for an additional hour or more. The key to this sustained maintenance of the ecdysis motor programme appears to be the activation of the peptidergic neurones of the cell 27/704 group (Fig. 8C). The CCAP released from these neurones in the periphery has been thought to be involved in behaviours such as air-swallowing that accompany ecdysis (Ewer *et al.* 1994; Braüning, 1990). Importantly, recent studies on desheathed abdominal ganglia of *M. sexta* show that CCAP also acts centrally to induce rapidly and reversibly the ecdysis motor programme itself (S. C. Gammie and J. W. Truman, in preparation). Thus, the sustained performance of the ecdysis motor programme appears to depend on the sustained presence of CCAP within the ganglionic neuropile.

A striking feature of the neurones in the cell 27/704 group is the elevated levels of intracellular cGMP that they show prior to ecdysis. Studies on one member of this group, cell 27, show that apart from the period around ecdysis, this cell is difficult to excite because its spike threshold is 45 mV above rest (Gammie and Truman, 1997). Importantly, the cell's threshold falls abruptly during its normal rise in cGMP level or in response to exogenous addition of the membrane-permeant form, 8-bromo-cGMP (Gammie and Truman, 1997). Thus, the appearance of cGMP in these cells appears to be a requirement for enabling these cells to fire and, hence, to release CCAP. Unlike the VM cells, however, there is

negligible spike-broadening seen in cell 27 (Gammie and Truman, 1997) and CCAP stores are not released to exhaustion. Consequently, cell 27 appears adapted to release peptide in a low-level, tonic fashion rather than in a massive phasic pulse as seen for the release of EH.

Previous experiments showed that the increase in cGMP levels in the neurones of the cell 27/704 group depended on a descending pathway from the brain (Ewer *et al.* 1994). There are a number of pieces of evidence that suggest that the descending VM neurones constitute this pathway. First, cGMP is the second messenger involved in mediating the action of EH (Morton and Truman, 1985; Morton and Giunta, 1992; Morton, 1996). Second, the appearance of cGMP in the neurones of the cell 27/704 group is invariably linked to the time of EH release (Figs 2, 6; J. Ewer and J. W. Truman, unpublished results). Third, as described in the Results, intracellular stimulation of the VM neurones results in the appearance of cGMP in members of this cell group. While cells 27 and 704 appear to respond to centrally released EH, it is unlikely that they also respond directly to circulating ETH (at least in terms of stimulating cGMP production). This conclusion is supported by the results of experiments in which the nerve cord was transected and animals were then challenged with epitracheal gland extracts. No EH release, cGMP increases and ecdysis behaviour were evident posterior to the transection, although the region showed pre-ecdysis behaviour. Hence, the pathway by which physiological levels of ETH may affect these cells is an indirect one through the ETH stimulation of the VM cells. Moreover, in order to cause EH release, ETH must act at an anterior site, most probably the brain. Whether ETH acts directly on the VM cells has not yet been determined.

Although the cell 27/704 neurones and the Inka cells show a simultaneous appearance of cGMP (Fig. 2B) in response to centrally and peripherally released EH, respectively, the subsequent time courses of this response differed markedly in the two cells. The short-lived cGMP response in the Inka cells (Fig. 2) may reflect a rapid clearing of EH from the circulation (Reynolds *et al.* 1979). The persistence of the cGMP increase in the neurones, however, is more problematic. We do not know whether this is due to the persistence of EH within the neuropile or to other factors downstream of peptide binding. Interestingly, in some insects, the duration of the cGMP response in the cell 27/704 neurones can be extended by behaviourally relevant stimuli. For example, in ecdysing first-instar grasshoppers *Locusta migratoria*, the cGMP response in these neurones remains elevated for 2–3 times its normal duration in response to stimuli that prolong the performance of ecdysis/digging behaviour (Truman *et al.* 1996). With the new realization that CCAP is a releaser of ecdysis behaviour, this extended cGMP response in these cells is probably needed for sustained CCAP release to support the continuation of the behaviour. Whether the elevated cGMP level results from the maintenance of guanylate cyclase activity or from the suppression of phosphodiesterase activity is unknown.

Applicability to other insects

EH is found throughout the insects (Truman *et al.* 1981; Horodyski *et al.* 1993; Horodyski, 1996) and is widely associated with the VM neurones. CCAP was originally identified from crustaceans (Stangier *et al.* 1987) and is now known to be present in a highly conserved set of neurones in both crustaceans and insects (Dirksen, 1994). Increases in intracellular cGMP levels are routinely seen in the CCAP neurones at ecdysis in a wide variety of insects with the notable exception of the higher Diptera, which express cGMP in a novel set of neurones (Ewer and Truman, 1996). The phylogenetic distribution of ETH is not yet known, but treatment of *Drosophila melanogaster* with synthetic *Manduca* ETH causes both EH release and ecdysis (J. D. Baker, unpublished results). Thus, the components of the ecdysis control system seen in *M. sexta* are probably ancient ones that are used widely throughout the insects.

We thank Dr J. De Vente for the rabbit and sheep anti-cGMP antisera, Professor N. J. Tublitz for the synthetic CCAP and Professor L. M. Riddiford for helpful discussions and use of equipment. This work was supported by grants from NSF (IBM 9511077) and NIH training grant (GM-07270, S.C.G.).

References

- BRAÜNIG, P. (1990). The morphology of suboesophageal ganglion cells innervating the nervus corporis cardiaci III of the locust. *Cell Tissue Res.* **260**, 95–108.
- CARLSON, J. R. (1977). The imaginal ecdysis of the cricket (*Teleogryllus oceanicus*), I. Temporal structure and organization into motor programmes. *J. comp. Physiol.* **115**, 299–317.
- COPENHAVER, P. F. AND TRUMAN, J. W. (1982). The role of eclosion hormone in the larval ecdysis of *Manduca sexta*. *J. Insect Physiol.* **28**, 695–701.
- CURTIS, A. T., HORI, M., GREEN, J. M., WOLFGANG, W. J., HIRUMA, K. AND RIDDIFORD, L. M. (1984). Ecdysteroid regulation of the onset of cuticular melanization in allatectomized and *black* mutant *Manduca sexta* larvae. *J. Insect Physiol.* **30**, 597–606.
- DAVIS, N. T., HOMBERG, U., DIRCKSEN, H., LEVINE, R. B. AND HILDEBRAND, J. G. (1993). Crustacean cardioactive peptide-immunoreactive neurons in the hawkmoth *Manduca sexta* and changes in their immunoreactivity during postembryonic development. *J. comp. Neurol.* **338**, 612–627.
- DE VENTE, J., STEINBUSCH, H. W. M. AND SCHIPPER, J. (1987). A new approach to immunocytochemistry of 3',5'-cyclic guanosine monophosphate: preparation, specificity and initial application of a new antiserum against formaldehyde-fixed 3',5'-cyclic guanosine monophosphate. *Neuroscience* **22**, 361–373.
- DIRCKSEN, H. (1994). Distribution and physiology of crustacean cardioactive peptide in arthropods. *Perspectives in Comparative Endocrinology* (ed. K. G. Davey, R. E. Peter and S. S. Tobe), pp. 139–148. Ottawa: National Research Council of Canada.
- DIRCKSEN, H., MULLER, A. AND KELLER, R. (1991). Crustacean cardioactive peptide in the nervous system of the locust, *Locusta migratoria*: an immunocytochemical study on the ventral nerve cord and peripheral innervation. *Cell Tissue Res.* **263**, 439–457.
- ELDRIDGE, R., HORODYSKI, F. M., MORTON, D. B., O'REILLY, D. R.,

- TRUMAN, J. W., RIDDIFORD, L. M. AND MILLER, L. K. (1991). Expression of an eclosion hormone gene in insect cells using baculovirus vectors. *Insect Biochem.* **21**, 341–351.
- EWER, J., DE VENTE, J. AND TRUMAN, J. W. (1994). Neuropeptide induction of cyclic GMP increases in the insect CNS: resolution at the level of single identifiable neurons. *J. Neurosci.* **14**, 7704–7712.
- EWER, J. AND TRUMAN, J. W. (1996). Increases in cyclic GMP occur at ecdysis in an evolutionarily conserved insect neuronal network. *J. comp. Neurol.* **370**, 330–341.
- GAMMIE, S. C. AND TRUMAN, J. W. (1997). An endogenous elevation of cGMP increases the excitability of identified insect neurosecretory cells. *J. comp. Physiol. A* (in press).
- HARLOW, E. AND LANE, D. B. (1988). *Antibodies. A Laboratory Manual*. Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
- HESTERLEE, S. AND MORTON, D. B. (1996). Insect physiology: the emerging story of ecdysis. *Current Biol.* **6**, 648–650.
- HEWES, R. S. AND TRUMAN, J. W. (1991). The roles of central and peripheral eclosion hormone release in the control of ecdysis behavior in *Manduca sexta*. *J. comp. Physiol. A* **168**, 697–707.
- HEWES, R. S. AND TRUMAN, J. W. (1994). Steroid regulation of excitability in identified insect neurosecretory cells. *J. Neurosci.* **14**, 1812–1819.
- HORODYSKI, F. M. (1996). Neuroendocrine control of ecdysis by eclosion hormone. *J. Insect Physiol.* **42**, 917–924.
- HORODYSKI, F. M., EWER, J., RIDDIFORD, L. M. AND TRUMAN, J. W. (1993). Isolation, characterization and expression of the eclosion hormone gene of *Drosophila melanogaster*. *Eur. J. Biochem.* **215**, 221–228.
- HUGHES, T. D. (1980). The imaginal ecdysis of the desert locust, *Schistocerca gregaria*. I. A description of the behaviour. *Physiol. Ent.* **5**, 47–54.
- MASINOVSKY, B., KEMPF, S. C., CALLAWAY, J. C. AND WILLOWS, A. O. D. (1988). Monoclonal antibodies to the molluscan small cardioactive peptide SCPB: immunolabeling of neurons in diverse invertebrates. *J. comp. Neurol.* **273**, 500–512.
- MILES, C. I. AND WEEKS, J. C. (1991). Developmental attenuation of the pre-ecdysis motor pattern in the tobacco hornworm, *Manduca sexta*. *J. comp. Physiol. A* **168**, 179–190.
- MORTON, D. B. (1996). Neuropeptide-stimulated cyclic guanosine monophosphate immunoreactivity in the neurosecretory terminals of a neurohemal organ. *J. Neurobiol.* **29**, 341–353.
- MORTON, D. B. AND GIUNTA, M. A. (1992). Eclosion hormone stimulates cyclic GMP levels in *Manduca sexta* nervous tissue via arachidonic acid metabolism with little or no contribution from the production of nitric oxide. *J. Neurochem.* **59**, 1522–1530.
- MORTON, D. B. AND TRUMAN, J. W. (1985). Steroid regulation of the peptide-mediated increase in cyclic GMP in the nervous system of the hawkmoth, *Manduca sexta*. *J. comp. Physiol. A* **157**, 423–432.
- NOVICKI, A. AND WEEKS, J. C. (1993). Organization of the larval pre-ecdysis motor pattern in the tobacco hornworm, *Manduca sexta*. *J. comp. Physiol. A* **173**, 151–162.
- NOVICKI, A. AND WEEKS, J. C. (1996). The initiation of pre-ecdysis and ecdysis behaviors in larval *Manduca sexta*: the roles of the brain, terminal ganglion and eclosion hormone. *J. exp. Biol.* **199**, 1757–1769.
- REYNOLDS, S. E. (1980). Integration of behaviour and physiology in ecdysis. *Adv. Insect Physiol.* **15**, 475–595.
- REYNOLDS, S. E., TAGHERT, P. H., AND TRUMAN, J. W. (1979). Eclosion hormone and bursicon titres and the onset of hormonal responsiveness during the last day of adult development in *Manduca sexta* (L.). *J. exp. Biol.* **78**, 77–86.
- RIDDIFORD, L. M., CURTIS, A. T. AND KIGUCHI, K. (1979). Culture of the epidermis of the tobacco hornworm *Manduca sexta*. *Tissue Culture Assn Man.* **5**, 975–985.
- SKOPIK, S. D., AND PITTENDRIGH, C. S. (1967). Circadian systems. II. The circadian oscillation in the individual *Drosophila* pupa; its independence of developmental stage. *Proc. natn. Acad. Sci. U.S.A.* **58**, 1862–1869.
- STANGIER, J., HILBICH, C., BEYREUTHER, K. AND KELLER, R. (1987). Unusual cardioactive peptide (CCAP) from pericardial organs of the shore crab, *Carcinus maenas*. *Proc. natn. Acad. Sci. U.S.A.* **84**, 575–579.
- TERZI, G., TRUMAN, J. W. AND REYNOLDS, S. E. (1988). Purification and characterization of eclosion hormone from the moth, *Manduca sexta*. *Insect Biochem.* **18**, 701–707.
- TRUMAN, J. W. (1972). Physiology of insect rhythms. I. Circadian organization of the endocrine events underlying the moulting cycle of larval tobacco hornworms. *J. exp. Biol.* **57**, 805–820.
- TRUMAN, J. W. (1978). Hormonal release of stereotyped motor programmes from the isolated nervous system of the Cecropia silkworm. *J. exp. Biol.* **74**, 151–174.
- TRUMAN, J. W. (1992). The eclosion hormone system of insects. *Progr. Brain Res.* **92**, 361–374.
- TRUMAN, J. W. AND COPENHAVER, P. F. (1989). The larval eclosion hormone neurones in *Manduca sexta*: identification of the brain–proctodeal neurosecretory system. *J. exp. Biol.* **147**, 457–470.
- TRUMAN, J. W., EWER, J. AND BALL, E. E. (1996). Dynamics of cyclic GMP levels in identified neurones during ecdysis behaviour in the locust *Locusta migratoria*. *J. exp. Biol.* **199**, 749–758.
- TRUMAN, J. W. AND RIDDIFORD, L. M. (1970). Neuroendocrine control of ecdysis in silkworms. *Science* **167**, 1624–1626.
- TRUMAN, J. W., ROUNTREE, D. B., REISS, S. E. AND SCHWARTZ, L. M. (1983). Ecdysteroids regulate the release and action of eclosion hormone in the tobacco hornworm, *Manduca sexta* (L.). *J. Insect Physiol.* **29**, 895–900.
- TRUMAN, J. W., TAGHERT, P. H., COPENHAVER, P. F., TUBLITZ, N. J. AND SCHWARTZ, L. M. (1981). Eclosion hormone may control all ecdyses in insects. *Nature* **291**, 70–71.
- TRUMAN, J. W., TAGHERT, P. H. AND REYNOLDS, S. E. (1980). Physiology of pupal ecdysis in the tobacco hornworm, *Manduca sexta*. I. Evidence for control by eclosion hormone. *J. exp. Biol.* **88**, 327–337.
- ZITNAN, D., KINGAN, T. G., HERMESMAN, J. L. AND ADAMS, M. E. (1996). Identification of ecdysis-triggering hormone from an epitracheal endocrine system. *Science* **271**, 88–91.