

Antioxidative defence alterations in skeletal muscle during prolonged acclimation to cold: role of L-arginine/NO-producing pathway

Vesna Petrović¹, Biljana Buzadžić¹, Aleksandra Korać², Ana Vasilijević¹, Aleksandra Janković¹, Ksenija Mićunović² and Bato Korać^{1,*}

¹Department of Physiology, Institute for Biological Research 'Siniša Stanković', University of Belgrade, Bulevar Despota Stefana 142, 11060 Belgrade, Serbia and ²Institute of Zoology, Faculty of Biology, University of Belgrade, Studentski trg 16, 11000 Belgrade, Serbia

*Author for correspondence (e-mail: koracb@ibiss.bg.ac.yu)

Accepted 18 October 2007

SUMMARY

Early in cold acclimation (1–7 days), heat is produced by shivering, while late in cold acclimation (12–45 days), skeletal muscle contributes to thermogenesis by tissue metabolism other than contractions. Given that both thermogenic phases augment skeletal muscle aerobic power and reactive species production, we aimed in this study to examine possible changes in skeletal muscle antioxidative defence (AD) during early and late cold acclimation with special emphasis on the influence of the L-arginine/nitric oxide (NO)-producing pathway on the modulation of AD in this tissue. Adult Mill Hill hybrid hooded rat males were divided into two main groups: a control group, which was kept at room temperature (22±1°C), and a group maintained at 4±1°C for 45 days. The cold-acclimated group was divided into three subgroups: untreated, L-arginine treated and N^ω-nitro-L-arginine methyl ester (L-NAME) treated. The AD parameters were determined in the gastrocnemius muscle on day 1, 3, 7, 12, 21 and 45 of cold acclimation. The results showed an improvement of skeletal muscle AD in both early and late cold acclimation. Clear phase-dependent changes were seen only in copper, zinc superoxide dismutase activity, which was increased in early cold acclimation but returned to the control level in late acclimation. In contrast, there were no phase-dependent changes in manganese superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase and glutathione S-transferase, the activities of which were increased during the whole cold exposure, indicating their engagement in both thermogenic phases. L-Arginine in early cold acclimation accelerated the cold-induced AD response, while in the late phase it sustained increases achieved in the early period. L-NAME affected both early and late acclimation through attenuation and a decrease in the AD response. These data strongly suggest the involvement of the L-arginine/NO pathway in the modulation of skeletal muscle AD.

Key words: skeletal muscle, antioxidative defence, nitric oxide, cold.

INTRODUCTION

Prolonged exposure of mammals to cold induces increases in heat production and enhances tolerance to cold through both shivering and non-shivering thermogenesis (NST) (Himms-Hagen, 1976). Shivering thermogenesis, achieved by increased skeletal muscle activity, occurs early in response to cold exposure (Jansky and Hart, 1968). A later decrease in shivering thermogenesis has been found to be accompanied by an increase in NST (Griggio, 1982). NST is achieved by the uncoupling of oxidative metabolism from ATP production, primarily in brown adipose tissue (Jansky, 1966; Foster and Frydman, 1979). In contrast to the clear role of skeletal muscle in shivering thermogenesis, the extent of NST in this tissue remains controversial, with some reports demonstrating NST in the skeletal muscle of pigeons (Skulachev and Maslov, 1960), ducklings (Barre et al., 1987) and rats (Mollica et al., 2005), whilst others found no NST in this tissue (Golozoubova et al., 2001). However, skeletal muscle, which represents a large percentage of body mass, significantly contributes to the intensification of metabolic activity of the whole organism during thermogenesis (Rolfe and Brand, 1996) due to its huge capacity for β -oxidation and the oxidative capacity of mitochondria (Hoppeler and Fluck, 2003).

Heat production increases metabolic rate and oxygen consumption in metabolically active tissues (Shiota and Masumi,

1988) that is implacably associated with the elevation of reactive oxygen species generation. Cellular homeostasis under conditions of increased reactive oxygen species production is achieved by a proportional increase in tissue antioxidative defence (AD) (Halliwell and Gutteridge, 1990; Buzadžić et al., 1997; Buzadžić et al., 1999; Korać and Buzadžić, 2001; Petrović et al., 2006). AD consists of enzymes – copper, zinc and manganese superoxide dismutase (CuZnSOD and MnSOD, EC 1.15.1.1), catalase (CAT, EC 1.11.1.6), glutathione peroxidase (GSH-Px, EC 1.11.1.9), glutathione S-transferase (GST, EC 2.5.1.18), thioredoxin reductase (TR, EC 1.6.4.5), glutathione reductase (GR, EC 1.6.4.2) and low molecular mass antioxidants such as vitamins E and C, glutathione (GSH), etc. (Chance et al., 1979; Cadenas et al., 1989; Aruoma, 1996). In recent years, considerable progress has been achieved in the field of redox regulation and the concept of the protective effect of AD was extended to its regulatory role, since AD activity determines reactive species availability (Mugge et al., 1991; Korać and Buzadžić, 2000; Buzadžić et al., 2006).

Reactive species in moderate concentrations, especially superoxide anion radical (O₂^{•-}) and nitric oxide (NO), play an important role as regulatory mediators in biological processes (Dröge, 2002). In skeletal muscle, NO is produced by the activity of constitutively expressed endothelial and neuronal NO synthase

(NOS), localized on the surface of the sarcolemmal membrane and the endothelial plasmalemmal caveolae, as well as by inducible NOS, the localization of which varies depending on the state of the cell (Kobzik et al., 1995; Bates et al., 1996). NO regulates many physiological functions of skeletal muscle including glucose uptake and oxidation (Young et al., 1997), mitochondriogenesis (Puigserver et al., 1998), contractile functions (Joneschild et al., 1999; Maréchal and Gailly, 1999), blood flow (Brevetti et al., 2003) and fatty acid oxidation (Jobgen et al., 2006), as well as muscle repair through satellite cell activation and the release of myotrophic factors (Anderson, 2000; Brunelli et al., 2007).

On the other hand, it is known that both endurance training and acclimation to cold increase skeletal muscle oxygen uptake and, consequently, production of reactive oxygen species. So far, most studies have focused on the changes of skeletal muscle AD during exercise (Tiidus et al., 1996; Tonkonogi et al., 2000; Pansarasa et al., 2002). However, knowledge of the changes in AD in skeletal muscle during acclimation to cold is very limited and there are no data concerning the possible role of NO in the modulation of skeletal muscle AD.

Hence, the aim of the present study was to assess changes in AD in skeletal muscle during early and late cold acclimation with a special focus on the possible role of NO in the modulation of AD in this tissue. For this purpose, adult male rats were kept at room temperature, or were exposed to cold for 45 days and received the NO-manipulating agents L-arginine or *N*^ω-nitro-L-arginine methyl ester (L-NAME) as drinking liquid, and the effects on AD over time were assessed.

MATERIALS AND METHODS

Mill Hill hybrid hooded, 4 month old *Rattus norvegicus* (Berkenhout 1769) males were divided into two main groups: a control group, kept at room temperature (22±1°C) for the duration of the experiment, and a second group, maintained at 4±1°C. The cold-acclimated group was divided into three subgroups: (1) untreated, (2) L-arginine treated and (3) L-NAME treated. Drugs were administered as a drinking liquid, as done previously by others (Saha et al., 1996) and by us (Vasiljević et al., 2007) in a dose (2.25% L-arginine HCl and 0.01% L-NAME HCl in tap water) that caused no toxic effects (Petrović et al., 2005). The animals were killed at the same time of the day after 1, 3, 7, 12, 21 or 45 days of cold acclimation. The rats were housed in individual plastic cages with drinking liquid and food *ad libitum*. Each experimental group consisted of six individuals. Body mass, food and fluid intake were recorded daily for each animal. The experimental protocol was approved by the Ethical Committee for the Treatment of Experimental Animals of the Institute for Biological Research, Belgrade.

The rats were killed by decapitation, and the gastrocnemius dissected and rinsed with physiological saline to wash out traces of blood. The tissue was homogenized (Ultra/Turrax homogenizer, Janke und Kunkel Ka/Werke, Staufen, Germany; 0–4°C) in 0.25 mol l⁻¹ sucrose, 0.1 mmol l⁻¹ EDTA and 50 mmol l⁻¹ Tris-HCl buffer, pH 7.4, and the homogenates sonicated (Takada et al., 1982).

Activity of antioxidative enzymes

Total SOD activity was examined by a modified method of Misra and Fridovich (Misra and Fridovich, 1972). MnSOD activity was determined after preincubation with 4 mmol l⁻¹ KCN. CuZnSOD activity was calculated as the difference between total SOD and MnSOD activities. Enzymatic activity was expressed in U mg⁻¹

protein. SOD units were defined as the amount of the enzyme inhibiting epinephrine (adrenaline) auto-oxidation under appropriate reaction conditions. CAT was assayed as suggested by the supplier (Sigma-Aldrich, St Louis, MO, USA) and the activity expressed in μmol H₂O₂ min⁻¹ mg⁻¹ protein. GSH-Px was determined with *t*-butylhydroperoxide as a substrate (Paglia and Valentine, 1967) and the activity expressed in nmol of reduced NADPH min⁻¹ mg⁻¹ protein. GST was measured by the method of Habig et al. (Habig et al., 1974) and the activity expressed in nmol GSH used min⁻¹ mg⁻¹ protein. GR activity was assayed according to Glatzle et al. (Glatzle et al., 1974) and expressed as nmol GSH min⁻¹ mg⁻¹ protein.

Determination of GSH

The content of GSH was examined in the tissue after deproteinization with sulphosalicylic acid. Total GSH was measured by enzyme recycling assay according to Griffith (Griffith, 1980) and expressed in nmol GSH g⁻¹ tissue.

Other assays and statistics

Protein content was estimated by the method of Lowry et al. (Lowry et al., 1951) using bovine serum albumin as a reference. Analysis of variance (ANOVA) was used for within-group comparison of the data. If the *F* test showed an overall difference, Tukey's test was applied to identify significant differences. Statistical significance was accepted at *P*<0.05.

RESULTS

Table 1 shows the changes in body mass during acclimation. Initially, in cold acclimation animals from all examined groups lost body mass (*P*<0.001). These decreases persisted during the whole of the early cold-acclimation period and body mass returned to the control level no sooner than day 45 of late cold acclimation.

During the whole experiment no changes in fluid intake were observed. In contrast, as acclimation to cold started, animals in all examined groups increased food intake by 100% and this increase remained constant to the end of experiment (data not shown).

Generally, the results showed improvement of skeletal muscle AD in both early and late cold acclimation. Clear phase-dependent changes were seen only in CuZnSOD activity (Fig. 1), which was increased in early cold acclimation but returned to the control level in late acclimation. In contrast, there were no phase-dependent changes in MnSOD (Fig. 2) and peroxidative- and GSH-related parts of AD (CAT, GSH-Px, GR and GST; Figs 3–6), the activities of which were increased during the whole period of cold exposure.

Alterations in CuZnSOD activity in gastrocnemius are shown in Fig. 1. In the untreated group, changes in CuZnSOD activity were quite phase dependent, i.e. an increase in its activity was observed

Table 1. Changes in body mass during cold acclimation

Day	Control	Cold		
		Untreated	L-arginine	L-NAME
1	2.0±1.0	-6.8±1.4***	-4.7±1.7***	-1.0±0.2***
3	4.7±2.8	-3.4±1.6**	-2.3±2.9*	-1.8±1.5**
7	8.2±1.7	-11.5±3.5***	-11.3±4.4***	-16.0±5.8***
12	14.4±4.4	-8.7±2.7***	-5.6±12.2***	2.7±3.3***
21	24.5±4.3	8.7±3.2***	9.5±3.3***	4.3±5.1***
45	49.0±16.0	42.8±9.6	39.0±16.9	49.0±4.4

Data for body mass gain (in g) are means ± s.e.m. (*N*=6 in all groups).

P*<0.05, *P*<0.01, ****P*<0.001 compared with the control acclimated to 22°C for the same time period.




Fig. 1. The effect of cold-exposure ($4\pm 1^\circ\text{C}$) duration on copper, zinc superoxide dismutase (CuZnSOD) activity in the skeletal muscle of untreated, L-arginine-treated and *N*^o-nitro-L-arginine methyl ester (L-NAME)-treated animals. * $P<0.05$, ** $P<0.01$, *** $P<0.001$ compared with the control kept at room temperature ($22\pm 1^\circ\text{C}$). # $P<0.05$, ## $P<0.01$, ### $P<0.001$ compared with the untreated group at the same time of cold acclimation.

during early cold exposure starting from day 3, which returned to the control level during late cold acclimation. On the other hand, in the L-arginine-treated group the activity of this enzyme increased from day 1 ($P<0.05$) in the early phase, but decreased below the control level after day 21 ($P<0.001$) of the late cold exposure, compared with both the control and the untreated group. In L-NAME-treated animals, CuZnSOD activity was decreased on day 1 of early cold exposure ($P<0.001$) and remained decreased until the end of cold exposure in comparison with the control and untreated group.

Changes in MnSOD activity during cold acclimation are depicted in Fig. 2. As shown, in the untreated group, MnSOD activity increased from day 3 ($P<0.01$), while in L-arginine-treated rats, MnSOD activity was increased on day 1 of early cold exposure compared with the control ($P<0.01$) and untreated group ($P<0.05$). During the late cold acclimation, increase in MnSOD activity achieved early was sustained in the untreated group and was unaffected by L-arginine, related to the untreated group. L-NAME treatment decreased MnSOD activity from day 3 to day 12 compared with the untreated group, and delayed the cold-induced increase in MnSOD activity on day 21 of late cold-acclimation ($P<0.001$).

CAT activity (Fig. 3) progressively increased in all three cold-acclimated groups with the highest increase on day 7 ($P<0.001$). A higher CAT activity was observed on day 3 of the early phase of acclimation in both untreated ($P<0.05$) and L-NAME-treated rats ($P<0.001$), while increased activity of this enzyme was recorded as early as day 1 of exposure to cold in L-arginine-treated animals ($P<0.05$) compared with the control ($P<0.05$) and untreated group ($P<0.05$). CAT activity remained elevated during the whole of the late cold acclimation in both untreated and L-arginine-treated rats, while in the L-NAME-treated group it returned to control values on day 21 and decreased compared with the untreated group at days 21 ($P<0.01$) and 45 ($P<0.001$) of late cold acclimation.




Fig. 2. Time course of changes in manganese superoxide dismutase (MnSOD) activity in skeletal muscle of untreated, L-arginine-treated and L-NAME-treated rats during early and late acclimation to cold ($4\pm 1^\circ\text{C}$). * $P<0.05$, ** $P<0.01$, *** $P<0.001$ compared with the control maintained at room temperature ($22\pm 1^\circ\text{C}$). # $P<0.05$, ## $P<0.01$, ### $P<0.001$ compared with the untreated group at the same time of cold acclimation.

Time-dependent changes of GSH-Px activity are shown in Fig. 4. Similar time-course changes in the activity of this enzyme were observed in all three cold-acclimated groups, during both the early and late phases, with the highest increase seen on day 7 in untreated ($P<0.001$) and L-NAME-treated animals ($P<0.01$). However, in the L-arginine-treated group the maximal increase of GSH-Px activity was recorded on day 3 of early cold exposure and it was higher ($P<0.05$) compared with that observed at the same




Fig. 3. Time-dependent changes in catalase (CAT) activity in the skeletal muscle of untreated, L-arginine-treated and L-NAME-treated rats during acclimation to cold ($4\pm 1^\circ\text{C}$). * $P<0.05$, ** $P<0.01$, *** $P<0.001$ compared with the control acclimated to room temperature ($22\pm 1^\circ\text{C}$). # $P<0.05$, ## $P<0.01$, ### $P<0.001$ compared with the untreated group at the same time of cold acclimation.




Fig. 4. Time course of changes in glutathione peroxidase (GSH-Px) activity in the skeletal muscle of untreated, L-arginine-treated and L-NAME-treated animals during exposure to cold ($4\pm 1^\circ\text{C}$). $*P<0.05$, $**P<0.01$, $***P<0.001$ compared with the control maintained at room temperature ($22\pm 1^\circ\text{C}$). $\#P<0.05$, $\##P<0.01$, $\###P<0.001$ compared with the untreated group at the same time of cold acclimation.




Fig. 5. The effect of short and prolonged cold exposure ($4\pm 1^\circ\text{C}$) on glutathione S-transferase (GST) activity in the skeletal muscle of untreated, L-arginine-treated and L-NAME-treated rats. $**P<0.01$, $***P<0.001$ compared with the control acclimated to room temperature ($22\pm 1^\circ\text{C}$).

time in the untreated group. The increase in GSH-Px activity originated in the early cold acclimation remained during the whole late cold acclimation in untreated and L-arginine-treated groups, while it returned to the control level on day 21 in L-NAME-treated rats and stayed below values observed in the untreated group at days 21 ($P<0.001$) and 45 ($P<0.01$).

As shown, L-arginine in early cold acclimation accelerated the cold-induced increase in MnSOD, CuZnSOD and CAT activity, while in the late phase it sustained increases achieved in the early period. L-NAME affected both early and late acclimation through attenuation and decreases in the AD response. That is, L-NAME treatment postponed the cold-induced increase of MnSOD activity and reversed CAT and GSH-Px activities to the control level on day 21 of late cold acclimation, and decreased CuZnSOD activity during the whole acclimation period. In contrast, neither L-arginine nor L-NAME changed GR and GST activities, which increased from day 1 of cold acclimation and stayed unchanged until the end of the experiment.

From Fig. 5 it can be seen that GST activity was increased in the untreated group on day 1 of early cold exposure ($P<0.001$) and remained elevated for the whole duration of the experiment compared with the control, with the maximum reached on day 7 of cold exposure ($P<0.001$). Both L-arginine and L-NAME failed to affect changes in GST activity seen in the untreated group.

GR activity (Fig. 6) showed time-dependent changes similar to those of GST activity, i.e. it was increased on day 1 of the early phase in all examined groups and remained elevated during the whole period of both early and late cold acclimation, with the maximum increase on day 7 ($P<0.001$).

GSH content in the gastrocnemius (Fig. 7) was decreased from day 1 of early cold exposure in all examined groups, and remained decreased during the whole period of cold acclimation in L-NAME-treated rats in relation to the control. However, in the untreated group the GSH content was restored to the control level on day 45 of late cold acclimation, while in the L-arginine-treated group it

restitution was observed on day 3 of early cold acclimation and remained unchanged until the end of the experiment.

DISCUSSION

In the present study, clear phase-dependent changes were seen only in CuZnSOD activity, which was increased in early cold acclimation (1–7 days) but returned to control in late acclimation (12–45 days). In contrast, there were no phase-dependent changes in MnSOD activity and the peroxidative- and GSH-related part of AD – CAT, GSH-Px, GR and GST – the activities of which were




Fig. 6. Changes in glutathione reductase (GR) activity in the skeletal muscle of untreated, L-arginine-treated and L-NAME-treated animals during early and late cold acclimation ($4\pm 1^\circ\text{C}$). $*P<0.05$, $**P<0.01$, $***P<0.001$ compared with the control kept at room temperature ($22\pm 1^\circ\text{C}$).




Fig. 7. Changes in GSH content in the skeletal muscle of untreated, L-arginine-treated and L-NAME-treated rats after different time periods of cold exposure ($4\pm 1^\circ\text{C}$). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with the control acclimated to room temperature ($22\pm 1^\circ\text{C}$). ## $P < 0.01$; ### $P < 0.001$ compared with the untreated group at the same time of cold acclimation.

increased during the whole cold exposure, owing to their functional engagement in both phases of thermogenesis. L-Arginine treatment exerted effects on early cold acclimation by acceleration of the cold-induced AD response and in the late phase it sustained increases achieved in the early period. In contrast, L-NAME attenuated and decreased the cold-induced skeletal muscle AD response in both early and late acclimation. The effects that these treatments exhibited on AD in both phases of thermogenesis are discussed in more detail below.

In the course of the early period of cold exposure, shivering thermogenesis is activated. During this time, to overcome cold, nerve stimulation induces permanent skeletal muscle contractions and relaxation. This period may therefore be seen as muscular training, which is known to increase the rate of oxygen consumption (Wickler, 1981) and the proportion of the O_2^- and H_2O_2 formed from oxygen utilized by mitochondria (Vasilaki et al., 2006). Accordingly, increases in MnSOD and CuZnSOD as well as CAT and GSH-Px activities observed in the untreated group during early cold acclimation could be explained as a response of these enzymes to an increased production of their substrates and are in line with several reports related to exercise-improved AD (Higuchi et al., 1985; Pansarasa et al., 2002). Supplementation with the physiological substrate for NOSs, L-arginine, in the early phase resulted in an acceleration of the increase in cold-induced AD enzymes (MnSOD, CuZnSOD, CAT and GSH-Px) activities, and restoration of GSH level compared with the untreated group. In contrast, L-NAME, the non-selective NOSs inhibitor, postponed the cold-induced increase in MnSOD activity seen in the untreated group in the early phase to day 21 of late cold exposure and decreased CuZnSOD activity during the acclimation to cold. It was found that L-arginine enhanced exercise-induced NOSs activation and NO production, which improves aerobic capacity (Maxwell et al., 2001), mechanical and metabolic muscle capability (Maréchal and Gailly, 1999; Ohta et al., 2007) and increases glucose oxidation (Fu et al., 2005). In contrast, L-NAME, through inhibition of NOSs and a decrease in

NO availability, diminished skeletal muscle contractions, as well as glucose uptake and oxidation (Roberts et al., 1997). Thus, the accelerated increase in MnSOD, CuZnSOD and CAT activities observed here in L-arginine-treated rats during the early acclimation to cold could be connected to improved skeletal muscle contraction and oxidative capacity by NO and, consequently, to O_2^- and H_2O_2 production.

However, after a period of 7 (Cannon and Nedergaard, 2004) or 10 days (Peralta et al., 2003) on cold, the impact of shivering decreases and cannot account for thermogenesis (Griggio, 1982), while NST markedly increases. During prolonged exposure to cold, despite decreased shivering, skeletal muscle retains an enhanced capacity for aerobic support of energy metabolism (although, less than in shivering), i.e. working as a supportive tissue to supply brown adipose tissue with oxygen, contributing to the overall acclimation to cold (Wickler, 1981; Wibom et al., 1992). In our study, after 7 days, CuZnSOD activity in the untreated group was restored to the control level, while MnSOD activity remained increased during the whole acclimation period. The different responses of SOD isoforms seen after early cold acclimation could be connected to their different subcellular organization; MnSOD is compartmentalized in the mitochondrial inner membrane, whereas CuZnSOD is found mainly in cytosol. Thus, it might be proposed that in the period when shivering ceases, production of O_2^- is such that mitochondrial MnSOD is sufficient to cope with it and to prevent an increase of O_2^- in the cytosol and a competent response in CuZnSOD activity.

The observed changes in enzyme activities during cold acclimation could be interpreted in terms of different tissue conditions characterized by different fuel utilization. It is known that maximal utilization of glucose by skeletal muscle is associated with acute cold exposure, while in long-term cold acclimation a marked increase in the utilization of lipid substrates occurs (Vallerand et al., 1990). Accordingly, a conspicuous increase of CAT and GSH-Px activities recorded here from day 7 of cold acclimation could be attributed to increased β -oxidation of fatty acids and, connected to that, increased H_2O_2 production. These data are in line with studies on various tissues that reported increases in CAT and GSH-Px activity during cold exposure, as a response to increased peroxidative pressure (Alptekin et al., 1996; Buzadžić et al., 1997; Selman et al., 2000).

L-arginine treatment sustained the shown increases in AD achieved in the early period until the end of cold acclimation. In contrast, L-NAME treatment postponed the cold-induced increase of MnSOD activity and reversed CAT and GSH-Px activities to the control level on day 21 of late cold exposure. These data showing attenuated and delayed cold-induced responses in AD components by L-NAME treatment in late cold acclimation strongly indicate the involvement of the L-arginine/NO pathway in the modulation of skeletal muscle AD. At this stage, considering our data, we cannot say how L-arginine/NO affects skeletal muscle AD. Fu et al. (Fu et al., 2005) showed that L-arginine, by increasing NOSs-derived NO production, enhances expression of the genes responsible for fatty acid oxidation and, thereby, H_2O_2 production. In contrast, Nagase et al. (Nagase et al., 1997) showed that incubation of L-arginine under peroxidative conditions leads to the non-enzymatic production of NO-dependent species, which are known to increase production of O_2^- and H_2O_2 (Navarro et al., 2005). Peralta et al. (Peralta et al., 2003) reported that L-arginine/NO, by mitochondrial NOS (mtNOS) activation, in early cold acclimation increased O_2^- uptake but significantly decreased it at day 24 of late cold exposure via inhibition of cytochrome *c* oxidase and an increase in O_2^-

production. Additionally, these authors emphasized that in skeletal muscle the major NOS isoform that participates in the response to cold acclimation is mtNOS, which in this tissue was described varyingly as post-translationally modified nNOS (Elfering et al., 2002) or as eNOS (Punkt et al., 2006). In the present study, restoration of GSH level was seen in the L-arginine-treated group at day 3 and in the untreated group at day 45, while in the L-NAME-treated group it stayed below the control level during the whole period of cold acclimation. Thus, it could be hypothesized that L-arginine, i.e. NO, acted by restoring the GSH amount to the control level. This hypothesis is based on the results of Moellering et al. (Moellering et al., 1998), who reported induction of GSH synthesis in skeletal muscle as well as in endothelial cells (Moellering et al., 1999) in response to NO via the activation of γ -glutamyl cysteine synthetase. Further studies along these lines are necessary for clarification of the precise mechanisms involved. The data presented here demonstrating that the cold-induced AD response is improved by L-arginine but attenuated by L-NAME clearly showed opposing effects of L-arginine and L-NAME on the modulation of AD in skeletal muscle. However, there are several reports, with different model systems, concerning the same effect of L-NAME treatment as that seen after L-arginine treatment. Conflicting observations in experiments with the NOSs inhibitor L-NAME are explained by non-enzymatic production of NO during incubation with L-NAME in the presence of NADPH, GSH and ascorbate (Moroz et al., 1998). Henningson et al. (Henningson et al., 2000) showed that chronic L-NAME treatment in diabetic rats blocked constitutive NOSs that evoked iNOS-derived NO production to compensate for NO level, while Krippel-Drews et al. (Krippel-Drews et al., 1996) reported that L-NAME can also act directly, not via intracellular messengers. We have recently reported that L-arginine and L-NAME affect pancreatic AD in the same manner (Vasiljević et al., 2007). Thus, tissue-specific effects of these two NO-manipulating agents could be proposed.

The presented results clearly show improvement of skeletal muscle AD in both early and late cold acclimation, indicating intensive oxidative metabolism in this tissue during both shivering and NST. Moreover, L-arginine in early cold acclimation accelerated the cold-induced AD response, while in the late phase it simply sustained increases achieved in the early period. L-NAME exerted effects on both early and late acclimation through attenuation and a decrease in the AD response. These data strongly suggest the involvement of the L-arginine/NO pathway in the modulation, i.e. the improvement of skeletal muscle AD during cold acclimation. However, in order to understand the precise mechanisms, additional studies, primarily related to the determination of the relative contributions of the L-arginine-dependent and NOSs-dependent effects, are needed. Our efforts along these lines are in progress.

This work was supported by the Ministry of Science, Republic of Serbia, grant no. 143050.

REFERENCES

- Alptekin, N., Seckin, S., Dogru-Abbasoglu, S., Yelkenci, F., Kocak-Toker, N., Toker, G. and Uysal, M. (1996). Lipid peroxides, glutathione gamma-glutamylcysteine synthetase and gamma-glutamyltranspeptidase activities in several tissues of rats following water-immersion stress. *Pharmacol. Res.* **34**, 167-169.
- Anderson, J. E. (2000). A role for nitric oxide in muscle repair: nitric oxide-mediated activation of muscle satellite cells. *Mol. Biol. Cell* **11**, 1859-1874.
- Aruoma, I. E. (1996). Characterization of drugs as antioxidant prophylactics. *Free Radic. Biol. Med.* **20**, 675-705.
- Barre, H., Bailly, L. and Rouanet, J. L. (1987). Increased oxidative capacity in skeletal muscle from cold-acclimated ducklings: a comparison with rats. *Comp. Biochem. Physiol.* **88B**, 519-522.
- Bates, T. E., Loesch, A., Burnstock, G. and Clark, J. B. (1996). Mitochondrial nitric oxide synthase: a ubiquitous regulator of oxidative phosphorylation? *Biochem. Biophys. Res. Commun.* **218**, 40-44.
- Brevetti, L. S., Chang, S. D., Tang, G. L., Sarkar, R. and Messina, L. M. (2003). Overexpression of endothelial nitric oxide synthase increases skeletal muscle blood flow and oxygenation in severe rat hind limb ischemia. *J. Vasc. Surg.* **38**, 820-826.
- Brunelli, S., Csiorati, C., D'Antona, G., Innocenzi, A., Covarello, D., Galvez, B. G., Perrotta, C., Monopoli, A., Sanvito, F., Bottinelli, R. et al. (2007). Nitric oxide release combined with nonsteroidal antiinflammatory activity prevents muscular dystrophy pathology and enhances stem cell therapy. *Proc. Natl. Acad. Sci. USA* **104**, 264-269.
- Buzadžić, B., Blagojević, D., Korać, B., Saičić, Z. S., Spasić, M. B. and Petrović, V. M. (1997). Seasonal variation in the antioxidant defense system of the brain of the ground squirrel (*Citellus citellus*) and response to low temperature compared with rat. *Comp. Biochem. Physiol.* **117C**, 141-149.
- Buzadžić, B., Korać, B. and Petrović, V. M. (1999). The effect of adaptation to cold and re-adaptation to room temperature on the level of glutathione in rat tissues. *J. Therm. Biol.* **24**, 373-377.
- Buzadžić, B., Korać, A., Petrović, V. and Korać, B. (2006). Redox regulation of brown adipocytes: molecular and cellular targets in tissue remodeling. *Acta Physiol. Pharmacol. Serb.* **42**, 141-159.
- Cadenas, E., Simić, M. G. and Sies, H. (1989). Antioxidant activity of 5-hydroxytryptophan, 5-hydroxyindole, and DOPA against microsomal lipid peroxidation and its dependence on vitamin E. *Free Radic. Res. Commun.* **6**, 11-17.
- Cannon, B. and Nedergaard, J. (2004). Brown adipose tissue: function and physiological significance. *Physiol. Rev.* **84**, 277-359.
- Chance, B., Sies, H. and Boveris, A. (1979). Hydroperoxide metabolism in mammalian organs. *Physiol. Rev.* **59**, 527-605.
- Dröge, W. (2002). Free radicals in the physiological control of cell function. *Physiol. Rev.* **82**, 47-95.
- Elfering, S. L., Sarkela, T. M. and Giulivi, C. (2002). Biochemistry of mitochondrial nitric oxide synthase. *J. Biol. Chem.* **277**, 38079-38086.
- Foster, D. O. and Frydman, M. L. (1979). Tissue distribution of cold-induced thermogenesis in conscious warm- or cold-acclimated rats reevaluated from changes in tissue blood flow: the dominant role of brown adipose tissue in the replacement of shivering by nonshivering thermogenesis. *Can. J. Physiol. Pharmacol.* **57**, 257-270.
- Fu, W. J., Haynes, T. E., Kohli, R., Hu, J., Shi, W., Spencer, T. E., Carroll, R. J., Meininger, C. J. and Wu, G. (2005). Dietary L-arginine supplementation reduces fat mass in Zucker diabetic fatty rats. *J. Nutr.* **135**, 714-721.
- Glatzle, D., Vuilleumier, J. P., Weber, F. and Decker, K. (1974). Glutathione reductase test with whole blood, a convenient procedure for the assessment of the riboflavin status in humans. *Experientia* **30**, 665-667.
- Golozubova, V., Hohtola, E., Matthias, A., Jacobsson, A., Cannon, B. and Nedergaard, J. (2001). Only UCP1 can mediate adaptive nonshivering thermogenesis in the cold. *FASEB J.* **15**, 2048-2050.
- Griffith, O. W. (1980). Determination of glutathione and glutathione disulfide using glutathione reductase and 2-vinylpyridine. *Anal. Biochem.* **106**, 207-212.
- Griggio, M. A. (1982). The participation of shivering and nonshivering thermogenesis in warm and cold-acclimated rats. *Comp. Biochem. Physiol.* **73A**, 481-484.
- Habig, W. H., Pabst, M. J. and Jakoby, W. B. (1974). Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J. Biol. Chem.* **249**, 7130-7139.
- Halliwell, B. and Gutteridge, J. M. (1990). The antioxidants of human extracellular fluids. *Arch. Biochem. Biophys.* **280**, 1-8.
- Henningson, R., Alm, P., Lindstrom, E. and Lindquist, I. (2000). Chronic blockade of NO synthase paradoxically increases islet NO production and modulates islet hormone release. *Am. J. Physiol.* **279**, 95-107.
- Higuchi, M., Cartier, L. J., Chen, M. and Holloszy, J. O. (1985). Superoxide dismutase and catalase in skeletal muscle: adaptive response to exercise. *J. Gerontol.* **40**, 281-286.
- Himms-Hagen, J. (1976). Cellular thermogenesis. *Annu. Rev. Physiol.* **38**, 315-351.
- Hoppeler, H. and Fluck, M. (2003). Plasticity of skeletal muscle mitochondria: structure and function. *Med. Sci. Sports Exerc.* **35**, 95-104.
- Jansky, L. (1966). Body organ thermogenesis of the rat during exposure to cold and at maximal metabolic rate. *Fed. Proc.* **25**, 1297-1305.
- Jansky, L. and Hart, J. S. (1968). Cardiac output and organ blood flow in warm- and cold-acclimated rats exposed to cold. *Can. J. Physiol. Pharmacol.* **46**, 653-659.
- Jobgen, W. H., Fried, S. K., Fu, W. J., Meininger, C. J. and Wu, G. (2006). Regulatory role for the arginine-nitric oxide pathway in metabolism of energy substrates. *J. Nutr. Biochem.* **17**, 571-588.
- Joneschild, E. S., Chen, L. E., Seaber, A. V., Frankel, E. S. and Urbaniak, J. R. (1999). Effect of a NOS inhibitor, L-NMMA, on the contractile function of reperfused skeletal muscle. *J. Reconstr. Microsurg.* **15**, 55-60.
- Kobzik, L., Stringer, B., Balligand, J. L., Reid, M. B. and Stamler, J. S. (1995). Endothelial type nitric oxide synthase in skeletal muscle fibers: mitochondrial relationships. *Biochem. Biophys. Res. Commun.* **211**, 375-381.
- Korać, B. and Buzadžić, B. (2000). Oxidative stress and antioxidative protection in skin. *Yugoslav. Med. Biochem.* **19**, 363-379.
- Korać, B. and Buzadžić, B. (2001). Doxorubicin toxicity to the skin: possibility of protection with antioxidants enriched yeast. *J. Dermatol. Sci.* **25**, 45-52.
- Krippel-Drews, P., Welkin, S. and Drews, G. (1996). Effects of the nitric oxide synthase inhibitor N-omega-nitro-L-arginine methyl ester on electrical activity and ion channels of mouse pancreatic B cells. *Biochem. Biophys. Res. Commun.* **224**, 199-205.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. and Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265-275.
- Maréchal, G. and Gailly, P. (1999). Effects of nitric oxide on the contraction of skeletal muscle. *Cell. Mol. Life Sci.* **55**, 1088-1102.

- Maxwell, A. J., Ho, H. V., Le C. Q., Lin, P. S., Berstein, D. and Cooke, J. P. (2001). L-arginine enhances aerobic capacity in association with augmented nitric oxide production. *J. Appl. Physiol.* **90**, 933-938.
- Misra, H. P. and Fridovich, I. (1972). The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J. Biol. Chem.* **247**, 3170-3175.
- Moellering, D., McAndrew, J., Patel, R. P., Cornwell, T., Lincoln, T., Cao, X., Messina, J. L., Forman, H. J., Jo, H. and Darley-Usmar, V. M. (1998). Nitric oxide-dependent induction of glutathione synthesis through increased expression of γ -glutamylcysteine synthetase. *Arch. Biochem. Biophys.* **358**, 74-82.
- Moellering, D., McAndrew, J., Patel, R. P., Forman, H. J., Mulcahy, R. T., Jo, H. and Darley-Usmar, V. M. (1999). The induction of GSH synthesis by nanomolar concentrations of NO in endothelial cells: a role for gamma glutamylcysteine synthetase and gamma glutamyl transpeptidase. *FEBS Lett.* **448**, 292-296.
- Mollica, M. P., Lionetti, L., Crescenzo, R., Tasso, R., Barletta, A., Liverini, G. and Iossa, S. (2005). Cold exposure differently influences mitochondrial energy efficiency in rat liver and skeletal muscle. *FEBS Lett.* **579**, 1978-1982.
- Moroz, L. L., Norby, S. W., Cruz, L., Sweedler, J. V., Gillette, R. and Clarkson, R. B. (1998). Non-enzymatic production of nitric oxide (NO) from NO synthase inhibitors. *Biochem. Biophys. Res. Commun.* **253**, 571-576.
- Mugge, A., Elwell, J. K., Peterson, T. E. and Harrison, D. G. (1991). Release of intact endothelium-derived relaxing factor depends on endothelial superoxide dismutase activity. *Am. J. Physiol.* **260**, 219-225.
- Nagase, S., Takemura, K., Ueda, A., Hirayama, A., Aoyagi, K., Kondoh, M. and Koyama, A. (1997). A novel nonenzymatic pathway for the generation of nitric oxide by the reaction of hydrogen peroxide and D- or L-arginine. *Biochem. Biophys. Res. Commun.* **233**, 150-153.
- Navarro, A., Torrejon, R., Bandez, M. J., Lopez-Cepero, J. M. and Boveris, A. (2005). Mitochondrial function and mitochondria-induced apoptosis in an overstimulated rat ovarian cycle. *Am. J. Physiol.* **289**, 1101-1109.
- Ohta, F., Takagi, T., Sato, H. and Ignarro, L. J. (2007). Low-dose L-arginine administration increases microperfusion of hindlimb muscle without affecting blood pressure in rats. *Proc. Natl. Acad. Sci. USA* **104**, 1407-1411.
- Paglia, D. E. and Valentine, W. N. (1967). Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidases. *J. Lab. Clin. Med.* **70**, 158-169.
- Pansarasa, O., D'Antona, G., Gualea, M. R., Marzani, B., Pellegrino, M. A. and Mazat�o, F. (2002). "Oxidative stress": effects of mild endurance training and testosterone treatment on rat gastrocnemius muscle. *Eur. J. Appl. Physiol.* **87**, 550-555.
- Peralta, J. G., Finocchietto, P. V., Converso, D., Schopfer, F., Carreras, M. C. and Poderoso, J. J. (2003). Modulation of mitochondrial nitric oxide synthase and energy expenditure in rats during cold acclimation. *Am. J. Physiol.* **284**, 2375-2383.
- Petrović, V., Korać, A., Buzadžić, B. and Korać, B. (2005). The effects of L-arginine and L-NAME supplementation on redox-regulation and thermogenesis in interscapular brown adipose tissue. *J. Exp. Biol.* **208**, 4263-4271.
- Petrović, V., Buzadžić, B., Korać, A., Vasiljević, A., Janković, A. and Korać, B. (2006). Free radical equilibrium in interscapular brown adipose tissue: relationship between metabolic profile and antioxidative defense. *Comp. Biochem. Physiol.* **142C**, 60-65.
- Puigserver, P., Wu, Z., Park, C. W., Graves, R., Wright, M. and Spiegelman, B. M. (1998). A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* **92**, 829-839.
- Punkt, K., Fritzsche, M., Stockmar, C., Hepp, P., Josten, C., Wellner, M., Schering, S. and Buchwalow, I. (2006). Nitric oxide synthase in human skeletal muscles related to defined fibre types. *Histochem. Cell Biol.* **125**, 567-573.
- Robbarts, C. K., Barnard, R. J., Scheck, S. H. and Balon, T. W. (1997). Exercise-stimulated glucose transport in skeletal muscle is nitric oxide dependent. *Am. J. Physiol.* **273**, 220-225.
- Rolfe, D. F. and Brand, M. D. (1996). Contribution of mitochondrial proton leak to skeletal muscle respiration and to standard metabolic rate. *Am. J. Physiol.* **271**, 1380-1389.
- Saha, S. K., Ohinata, H. and Kuroshima, A. (1996). Effects of acute and chronic inhibition of nitric oxide synthase on brown adipose tissue thermogenesis. *Jpn. J. Physiol.* **46**, 375-382.
- Selman, C., McLaren, J. S., Himanka, M. J. and Speakman, J. R. (2000). Effect of long-term cold exposure on antioxidant enzyme activities in a small mammal. *Free Radic. Biol. Med.* **28**, 1279-1285.
- Shiota, M. and Masumi, S. (1988). Effect of norepinephrine on consumption of oxygen in perfused skeletal muscle from cold-exposed rats. *Am. J. Physiol.* **254**, 482-489.
- Skulachev, V. P. and Maslov, S. P. (1960). Role of non-phosphorylating oxidation in thermoregulation. *Biokhimiia* **25**, 1058-1064.
- Takada, Y., Noguchi, T., Okabe, T. and Kajiyama, M. (1982). Superoxide dismutase in various tissues from rabbits bearing the Vx-2 carcinoma in the maxillary sinus. *Cancer Res.* **42**, 4233-4235.
- Tiitus, P. M., Pushkarenko, J. and Houston, M. E. (1996). Lack of antioxidant adaptation to short-term aerobic training in human muscle. *Am. J. Physiol.* **271**, 832-836.
- Tonkonogi, M., Walsh, B., Svensson, M. and Sahlin, K. (2000). Mitochondrial function and antioxidative defence in human muscle: effects of endurance training and oxidative stress. *J. Physiol.* **528**, 379-388.
- Vallerand, A. L., Perusse, F. and Bukowiecki, L. J. (1990). Stimulatory effects of cold exposure and cold acclimation on glucose uptake in rat peripheral tissues. *Am. J. Physiol.* **259**, 1043-1049.
- Vasilaki, A., Mansouri, A., Remmen, H., van der Meulen, J. H., Larkin, L., Richardson, A. G., McArdle, A., Faulkner, J. A. and Jackson, M. J. (2006). Free radical generation by skeletal muscle of adult and old mice: effect of contractile activity. *Aging Cell* **5**, 109-117.
- Vasiljević, A., Buzadžić, B., Korać, A., Petrović, V., Janković, A., Mićunović, K. and Korać, B. (2007). The effects of cold acclimation and nitric oxide on antioxidative enzymes in rat pancreas. *Comp. Biochem. Physiol.* **145C**, 641-647.
- Wibom, R., Hultman, E., Johansson, M., Matheri, K., Constantin-Teodosiu, D. and Schantz, P. G. (1992). Adaptation of mitochondrial ATP production in human skeletal muscle to endurance training and detraining. *J. Appl. Physiol.* **73**, 2004-2010.
- Wickler, S. J. (1981). Nonshivering thermogenesis in skeletal muscle of seasonally acclimatized mice, *Peromyscus*. *Am. J. Physiol.* **241**, 185-189.
- Young, M. E., Radda, G. K. and Leighton, B. (1997). Nitric oxide stimulates glucose transport and metabolism in rat skeletal muscle *in vitro*. *Biochem. J.* **322**, 223-228.