

## Thermodynamics of oxygenation-linked proton and lactate binding govern the temperature sensitivity of O<sub>2</sub> binding in crustacean (*Carcinus maenas*) hemocyanin

Roy E. Weber\*, Jane W. Behrens†, Hans Malte and Angela Fago

Zoophysiology, Department of Biological Sciences, University of Aarhus, CF Møllers Alle 1131, DK 8000 Aarhus, Denmark

\*Author for correspondence (e-mail: roy.weber@biology.au.dk)

†Present address: Marine Biological Laboratory, University of Copenhagen, Strandpromenaden 5, 3000 Helsingør, Denmark

Accepted 16 January 2008

### SUMMARY

With the aim of understanding the molecular underpinnings of the enormous variation in the temperature sensitivity of hemocyanin–O<sub>2</sub> affinity encountered in crustaceans, we measured O<sub>2</sub> binding to *Carcinus maenas* hemocyanin at two temperatures, varying pH values and in the absence and presence of lactate ions in order to assess the contributions of oxygenation-linked binding of protons (the Bohr effect) and lactate ions to the overall enthalpies of oxygenation ( $\Delta H'$ ). The hemocyanin binds maximally 0.35 lactate ions per functional subunit. Lactate (which accumulates under hypoxic conditions) increases O<sub>2</sub> affinity by preferentially raising the association equilibrium constant of the hemocyanin in the low-affinity Tense state ( $K_T$ ), without significantly affecting that of the high-affinity Relaxed state ( $K_R$ ). In the absence of lactate, the variation in the temperature sensitivity observed with decreasing pH tallies neatly with changes in the nature and magnitude of the Bohr effect. Accordingly, the normal, absent and reverse Bohr effects observed under alkaline, neutral and acid conditions, respectively, reflect endothermic proton dissociation, absence of proton binding and exothermic proton association, respectively, upon oxygen binding. Oxygenation-linked lactate binding is exothermic, highly pH dependent and peaks near pH 7.6, where it contributes approximately  $-30 \text{ kJ mol}^{-1}$  to the overall heat of oxygenation. This predictably increases the temperature sensitivity of O<sub>2</sub> affinity, potentially hampering O<sub>2</sub> loading in warm, hypoxic habitats. The data demonstrate governing roles for lactate and proton ions in determining the temperature sensitivity of hemocyanin–O<sub>2</sub> affinity in crustaceans.

Key words: *Carcinus maenas*, crab, hemocyanin, lactate, oxygen binding, temperature effect.

### INTRODUCTION

The oxygenation of gas-transporting proteins is a function of their intrinsic O<sub>2</sub> affinities, as well as their interactions with allosteric effectors. The latter interactions govern O<sub>2</sub> affinity, and may modulate O<sub>2</sub> loading at the respiratory surfaces and/or O<sub>2</sub> unloading and in the metabolizing tissues in response to specific environmental conditions and metabolic requirements. The interactions of hemoglobins (Hbs) and hemocyanins (Hcs) with protons and CO<sub>2</sub> (the Bohr effect, which decreases O<sub>2</sub> affinity and facilitates O<sub>2</sub> in the respiring tissues) are a well-known example.

Unlike the intensively investigated vertebrate Hbs, where protons, chloride and organic phosphate anions are major effectors that decrease Hb–O<sub>2</sub> affinity, crustacean Hcs commonly exhibit marked sensitivities to L-lactate and urate anions and to divalent cations that increase Hc–O<sub>2</sub> affinity (Mangum and Towle, 1977; Truchot, 1980; Mason et al., 1983). Lactate is an end product of anaerobic metabolism that enhances O<sub>2</sub> loading, so it may be a link in the negative feedback that favours aerobiosis in the face of decreasing O<sub>2</sub> availability (Truchot, 1980). In contrast to the tetrameric, intraerythrocytic vertebrate Hbs that consist of two  $\alpha$  and two  $\beta$  chains and have a sedimentation constants of 4.4 S, crustacean Hcs commonly consist of 24 S dodecamers and/or 16 S hexamers which, respectively, comprise twelve and six 5 S polypeptide chains (Markl et al., 1979; Dainese et al., 1998; Molon et al., 2000; Podda et al., 2007). However, *Carcinus maenas* Hc consists of 24 S dodecamers only (Eriksson-Quensel and Svedberg, 1936; Markl et al., 1979).

A universal, although often neglected, factor governing O<sub>2</sub> affinity in ectothermic invertebrates is temperature. Based on the exothermic nature of their oxygenation reactions (Klotz and Klotz, 1955), the O<sub>2</sub> affinities of metal-containing, gas-binding proteins decrease with increasing temperature. Apart from the intrinsic heat of oxygenation ( $\Delta H^{O_2}$ ), the overall heat of oxygenation ( $\Delta H'$ ) includes the heat of solution of oxygen ( $\Delta H^{sol}$ ,  $-13 \text{ kJ mol}^{-1}$ ) and heats of processes linked to oxygenation, such as the release or binding of protons [including the heat of ionization of buffers ( $\Delta H^{H^+}$ )] and of other allosteric effectors ( $\Delta H^{X,Y,etc}$ ).

The  $\Delta H'$  values of crustacean Hc at physiological pH (7.6–7.8) vary tremendously (from  $-67$  to  $+134 \text{ kJ mol}^{-1}$ ) (cf. Jokumsen and Weber, 1982; Burnett et al., 1988; Brix et al., 1989; Adamczewska and Morris, 1998; Chausson et al., 2004). The available literature has variously attributed adaptive advantages to both low and high temperature sensitivities, arguing that the former may stabilize tissue O<sub>2</sub> supply in the face of environmental thermal variations, and the latter may increase O<sub>2</sub> unloading in the respiring tissues, in parallel with temperature-induced increases in metabolic O<sub>2</sub> requirement. Extensive studies of vertebrate Hbs attribute adaptive reductions in temperature sensitivity to endothermic processes coupled to O<sub>2</sub> binding, such as changes in protein conformation (Wyman et al., 1977) or, more commonly, dissociation of proton, phosphate and chloride ions (Wyman, 1964; Weber et al., 1985; Fago et al., 1997b; Weber et al., 2003). However, no systematic studies appear to have been carried out on the interactive effects of ligand binding on the

enthalpies of arthropod Hcs, notwithstanding the enormous variation in their temperature sensitivities and the radically different allosteric control mechanisms encountered compared to Hbs. In contrast to vertebrate Hbs, where protons and organic phosphates preferentially decrease O<sub>2</sub> affinity of the Hb molecules in the low-affinity Tense state ( $K_T$ ) without significantly affecting that of the Relaxed state ( $K_R$ ), in crab *Callinectes sapidus* Hc, L-lactate ions increase both  $K_T$  and  $K_R$ , whereas protons decrease both  $K_T$  and  $K_R$ , and Ca<sup>2+</sup> ions increase  $K_R$  (Johnson et al., 1988). Thus "...the allosteric interactions of L-lactate and crustacean hemocyanins ... provide an interesting contrast to the extremely well-studied allosteric interactions of the vertebrate hemoglobins" (Graham, 1985).

Aiming to probe the contributions of the major endogenous allosteric effectors (protons and lactate) to the overall enthalpies of oxygenation, we measured the interactive effects of lactate, temperature and pH on the O<sub>2</sub> affinity of Hc in the blood of the shore crab *Carcinus maenas*, which is exposed to large variations in ambient temperature on a daily and seasonal basis.

## MATERIALS AND METHODS

### Animals

Specimens of shore crabs *Carcinus maenas* L., with carapace width of 5–7 cm, were collected in Aarhus Bay, Denmark, and maintained in well-aerated seawater (30–32 p.p.t., 12°C,  $P_{O_2}$  >135 mmHg; 12 h:12 h photoperiod) for at least 4 days without feeding. Prebranchial hemolymph samples (1–2 ml) were withdrawn from the branchial sinuses by piercing the arthropodial membrane at the bases of the walking legs using a syringe fitted with a 22-G hypodermic needle. Samples from individual crabs were kept separately on ice. Those that did not coagulate were pooled into a Petri dish and whipped with a glass spatula to separate coagulating fibrinogen that was removed by filtration. Hemolymph was frozen at –80°C in 120 µl aliquots that were freshly thawed for oxygen binding studies.

### [Hc] and [L-lactate] measurements

Hc concentrations were estimated from peak absorbances near 335 nm, using an extinction coefficient of 17.5 mmol l<sup>-1</sup> cm<sup>-1</sup>, based on the  $A_{1\%1\text{cm}}$  value of 2.33 reported for *C. maenas* Hc (Nickerson and Van Holde, 1971) and a functional-unit mass of 75 kDa. Reagent grade L(+)-lactate, lithium salt (C<sub>3</sub>H<sub>5</sub>O<sub>3</sub>Li) was purchased from Sigma-Aldrich Chemicals (St Louis, MO, USA). L-lactate concentration in freshly collected hemolymph samples of crabs kept in normoxic seawater was measured according to Lowry and Passonneau's 'Method I' (Lowry and Passonneau, 2006).

### O<sub>2</sub> binding measurements of native hemolymph

Hemolymph samples were buffered at varying pH by adding 1 mol l<sup>-1</sup> Bis-Tris buffers to a final buffer concentration of 0.1 mol l<sup>-1</sup>. O<sub>2</sub> equilibria of 4 µl hemolymph samples were recorded at 10 and 20°C using a modified gas diffusion chamber connected to Wösthoff gas mixing pumps (Bochum, Germany) that mix air and ultrapure (>99.998%) N<sub>2</sub> to increase O<sub>2</sub> tensions stepwise (Weber, 1981; Weber et al., 1987), while absorbance is continuously monitored at 365 nm. pH values were measured in triplicate in separate 110 µl sub-samples (to avoid KCl contamination from the calomel electrode) at the same temperatures as the O<sub>2</sub> equilibrium measurements, using a BMS 2 Mk 2 microelectrode coupled to a PHM 64 Research pH meter (Radiometer, Copenhagen, Denmark). For each O<sub>2</sub> binding curve at least four equilibrium steps between 20% and 80% saturation were recorded, and  $P_{50}$  and  $n_{50}$  values (O<sub>2</sub>

tensions and Hill's cooperativity coefficients at 50% O<sub>2</sub> saturation, respectively) were interpolated from Hill plots,  $\log[S/(1-S)]$  vs  $\log P_{O_2}$ , where  $S$  is the fractional O<sub>2</sub> saturation.

The effects and stoichiometry of L-lactate binding were investigated by recording Hc–O<sub>2</sub> equilibria at varying L-lactate concentrations, and at pH values close to 6.9 and 7.7, and thereafter interpolating the  $P_{50}$  and  $n_{50}$  values at these exact pH values (6.90 and 7.70) from regression analysis of  $\log P_{50}$  vs pH.

For precise O<sub>2</sub> equilibrium measurements focusing on extreme, low and high O<sub>2</sub> saturations (extended Hill plots, see below), a subsample of the freshly prepared hemolymph was concentrated approximately twofold by centrifugation at 2500 g and 4°C for 1 h in Ultrafree-4 Millipore tubes with 10 000 Da molecular mass cut-off membranes.

The data were analysed in terms of the two-state MWC model, according to the equation:

$$S = [LK_T P(1+K_T P)^{(q-1)} + K_R P(1+K_R P)^{(q-1)}] / [L(1+K_T P)^q + (1+K_R P)^q], \quad (1)$$

where  $S$  denotes saturation,  $L$  is the equilibrium constant between the tense (T) and relaxed (R) state in fully deoxygenated form,  $K_T$  and  $K_R$  are the O<sub>2</sub> association equilibrium constants for the low-affinity (T, tense) and high-affinity (R, relaxed) forms, respectively,  $P$  is the partial pressure of O<sub>2</sub>, and  $q$  is the number of interacting binding sites (Monod et al., 1965). Curve-fitting to obtain  $K_T$ ,  $K_R$  and the allosteric constant  $L$ , estimation of the standard errors and calculation of the derived parameters  $P_{50}$ ,  $P_m$  (the median O<sub>2</sub> tension),  $n_{50}$ ,  $n_{\max}$  (the maximum cooperativity along the equilibrium curve) and  $\Delta G$  (the free energy of cooperativity) were carried out as detailed earlier (Weber et al., 1995; Fago et al., 1997a; Bugge and Weber, 1999).

The overall heat of oxygenation at varying pH values (range 6.1–8.6) and in the presence and absence of 10 mmol l<sup>-1</sup> L-lactate was evaluated from the difference in  $P_{50}$  values at 10 and 20°C, using the van't Hoff isochore (Wyman, 1964):

$$\Delta H' = 2.303R \cdot \Delta \log P_{50} / \Delta(1/T), \quad (2)$$

where  $R$  is the gas constant and  $T$  the absolute temperature. The heat of effector binding was then interpolated as the difference between  $\Delta H'$  values in the presence and absence of the effector (cf. Weber et al., 1985).

## RESULTS AND DISCUSSION

### Allosteric lactate interaction

Lactate concentration in hemolymph of crabs kept in aerated seawater was 0.034±0.023 mmol l<sup>-1</sup> ( $N=5$ ), confirming the low level previously reported for normoxic, resting *C. maenas* [0.1 mmol l<sup>-1</sup> (Lallier and Truchot, 1989)].

Dose–response curves measured at two pH values (Fig. 1) show that lactate increases O<sub>2</sub> affinity, exerting the greatest effect in the 50–100 mmol l<sup>-1</sup> concentration range, without markedly affecting the Bohr factor (depicted by the vertical distance between the two curves). Interpolated from the slopes of  $\log P_{50}$  vs  $\log[\text{lactate}]$  plots in this range, *C. maenas* Hc binds maximally 0.30 and 0.35 lactate ions per O<sub>2</sub> molecule at pH 7.0 and 7.7, respectively. Plotting  $\log P_{50}$  values against free lactate concentrations (estimated assuming that each functional Hc subunit binds 0.3 lactate ions at half-saturation) does not tangibly alter these coefficients, which compare with values of 0.11–0.54 observed in decapod crustacean Hc under different experimental conditions [(cf. Zeis et al., 1992; Adamczewska and Morris, 1998) and studies cited therein].




Fig. 1. (A) Effects of increasing lactate concentrations on hemolymph  $P_{50}$  values at pH 7.00 and 7.70 and 20°C. Hc subunit concentration, 0.85 mmol l<sup>-1</sup>.

Extended Hill plots of precise O<sub>2</sub> equilibria of the hemolymph at pH 7.84 (Fig. 2) show slopes of unity in the lower and upper extremities of the curves, reflecting non-cooperative binding of the first and last O<sub>2</sub> molecules to the functional units. Fitting the MWC model to the data in the absence and in the presence of 10 and 50 mmol l<sup>-1</sup> lactate, yields  $q$ -values (the number of interacting O<sub>2</sub>-binding sites) of 4.5, 4.7 and 3.2, respectively (Table 1). Fitting the model with  $q$  fixed at 4 (the average integral value found when the model was fitted with  $q$  floating) shows  $P_{50}$  to be 7.7 mmHg, O<sub>2</sub>




Fig. 2. Extended Hill plots of *C. maenas* Hc measured at 10°C in the absence (circles) and presence of 10 mmol l<sup>-1</sup> (triangles) and 50 mmol l<sup>-1</sup> (squares) L-lactate. Hc subunit concentration, 2.76 mmol l<sup>-1</sup>. Open and closed symbols are data points where the MWC equation was fitted with  $q$  floating and fixed at 4, respectively, during the iteration (see Eqn 1). Mean pH=7.843±0.009.

binding to be strongly cooperative ( $n_{50}$  and  $n_{max}=3.2$ ), and O<sub>2</sub> association constants for the Hc in the deoxy (Tense) and the oxygenated (Relaxed) states ( $K_T$  and  $K_R$ ) to be 0.01 and 1.61 mmHg<sup>-1</sup>, respectively (Table 1). The identity between  $n_{50}$  and  $n_{max}$  values and the close agreement between  $P_{50}$  and  $P_m$  values (Table 1) indicate symmetry of the O<sub>2</sub> binding curves, justifying rigorous analysis of the allosteric effects in terms of  $P_{50}$  values (cf. Wyman, 1964).

Calculated as  $\Delta G = RT \ln \frac{\langle(L+1)(Lc^q+1)\rangle}{\langle(Lc+1)(L^q+1)\rangle}$ , where  $L$  is the allosteric constant and  $c=K_T/K_R$  (Imai, 1982), the free energy of cooperativity equals 11.84 kJ mol<sup>-1</sup> in the absence of lactate and  $q=4$  (Table 1). As shown (Fig. 2, Table 1) lactate increases O<sub>2</sub> affinity (decreases  $P_{50}$ ) by lowering  $L$  and raising  $K_T$ , without markedly altering  $K_R$ . This indicates that the underlying molecular mechanism of the lactate effect is to shift the allosteric equilibrium towards the high affinity R state and to increase the O<sub>2</sub> affinity of the T state molecules, indicating that the T-state Hc becomes less stable in the presence of added lactate. The decreased  $K_R/K_T$  ratio decreases  $n_{50}$  values (slopes of the plots at  $\log[Y/(1-Y)]=0$ ) from 3.2 in the absence of lactate to 2.4 and 2.2 in the presence of 10 and 50 mmol l<sup>-1</sup> lactate, respectively, and correspondingly lowers  $\Delta G$  from 11.84 to 7.90 and 6.77, respectively.

#### Interactive effects of lactate and pH

Measurement of the O<sub>2</sub> affinity of *C. maenas* Hc at 10 and 20°C, in the absence and presence of lactate over a wide pH range where the Bohr effect (oxygenation-linked proton binding) changes from normal to absent and reverse (Fig. 3) provides an ideal opportunity to analyse the contributions of individual, oxygenation-linked processes to the overall enthalpy of oxygenation.

#### pH

As shown (Fig. 3) the Hc exhibits a strong normal Bohr effect  $\phi$  ( $=\Delta \log P_{50}/\Delta \text{pH}$ )=-0.80 at alkaline pH range, which encompasses physiological values in decapods [7.44-7.84 (Mangum and Shick, 1972)], no Bohr effect ( $\phi=0$ ) at pH 6.9, and a distinct, reversed Bohr effect ( $\phi=+0.46$ ) at lower pH. The data show a drastic increase in the temperature effect with decreasing pH (Fig. 4), indicating that high temperatures will adversely affect O<sub>2</sub> loading under acidotic and hypoxic conditions.

Fig. 5 shows the relationship between  $\Delta H'$  and pH in the absence and presence (Fig. 5 top, curves A and B, respectively) of lactate. In the absence of lactate (Fig. 5 top, curve A) the numerical value of  $\Delta H'$  increases from ~-10 kJ mol<sup>-1</sup> at pH 8 to plateaus of approx. -42 at pH 7.0 and ~60 at pH 6.5, revealing a clear inverse relationship with the Bohr factors (-0.8, 0 and +0.46) at these respective pH values. This indicates that (i) as with vertebrate Hb, oxygenation-linked proton dissociation is endothermic and thus counterbalances the heat released upon Hc oxygenation ( $\Delta H^{O_2}$ ) in the pH range where the Bohr effect is normal, (ii) the overall  $\Delta H'$  value at pH 7.0 is devoid of contributions from proton binding ( $\Delta H^{H^+}$ ) and thus reflects the intrinsic heat of O<sub>2</sub> binding by the Hc, (iii) oxygenation-linked proton association (reverse Bohr effect) is exothermic and thus increases the temperature effect contributing about +18 (= -60 to -42) kJ mol<sup>-1</sup> and (iv) in the absence of other effectors, proton concentrations play the dominating role in determining the temperature sensitivity of O<sub>2</sub> affinity in crustacean Hc.

This contention is supported by numerous data in the literature. Particular examples are Hcs of (i) crab *Holthuisana transversa*, where an exceptionally low Bohr factor (-0.13) correlates with

Table 1. MWC and derived parameters for *Carcinus maenas* hemolymph with 0, 10 and 50 mmol l<sup>-1</sup> L-lactate at 10°C

[L-lactate] (mmol l <sup>-1</sup> )	pH	P <sub>50</sub> (mmHg)	P <sub>m</sub> (mmHg)	n <sub>50</sub>	n <sub>max</sub>	logK <sub>T</sub>	logSE	logK <sub>R</sub>	logSE	logL	q*	ΔG (kJ mol <sup>-1</sup> )
A												
0	7.83	7.81	7.83	3.4	3.4	-1.9651	0.05169	0.1483	0.0879	4.6990	4.51	11.39
10	7.85	4.66	4.69	2.6	2.6	-1.3854	0.0839	0.0128	0.1010	3.1980	4.67	7.51
50	7.85	3.00	3.11	2.0	2.0	-1.4589	0.2894	0.0903	0.0455	1.8727	3.22	7.45
B												
0	7.83	7.69	7.70	3.2	3.2	-1.9861	0.0505	0.2055	0.0775	4.3670	4	11.84
10	7.85	4.63	4.67	2.4	2.4	-1.4364	0.0700	0.0495	0.0817	2.8739	4	7.90
50	7.85	3.00	3.10	2.2	2.2	-1.2314	0.0604	0.0708	0.0347	2.2671	4	6.77

\*The MWC model was fitted to the data with  $q$  (the number of interacting O<sub>2</sub>-binding sites) floating (A) or fixed at 4 (B). SE, standard errors (of K<sub>T</sub> and K<sub>R</sub> values); for other abbreviations, see text.

highly exothermic oxygenation reactions ( $\Delta H' = -48$  to  $-62$  kJ mol<sup>-1</sup>) observed over a range of operating conditions (Morris et al., 1988), (ii) hermit crab *Pagurus bernhardus*, where a massive Bohr effect [ $\varphi = -1.55$  (Jokumsen and Weber, 1982)] tallies with a complete lack of temperature sensitivity at pH 7.2–7.8, whereas the temperature effect is large at low pH where the Bohr effect is zero (A. Jokumsen and R.E.W., unpublished), and (iii) krill *Meganctiphanes norvegica*, and the hydrothermal vent crab *Segonzacia mesatlantica*, where even larger Bohr factors ( $\varphi = -1.9$  and  $-2.9$ , respectively) are associated with overall endothermic heats of oxygenation [ $\Delta H' = +134$  and  $+17$  kJ mol<sup>-1</sup> (Brix et al., 1989; Chausson et al., 2004)].

The differences between  $\Delta H'$  in the presence and absence of lactate (depicted by curve B–A, Fig. 5 bottom) reflects the pH-dependent heats of reaction with lactate ions and of coupled reactions. As evident, lactate drastically raises the  $\Delta H'$  values at alkaline (physiological) pH. Given that lactate increases O<sub>2</sub> affinity (Fig. 1) and thus undergoes oxygenation-linked binding (and

deoxygenation-linked dissociation), the data show that lactate binding is exothermic, strongly pH-dependent, and peaks at a pH of approximately 7.6, where it contributes approximately  $-30$  kJ mol<sup>-1</sup> per O<sub>2</sub> molecule bound/released and decreases at higher and lower pH values. The exothermic, oxygenation-linked binding of lactate ions that increases the overall oxygenation enthalpy in crab Hc contrasts sharply with vertebrate Hbs, where endothermic, oxygenation-linked dissociation of organic effectors (2,3-diphosphoglycerate, DPG, and ATP) reduces the overall heat of oxygenation and thus the temperature sensitivity of O<sub>2</sub> affinity.

The single previously reported value for the enthalpic contribution of lactate binding to Hc [ $-25$  kJ mol<sup>-1</sup> at pH 7.5 for Hc of the crab *Calappa granulata* (Olianas et al., 2006)] is in good agreement with our data. Remarkably, when measured directly by calorimetric methods, the enthalpy change for urate binding to *Homarus vulgaris* Hc is much higher [ $135$  kJ mol<sup>-1</sup> at pH 8.0 (Menze et al., 2001)]. The enthalpies of lactate binding to crab Hc are also considerably lower than for DPG binding to human Hb, where similar values ( $\sim 46$ – $55$  kJ mol<sup>-1</sup>) were obtained from van't Hoff plots and calorimetric methods (Benesch et al., 1969; Bunn et al., 1971; Nelson et al., 1974). Literature on mammalian Hbs illustrate how the tyranny of the van't Hoff equation may be bypassed by endothermic processes (e.g. conformational changes and proton and chloride binding) (Wyman et al., 1977; Ikeda-Saito et al., 1983; De Rosa et al., 2004) that reduce the temperature sensitivity of O<sub>2</sub> affinity. Arguably, a low  $\Delta H'$  value is an advantageous trait for O<sub>2</sub> loading at high temperature when an effector (such as lactate) increases O<sub>2</sub> affinity and thereby raises the exothermic nature of oxygenation,




Fig. 3. P<sub>50</sub> and n<sub>50</sub> values of *C. maenas* Hc measured at 10°C (circles) and 20°C (triangles) in the absence of added lactate (open symbols) and presence (closed symbols) of 10 mmol l<sup>-1</sup> L-lactate. Hc subunit concentration, 1.11 mmol l<sup>-1</sup>.




Fig. 4. O<sub>2</sub> equilibrium curves of *C. maenas* Hc at 10 and 20°C and pH 7.8 and 7.0, interpolated from data in Fig. 3.






Fig. 5. (Top) Overall oxygenation enthalpies ( $\Delta H'$ ) derived using the van't Hoff isochore and  $P_{50}$  values at 10 and 20°C and different pH values (Fig. 3), in the absence of added lactate (A) and the presence of 10 mmol l<sup>-1</sup> lactate (B). (Bottom) The interpolated, pH-dependent difference between these values (B-A).

just as low  $\Delta H$  values resulting from ligand reactions coupled to oxygen binding are advantageous for maintaining a high O<sub>2</sub> affinity in vertebrate Hbs.

Although lactate that is produced under anaerobic conditions increases Hc-O<sub>2</sub> affinity and thus may favour O<sub>2</sub> loading and survival under hypoxic conditions (Truchot, 1980), our data show that a thermodynamic consequence of oxygenation-linked lactate binding is an increased temperature sensitivity in the physiological pH range (cf. Fig. 5), which may compromise O<sub>2</sub> loading at high temperature. While this specific 'lactate effect' ( $\Delta H^{\text{lact}}$ ) is maximal near 7.5, the overall temperature sensitivity ( $\Delta H'$ ) remains high at lower pH values (imparted by lactate accumulation), where the Bohr effect successively disappears and reverses (resulting in loss of O<sub>2</sub>-linked endothermic proton dissociation and initiation of ectothermic proton binding, respectively). However, apart from hampering O<sub>2</sub> binding at high temperature, increased temperature sensitivity will promote O<sub>2</sub> loading in the gills at low temperature. In this regard it may be significant that increases in lactate levels may induce behavioural hypothermia, since lactate injection reduces the preferred temperature in *C. maenas* (De Wachter et al., 1997).

In conclusion, the large variations in the effects of temperature on O<sub>2</sub> affinity of crustacean Hcs (observed between different species and within the same individuals at different times) appear to be directly related to variations in the heats of reaction of processes coupled to the oxygenation reaction, notably proton and lactate binding, indicating that hemolymph levels of these ions are major factors controlling the temperature sensitivity of oxygen binding to crustacean hemocyanin.

In this light it would be interesting to investigate the enthalpic components in Hcs that lack a lactate effect [as in the hydrothermal vent crab *Syanagraea praedator* (Chausson et al., 2001)] or have

an opposite lactate effect [as in the land crab *Gecarcoidea natalis*, where lactate decreases O<sub>2</sub> affinity (Adamczewska and Morris, 1998)] as well as the thermodynamic consequences of binding other Hc effectors, such as divalent cations, whose concentrations may vary greatly (with water salinity, ionoregulatory capacities and the moult cycle) and urate, which may bind to Hc with a higher affinity than L-lactate [40-fold higher in *Homarus vulgaris* Hc (Nies et al., 1992)]. In *Carcinus maenas*, whose Hc has about twice as many urate (and caffeine) binding sites than that of *H. vulgaris* (Hellmann et al., 2004), hemolymph urate concentrations increase drastically in response to hypoxic exposure (Lallier et al., 1987). In line with the effect of lactate at pH > 7.2 reported here (Fig. 5), urate increases the temperature sensitivity of slipper lobster *Scyllarides latus* Hc, which has four urate binding sites per hexamer, at pH 7.5 (Sanna et al., 2004).

Supported by the Danish Natural Science Research Council (FNU) and the Carlsberg Foundation. We thank Anny Bang and Dorte Olsson for valuable technical assistance.

## REFERENCES

- Adamczewska, A. M. and Morris, S. (1998). The functioning of the haemocyanin of the terrestrial Christmas Island red crab *Gecarcoidea natalis* and roles for organic modulators. *J. Exp. Biol.* **201**, 3233-3244.
- Benesch, R. E., Benesch, R. and Yu, C. I. (1969). The oxygenation of hemoglobin in the presence of 2,3-diphosphoglycerate. Effect of temperature, pH, ionic strength, and hemoglobin concentration. *Biochemistry* **8**, 2567-2571.
- Brix, O., Borgund, S., Barnung, T., Colosimo, A. and Giardina, B. (1989). Endothermic oxygenation of hemocyanin in the krill *Meganycitaphanes norvegica*. *FEBS Lett.* **247**, 177-180.
- Bugge, J. and Weber, R. E. (1999). Oxygen binding and its allosteric control in hemoglobin of the pulmonate snail, *Biomphalaria glabrata*. *Am. J. Physiol.* **276**, R347-R356.
- Bunn, H. F., Ransil, B. J. and Chao, A. (1971). The interaction between erythrocyte organic phosphates, magnesium ion, and hemoglobin. *J. Biol. Chem.* **246**, 5273-5279.
- Burnett, L. E., Scholnick, D. A. and Mangum, C. P. (1988). Temperature sensitivity of molluscan and arthropod hemocyanins. *Biol. Bull.* **174**, 153-162.
- Chausson, F., Bridges, C. R., Sarradin, P. M., Green, B. N., Riso, R., Caprais, J. C. and Lallier, F. H. (2001). Structural and functional properties of hemocyanin from *Cyanograea praedator*, a deep-sea hydrothermal vent crab. *Proteins* **45**, 351-359.
- Chausson, F., Sanglier, S., Leize, E., Hagege, A., Bridges, C. R., Sarradin, P. M., Shillito, B., Lallier, F. H. and Zal, F. (2004). Respiratory adaptations to the deep-sea hydrothermal vent environment: the case of *Segonzacia mesatlantica*, a crab from the Mid-Atlantic Ridge. *Micron* **35**, 31-41.
- Dainese, E., Di Muro, P., Beltramini, M., Salvato, B. and Decker, H. (1998). Subunits composition and allosteric control in *Carcinus aestuarii* hemocyanin. *Eur. J. Biochem.* **256**, 350-358.
- De Rosa, M. C., Castagnola, M., Bertonati, C., Galtieri, A. and Giardina, B. (2004). From the Arctic to fetal life: physiological importance and structural basis of an 'additional' chloride-binding site in haemoglobin. *Biochem. J.* **380**, 889-896.
- De Wachter, B., Sartoris, F. J. and Pörtner, H. O. (1997). The anaerobic endproduct lactate has a behavioural and metabolic signalling function in the shore crab *Carcinus maenas*. *J. Exp. Biol.* **200**, 1015-1024.
- Eriksson-Quensel, I.-B. and Svedberg, T. (1936). The molecular weights and pH-stability regions of the hemocyanins. *Biol. Bull.* **71**, 498-547.
- Fago, A., Bendixen, E., Malte, H. and Weber, R. E. (1997a). The anodic hemoglobin of *Anguilla anguilla*. Molecular basis for allosteric effects in a Root-effect hemoglobin. *J. Biol. Chem.* **272**, 15628-15635.
- Fago, A., Wells, R. M. G. and Weber, R. E. (1997b). Temperature-dependent enthalpy of oxygenation in Antarctic fish hemoglobins. *Comp. Biochem. Physiol.* **118B**, 319-326.
- Graham, R. A. (1985). A model for L-lactate binding to *Cancer magister* hemocyanin. *Comp. Biochem. Physiol.* **81B**, 885-887.
- Hellmann, N., Hornemann, J., Jaenicke, E. and Decker, H. (2004). Urate as effector for crustacean hemocyanins. *Micron* **35**, 109-110.
- Ikeda-Saito, M., Yonetani, T. and Gibson, Q. H. (1983). Oxygen equilibrium studies on hemoglobin from the bluefin tuna (*Thunnus thynnus*). *J. Mol. Biol.* **168**, 673-686.
- Imai, K. (1982). *Allosteric Effects in Haemoglobin*. Cambridge: Cambridge University Press.
- Johnson, B. A., Bonaventura, C. and Bonaventura, J. (1988). Allosterism in *Callinectes sapidus* hemocyanin: cooperative oxygen binding and interactions with L-lactate, calcium, and protons. *Biochemistry* **27**, 1995-2001.
- Jokumsen, A. and Weber, R. E. (1982). Hemocyanin-oxygen affinity in hermit crab blood is temperature independent. *J. Exp. Zool.* **221**, 389-394.
- Klotz, I. M. and Klotz, T. A. (1955). Oxygen-carrying proteins: a comparison of the oxygenation reaction in hemocyanin and hemerythrin with that in hemoglobin. *Science* **121**, 477-480.
- Lallier, F. and Truchot, J. P. (1989). Hemolymph oxygen transport during environmental hypoxia in the shore crab, *Carcinus maenas*. *Respir. Physiol.* **77**, 323-336.

- Lallier, F., Boitel, F. and Truchot, J. P.** (1987). The effect of ambient oxygen and temperature on haemolymph L-lactate and urate concentrations in the shore crab *Carcinus maenas*. *Comp. Biochem. Physiol.* **86A**, 255-260.
- Lowry, O. H. and Passonneau, J. V.** (2006). *A Flexible System of Enzymatic Analysis*. Orlando: Academic Press.
- Mangum, C. P. and Shick, J. M.** (1972). The pH of body fluids of marine invertebrates. *Comp. Biochem. Physiol.* **42A**, 693-697.
- Mangum, C. and Towle, D.** (1977). Physiological adaptation to unstable environments. Inconstancy of the internal milieu in an animal may be a regulatory mechanism. *Am. Sci.* **65**, 67-75.
- Markl, J., Hofer, A., Bauer, G., Markl, A., Kempfer, B., Brenzinger, M. and Linzen, B.** (1979). Subunit heterogeneity in arthropod hemocyanins. II. Crustacea. *J. Comp. Physiol.* **133**, 167-175.
- Mason, R. P., Mangum, C. P. and Godette, G.** (1983). The influence of inorganic ions and acclimation salinity on hemocyanin-oxygen binding in the blue crab *Callinectes sapidus*. *Biol. Bull.* **164**, 104-123.
- Menze, M. A., Hellmann, N., Decker, H. and Grieshaber, M. K.** (2001). Binding of urate and caffeine to haemocyanin analysed by isothermal titration calorimetry. *J. Exp. Biol.* **204**, 1033-1038.
- Molon, A., Di Muro, P., Bubacco, L., Vasilyev, V., Salvato, B., Beltrami, M., Conze, W., Hellmann, N. and Decker, H.** (2000). Molecular heterogeneity of the hemocyanin isolated from the king crab *Paralithodes camtschaticae*. *Eur. J. Biochem.* **267**, 7046-7057.
- Monod, J., Wyman, J. and Changeux, J.-P.** (1965). On the nature of allosteric transitions: a plausible model. *J. Mol. Biol.* **12**, 88-118.
- Morris, S., Greenaway, P. and McMahon, B. R.** (1988). Oxygen and carbon dioxide transport by the haemocyanin of an amphibious crab, *Holthuisana transversa*. *J. Comp. Physiol. B* **157**, 873-882.
- Nelson, D. P., Miller, W. D. and Kiesow, L. A.** (1974). Calorimetric studies of hemoglobin function, the binding of 2,3-diphosphoglycerate and inositol hexaphosphate to human hemoglobin A. *J. Biol. Chem.* **249**, 4770-4775.
- Nickerson, K. W. and Van Holde, K. E.** (1971). A comparison of molluscan and arthropod hemocyanin-I. Circular dichroism and absorption spectra. *Comp. Biochem. Physiol.* **39B**, 855-872.
- Nies, A., Zeis, B., Bridges, C. R. and Grieshaber, M. K.** (1992). Allosteric modulation of haemocyanin oxygen-affinity by L-lactate and urate in the lobster *Homarus vulgaris*. II. Characterization of specific effector binding sites. *J. Exp. Biol.* **168**, 111-124.
- Olianas, A., Sanna, M. T., Messana, I., Castagnola, M., Masia, D., Manconi, B., Cau, A., Giardina, B. and Pellegrini, M.** (2006). The hemocyanin of the shamed crab *Calappa granulata*: structural-functional characterization. *J. Biochem.* **139**, 957-966.
- Podda, G., Manconi, B., Olianas, A., Pellegrini, M., Messana, I., Mura, M., Castagnola, M., Giardina, B. and Sanna, M. T.** (2007). Structural and functional characterization of hemocyanin from the anemone hermit crab *Dardanus calidus*. *J. Biochem.* doi:10.1093/jb/mvm210.
- Sanna, M. T., Olianas, A., Castagnola, M., Sollai, L., Manconi, B., Salvadori, S., Giardina, B. and Pellegrini, M.** (2004). Oxygen-binding modulation of hemocyanin from the slipper lobster *Scyllarides latus*. *Comp. Biochem. Physiol.* **139B**, 261-268.
- Truchot, J.-P.** (1980). Lactate increases the oxygen affinity of crab hemocyanin. *J. Exp. Zool.* **214**, 205-208.
- Weber, R. E.** (1981). Cationic control of O<sub>2</sub> affinity in lugworm erythrocrurin. *Nature* **292**, 386-387.
- Weber, R. E., Wells, R. M. G. and Rossetti, J. E.** (1985). Adaptations to neoteny in the salamander, *Necturus maculosus*. Blood respiratory properties and interactive effects of pH, temperature and ATP on hemoglobin oxygenation. *Comp. Biochem. Physiol.* **80A**, 495-501.
- Weber, R. E., Jensen, F. B. and Cox, R. P.** (1987). Analysis of teleost hemoglobin by Adair and Monod-Wyman-Changeux models. Effects of nucleoside triphosphates and pH on oxygenation of tench hemoglobin. *J. Comp. Physiol.* **157B**, 145-152.
- Weber, R. E., Malte, H., Braswell, E. H., Oliver, R. W. A., Green, B. N., Sharma, P. K., Kuchumov, A. and Vinogradov, S. N.** (1995). Mass spectrometric composition, molecular mass and oxygen binding of *Macrobdella decora* hemoglobin and its tetramer and monomer subunits. *J. Mol. Biol.* **251**, 703-720.
- Weber, R. E., Hourdez, S., Knowles, F. and Lallier, F.** (2003). Hemoglobin function in deep-sea and hydrothermal-vent endemic fish: *Symenichelis parasitica* (Anguillidae) and *Thermarces cerberus* (Zoarcidae). *J. Exp. Biol.* **206**, 2693.
- Wyman, J., Jr** (1964). Linked functions and reciprocal effects in hemoglobin: a second look. *Adv. Protein Chem.* **19**, 223-286.
- Wyman, J., Gill, S. J., Noll, L., Giardina, B., Colosimo, A. and Brunori, M.** (1977). The balance sheet of a hemoglobin. Thermodynamics of CO binding by hemoglobin Trout I. *J. Mol. Biol.* **109**, 195-205.
- Zeis, B., Nies, A., Bridges, C. R. and Grieshaber, M. K.** (1992). Allosteric modulation of haemocyanin oxygen-affinity by L-lactate and urate in the lobster *Homarus vulgaris*. I. Specific and additive effects on haemocyanin oxygen-affinity. *J. Exp. Biol.* **168**, 93-110.