

RESEARCH ARTICLE

Cardiac responses to hypercapnia in larval zebrafish (*Danio rerio*): the links between CO₂ chemoreception, catecholamines and carbonic anhydrase

Scott Miller¹, Jacob Pollack¹, Julia Bradshaw², Yusuke Kumai³ and Steve F. Perry^{1,*}

ABSTRACT

The ontogeny of carbon dioxide (CO₂) sensing in zebrafish (*Danio rerio*) has not been examined. In this study, CO₂-mediated increases in heart rate were used to gauge the capacity of zebrafish larvae to sense CO₂. CO₂ is thought to be detected via neuroepithelial cells (NECs), which are homologous to mammalian carotid body glomus cells. Larvae at 5 days post-fertilization (d.p.f.) exhibited tachycardia when exposed for 30 min to 0.75% CO₂ (~5.63 mmHg); at 7 d.p.f., tachycardia was elicited by 0.5% CO₂ (~3.75 mmHg). Based on pharmacological evidence using β -adrenergic receptor (β -AR) antagonists, and confirmed by β_1 -AR translational gene knockdown using morpholinos, the reflex tachycardia accompanying hypercapnia was probably mediated by the interaction of catecholamines with cardiac β_1 receptors. Because the cardiac response to hypercapnia was abolished by the ganglionic blocker hexamethonium, it is probable that the reflex cardio-acceleration was mediated by catecholamines derived from sympathetic adrenergic neurons. Owing to its likely role in facilitating intracellular acidification during exposure to hypercapnia, it was hypothesized that carbonic anhydrase (CA) is involved in CO₂ sensing, and that inhibition of CA activity would blunt the downstream responses. Indeed, the cardiac response to hypercapnia (0.75% CO₂) was reduced in fish at 5 d.p.f. exposed to acetazolamide, a CA inhibitor, and in fish experiencing zCac (CA2-like a) knockdown. Successful knockdown of zCac was confirmed by CA activity measurements, western blotting and immunocytochemistry. Co-injection of embryos with zCac morpholino and mRNA modified at the morpholino binding site restored normal levels of CA activity and protein levels, and restored (rescued) the usual cardiac responses to hypercapnia. These data, combined with the finding that zCac is expressed in NECs located on the skin, suggest that the afferent limb of the CO₂-induced cardiac reflex in zebrafish larvae is initiated by coetaneous CO₂-sensing neuroepithelial cells.

KEY WORDS: Hypercapnia, Chemoreception, Neuroepithelial cell, Tachycardia cardiorespiratory control

INTRODUCTION

Adult fish including zebrafish (*Danio rerio*) exhibit well-defined cardiorespiratory responses to elevated environmental CO₂ (hypercapnia). Typically, most species that have been examined

exhibit hyperventilation that is mediated by increases in breathing frequency and/or ventilatory stroke volume (Dejours, 1973; Janssen and Randall, 1975; Randall et al., 1976; Smatresk and Cameron, 1982; Perry and Gilmour, 1996; Crocker et al., 2000; Vulesevic et al., 2006; Perry et al., 2009b) (for reviews, see Shelton et al., 1986; Perry and Wood, 1989; Milsom, 1995; Gilmour, 2001; Perry and Gilmour, 2002; Gilmour and Perry, 2007; Perry et al., 2009a; Perry and Abdallah, 2012). The cardiovascular responses to hypercapnia have received less attention but in those few species that have been examined, a conserved response appears to be bradycardia (Perry et al., 1999; Sundin et al., 2000; Reid et al., 2000; McKendry et al., 2001; Perry and McKendry, 2001; Gilmour et al., 2005) which may, or may not, be associated with an increase in arterial blood pressure (reviewed by Gilmour and Perry, 2007). Bradycardia, however, is not a universal response to hypercapnia; two species were shown to exhibit tachycardia [*Acipenser transmontanus* (Crocker et al., 2000) and *Tinca tinca* (Randall and Shelton, 1963)] and in others, cardiac frequency is unaltered (Gilmour and Perry, 2007).

Because cardiorespiratory adjustments in fish are principally keyed to O₂ status (Dejours, 1973; Randall and Jones, 1973; Smith and Jones, 1982) (reviewed by Perry and Wood, 1989), the hyperventilatory responses to hypercapnia were originally thought to reflect the indirect effects of respiratory acidosis on lowering blood O₂ content (hypoxaemia) via Bohr and Root shifts (Smith and Jones, 1982). The striking inverse correlation between blood oxygenation status and ventilation in rainbow trout (*Oncorhynchus mykiss*) led to the novel, yet still unproven concept of a blood O₂ content receptor (Randall, 1982). Although hypoxaemia may contribute indirectly to the reflex responses to hypercapnia, it is now well established that fish possess specific peripheral CO₂ chemoreceptors that are directly involved in the initiation of downstream responses (Kinkead and Perry, 1991; Perry and McKendry, 2001; McKendry and Perry, 2001; McKendry et al., 2001; Gilmour et al., 2005) and which function independently of O₂ (Heisler et al., 1988; Perry and Gilmour, 1996). In adult fish, the CO₂ chemoreceptors are localized to the gills (McKendry et al., 2001; Perry and Reid, 2002) or buccal cavity (Reid et al., 2000) where they typically respond to external (but not internal) CO₂ (McKendry and Perry, 2001; Perry and McKendry, 2001) and exhibit little, if any, reactivity to external [H⁺] (Reid et al., 2000; Perry and McKendry, 2001; Gilmour et al., 2005). Despite the weight of evidence supporting the existence of externally oriented CO₂ receptors (reviewed by Gilmour and Perry, 2007), several studies have indeed provided evidence supporting a role for internally oriented CO₂/H⁺ receptors (Wood et al., 1990; Wood and Munger, 1994).

Regardless of their modality (CO₂ versus H⁺) or orientation (external versus internal), there is compelling *in vitro* evidence that neuroepithelial cells (NECs) (Dunel-Erb et al., 1982) of the adult gill

¹Department of Biology, University of Ottawa, 30 Marie Curie, Ottawa, ON, Canada, K1N 6N5. ²Department of Fisheries and Oceans, Pacific Biological Station, 3190 Hammond Bay Road, Nanaimo, BC, Canada, V9T 6N7. ³Department of Physiology and Biophysics, Case Western Reserve University School of Medicine, 10900 Euclid Avenue, Cleveland, OH 44106, USA.

*Author for correspondence (sfperry@uottawa.ca)

filament are dual O₂/CO₂ chemoreceptors (Jonz et al., 2004; Bursleson et al., 2006; Olson et al., 2008; Qin et al., 2010; Abdallah et al., 2012; Abdallah et al., 2014) and thus functionally analogous to the type I (glomus) cells of the mammalian carotid body (Milsom and Bursleson, 2007). Similar to O₂ sensing (Jonz et al., 2004), CO₂ chemoreception by zebrafish NECs is associated with membrane depolarization following inhibition of background potassium conductance (Qin et al., 2010). Ultimately, the membrane depolarization initiates an elevation of intracellular [Ca²⁺] owing to mobilization of intracellular Ca²⁺ stores (Abdallah et al., 2014), which is believed to trigger neurotransmitter release that promotes the downstream reflex responses. Although there is no direct *in vivo* evidence implicating the NECs in reflex cardiorespiratory responses to changing ambient O₂ levels, alterations in NEC morphology or abundance during hypoxia (Jonz et al., 2004) or hyperoxia (Vulesevic et al., 2006) are consistent with their presumed role in O₂ sensing. In contrast, the density of NECs on the adult gill of zebrafish is unaltered by chronic elevation of ambient CO₂ (Vulesevic et al., 2006).

In mammals, CO₂ chemoreception by type 1 cells is facilitated by the enzyme carbonic anhydrase (CA) because it promotes rapid intracellular acidification via the catalysed hydration of molecular CO₂ (Lahiri and Forster, 2003). Neuroepithelial cells isolated from adult zebrafish gill exhibit CA immunoreactivity and inhibition of CA with acetazolamide, while slowing the rate of intracellular acidification (Abdallah et al., 2014) delayed and blunted the membrane depolarization in isolated NECs exposed to hypercapnia (Qin et al., 2010). Interestingly, however, the CO₂-mediated rise in intracellular [Ca²⁺] was unaffected by CA inhibition (Abdallah et al., 2014). Thus given the conflicting *in vitro* data and in the absence of *in vivo* data, it is difficult to assign a role to CA in the facilitation of the cardiorespiratory reflex responses to hypercapnia.

Little is known about the ontogeny of O₂ sensing in fish and there are no data concerning CO₂ chemoreception in developing larvae. With respect to O₂ sensing, it is thought that the NECs of the skin are critical in promoting the ventilatory responses to hypoxia prior to development and innervation of gill NECs (Jonz and Nurse, 2005; Coccimiglio and Jonz, 2012). Given that gill NECs (or more likely a subset of gill NECs) are dual O₂/CO₂ chemoreceptors, it seems likely that the responses to CO₂ in larval zebrafish (if present) also are initiated by the coetaneous NECs. Thus the initial objective of the present study was simply to determine whether zebrafish larvae display cardiorespiratory responses to elevated external CO₂. Initial experiments demonstrated a robust and reproducible elevation in cardiac frequency to hypercapnia, while the ventilatory responses were more variable. Thus throughout this study, a change in heart rate was used as the physiological determinant of CO₂ sensing. After establishing the neurohumoral mechanisms underlying hypercapnic tachycardia, the principal goal of this study was to test the hypothesis that CA is involved in CO₂ sensing and ultimately in promoting the ensuing downstream physiological reflex responses. The sensing of CO₂ and the pathways associated with it were assessed using standard pharmacological methods, coupled with translational gene knockdown.

RESULTS

At 4 d.p.f., none of the concentrations of CO₂ that were tested affected heart rate (f_H ; Fig. 1A). Note, however, that the resting f_H in the fish at 4 d.p.f. was somewhat higher than in the other groups and thus it is possible that the fish at this age lacked the scope to increase f_H . At 5 d.p.f., only the highest level of CO₂ (0.75%) produced a significant increase in f_H when compared with the normocapnic




Fig. 1. The effect of 30 min of hypercapnia (CO₂ ~zero (air), 0.25, 0.5 or 0.75%) on heart rate (f_H) in zebrafish (*Danio rerio*) larvae. Effect at (A) 4 d.p.f., (B) 5 d.p.f. and (C) 7 d.p.f.; values are expressed as means \pm s.e.m.; $N=6$ different fish at each developmental stage. Data points not sharing common letters are statistically different from each other ($P<0.05$; one-way repeated measures ANOVA).

group ($P=0.002$; Fig. 1B). At 7 d.p.f., f_H was increased, at 0.5% and 0.75% CO₂ ($P<0.001$; Fig. 1C). Because 5 d.p.f. was the earliest stage of development when fish exhibited CO₂ sensitivity, this stage was chosen for all subsequent experiments, except for those involving CA rescue (performed at 4 d.p.f.) inhibition of CA by acetazolamide (ACTZ) for which experiments were conducted at 5 and 7 d.p.f. The more robust f_H response to CO₂ at 7 d.p.f. made it easier to detect potential inhibitory effects following CA inhibition.

At 0.75% CO₂ (the highest concentration used in the initial phase of this study), the pH of the water decreased from 7.2 to 6.6. Exposing zebrafish to normocapnic water at pH 6.6 (Fig. 2) caused a significant decrease in f_H ($P=0.037$), thereby demonstrating that the tachycardia observed during exposure to hypercapnic acidosis was probably the result of CO₂, and not the associated reduction in pH.

The addition of hexamethonium, a nicotinic receptor antagonist and ganglionic blocker within the parasympathetic and sympathetic divisions of the autonomic nervous system, prevented a significant increase in f_H in fish exposed to 0.75% CO₂ (Fig. 3). The heart rate of the control fish exposed to 0.75% CO₂ was significantly higher




Fig. 2. The effect of isocapnic acidosis (pH was lowered from 7.2 to 6.6 which corresponded to the reduction in pH observed when fish were exposed to 0.75% CO_2) on heart rate (f_H) in zebrafish (*Danio rerio*) larvae at 5 d.p.f. Values are expressed as means \pm s.e.m.; $N=6$. *Statistically significant difference ($P<0.05$) from the data obtained at pH 7.2 (Student's paired t -test).

than either the air-exposed (normocapnic) controls or the CO_2 -exposed fish treated with hexamethonium. Therefore, it can be concluded that the downstream responses to CO_2 at 5 d.p.f. are under neural control.

Addition of propranolol, a non-specific β -adrenergic receptor blocker (Fig. 4A), caused a significant decrease in heart rate in the normocapnic fish (from to 137.3 ± 2.2 to 123.3 ± 4.4 beats min^{-1}) and prevented the increase in f_H that was observed in the hypercapnic control fish (Fig. 4). Unlike propranolol, the specific β_1 -adrenergic receptor antagonist, atenolol, did not affect f_H in normocapnic fish; however, atenolol did prevent the usual increase in f_H accompanying 0.75% CO_2 (Fig. 4B).

The studies using pharmacological blockade of β -adrenergic receptors were complemented by additional experiments employing translational gene knockdown of the β_1 -AR (Fig. 5). Unlike the sham-injected group that significantly increased their f_H when exposed to 0.75% CO_2 ($P=0.003$), the group of fish experiencing β_1 -AR knockdown did not increase its f_H during hypercapnia (157.3 ± 6.8 beats min^{-1} in normocapnia versus 151.7 ± 3.9 beats min^{-1} ; $N=6$). At 0.75% CO_2 , the fish experiencing β_1 -AR knockdown had a significantly lower f_H than the sham-injected fish ($P<0.001$).

To determine whether CA was involved in CO_2 sensing, cardiac responses to hypercapnia were assessed with and without ACTZ, a membrane-permeable CA inhibitor. At 5 d.p.f., the ACTZ-treated fish failed to increase f_H at all levels of CO_2 (data for exposure to




Fig. 3. The effect of the ganglionic blocker hexamethonium (Hex; 10^{-4} mol l^{-1}) on the cardiac frequency (f_H) response of zebrafish (*Danio rerio*) larvae at 5 d.p.f. to hypercapnia (30 min of 0.75% CO_2). Values are expressed as means \pm s.e.m.; $N=6$. Letters indicate differences among the group ($P<0.05$; one-way repeated measures ANOVA).




Fig. 4. The effects of β -adrenergic receptor (β -AR) blockade. Effects of (A) the non-specific β -AR blocker propranolol (10^{-4} mol l^{-1}) or (B) the β_1 -AR antagonist atenolol (10^{-4} mol l^{-1}) on the cardiac frequency (f_H) response to hypercapnia (30 min of 0.75% CO_2) in zebrafish (*Danio rerio*) larvae at 5 d.p.f. Values are expressed as means \pm s.e.m. *Significant difference ($P<0.05$) between normocapnia (air) and hypercapnia (0.75% CO_2); [‡]significant difference ($P<0.05$) between control (unfilled bars, $N=6$) and propranolol-treated fish (filled bars, $N=6$); two-way repeated measures ANOVA.

0.75% CO_2 are shown in Fig. 6A). When 7 d.p.f. zebrafish (which exhibited a more robust response to CO_2 in comparison to fish at 5 d.p.f.) were tested, ACTZ treatment again prevented a significant increase in f_H (Fig. 6B).

To ensure that the ACTZ-treated fish were still capable of increasing f_H , fish at 5 and 7 d.p.f. were exposed to the cardiac stimulant noradrenaline (Fig. 7A,B). In both sets of ACTZ-treated fish, the addition of noradrenaline caused a significant increase in




Fig. 5. The effects of β_1 -AR knockdown on the cardiac frequency (f_H) response to hypercapnia (30 min of 0.75% CO_2) in zebrafish (*Danio rerio*) larvae at 5 d.p.f. Values are expressed as means \pm s.e.m. *Significant difference ($P<0.05$) between normocapnia (air) and hypercapnia (0.75% CO_2); [‡]significant difference ($P<0.05$) between sham (unfilled bars, $N=6$) and fish experiencing β_1 -AR knockdown (filled bars, $N=6$); two-way repeated measures ANOVA.




Fig. 6. The effects of the carbonic anhydrase inhibitor acetazolamide (ACTZ) on the cardiac frequency (f_H) response to hypercapnia (30 min of 0.75% CO₂) in zebrafish (*Danio rerio*) larvae. Effects at (A) 5 d.p.f. and (B) 7 d.p.f.; values are expressed as means \pm s.e.m. *Significant difference ($P < 0.05$) between normocapnia (air) and hypercapnia (0.75% CO₂); †significant difference ($P < 0.05$) between sham (unfilled bars, $N=6$) and ACTZ-treated fish (filled bars, $N=6$); two-way repeated measures ANOVA.

f_H . This showed that ACTZ was not preventing the heart from responding normally to β -adrenergic receptor stimulation.

Because acetazolamide is likely to inhibit all isoforms of zebrafish CA, an alternate approach employing selective gene knockdown was used to specifically assess the role of the zCAc isoform. The data summarized in Fig. 8 clearly demonstrate that knockdown of zCAc prevented the increase in f_H that was observed in the fish injected with the control morpholino. The effectiveness of the CA knockdown and its rescue using zCAc mRNA (slightly modified to prevent binding of the morpholino) are depicted in Figs 9 and 10. Preliminary experiments revealed that although CA knockdown was effective until 5 d.p.f. (CA activity was reduced from 425.9 to 141.2 pmol min⁻¹ μ g⁻¹), rescue of CA activity could not be reliably sustained longer than 4 d.p.f. At 4 d.p.f., CA activity in morphants was reduced by 84% (from 305.5 to 49.6 pmol min⁻¹ μ g⁻¹; Fig. 9A) and protein levels were undetectable by western blotting (Fig. 9B). Additionally, CA localized by immunocytochemistry (ICC) to cells of the yolk sac in sham fish was not detectable in the zCAc morphants (Fig. 10A–D). Co-injection of the zCAc mRNA with morpholino resulted in normal levels of CA activity and protein levels (Fig. 9A,B). The elevation in heart rate during hypercapnia exposure, normally absent in the larvae experiencing zCAc knockdown, was restored after zCAc rescue (Fig. 9C). However, the rescue was only partial because the increase in heart rate, although significant, was less pronounced than in the sham fish (Fig. 9C).

The NECs on the skin were readily identified by immunocytochemical detection of 5-HT (Fig. 10C). A subset of the 5-HT-containing NECs also expressed zCAc (Fig. 10D,E). The majority of cells expressing zCAc, however, did not co-express 5-




Fig. 7. The effects of noradrenaline (NA; 10^{-4} mol l⁻¹) on cardiac frequency (f_H) in zebrafish (*Danio rerio*) larvae exposed to acetazolamide (ACTZ). Effects at (A) 5 d.p.f. ($N=6$) and (B) 7 d.p.f. ($N=6$). Values are expressed as means \pm s.e.m. *Significant effect ($P < 0.05$; Student's paired t -test) of noradrenaline (filled bars) on f_H .

HT; these cells were most likely to be H⁺-ATPase-enriched ionocytes (data not shown).

DISCUSSION

The purpose of this study was to examine CO₂ sensing in developing zebrafish and to establish the mechanisms of the ensuing downstream effects on cardiac function. The major findings emerging from this research are that (i) zebrafish larvae respond to hypercapnia with tachycardia that is probably mediated by a neuronal reflex involving adrenergic activation of cardiac β -adrenergic receptors and (ii) the CO₂-mediated tachycardia relies on the activity of CA presumably owing to its role in CO₂ chemoreception.




Fig. 8. The effect of zCAc knockdown on the cardiac frequency (f_H) response to hypercapnia (30 min of 0.75% CO₂) in zebrafish (*Danio rerio*) larvae at 5 d.p.f. Values are expressed as means \pm s.e.m.; *significant difference ($P < 0.05$) between normocapnia (air) and hypercapnia (0.75% CO₂); †significant difference ($P < 0.05$) between sham (unfilled bars, $N=6$) and fish experiencing zCAc knockdown (filled bars, $N=6$); two-way repeated measures ANOVA.




Fig. 9. zCac rescue restores CA activity and re-enables the hypercapnic tachycardia during zCac knockdown. The effects of zCac knockdown (filled bars; $N=4$) and rescue with zCac mRNA (hatched bars; $N=3$) on (A) whole body CA activity in zebrafish (*Danio rerio*) larvae at 4 d.p.f. Values are expressed as means \pm s.e.m.; *significant difference ($P<0.05$) from the sham-injected (unfilled bars; $N=5$) larvae (one-way ANOVA). (B) zCac protein levels as depicted in a representative western blot. Note that the standard conditions used in all physiological trials involving knockdown and/or rescue were 4 ng morpholino (MO) per embryo and 200 pg CAc mRNA per embryo. (C) Cardiac frequency (f_H) response to hypercapnia (30 min of 1.0% CO_2) in larvae at 4 d.p.f. Values are expressed as means \pm s.e.m.; *significant difference ($P<0.05$) between normocapnia (unfilled bars) and hypercapnia (filled bars; 1.0% CO_2); ‡significant difference ($P<0.05$) between sham ($N=6$) and fish experiencing zCac knockdown ($N=6$) or zCac knockdown/rescue ($N=6$); two-way repeated measures ANOVA.

Recently, it was observed that NECs of the skin are innervated after 1 d.p.f. (Coccimiglio and Jonz, 2012) and appear to function in O_2 sensing well before gill NECs become innervated at ~ 7 d.p.f. (Jonz and Nurse, 2005). Because at least a portion of the NECs also act as CO_2 chemoreceptors (Qin et al., 2010), an underlying assumption of the present study was that any effects of CO_2 on cardiac function would presumably be initiated by the NECs either on the skin (at 4 and 5 d.p.f.) and/or gill (at 7 d.p.f.). Despite the likelihood that NECs function as CO_2 sensors *in vivo*, strictly speaking and similar to O_2 sensing, there are as yet no direct *in vivo* data implicating NECs in CO_2 chemoreception.

The cardiac response to hypercapnia

The basic response to elevated CO_2 at 5–7 d.p.f. was an elevation in heart rate. An increase in heart rate in zebrafish larvae during

hypercapnia differs from the bradycardia that is usually observed in adult fish (note that no data are yet available for adult zebrafish). Similar to the bradycardia in adult fish (e.g. Perry et al., 1999), the physiological significance of the hypercapnia-mediated tachycardia in larvae is unknown. Given that the results of a recent study (Gilmour et al., 2009) suggested that the red blood cells of larval zebrafish may be involved in CO_2 excretion, it is conceivable that internal convection might selectively promote CO_2 excretion and thereby be increased by any elevation of cardiac output associated with tachycardia. Thus the tachycardia could serve to minimize the extent of the respiratory acidosis associated with exposure to hypercapnia.

Zebrafish at 5 d.p.f. increased their heart rate when exposed to elevated P_{CO_2} (0.75% CO_2 or 5.6 mmHg); the response was more robust at 7 d.p.f. with heart rate being increased at CO_2 levels of 0.5% or 3.75 mmHg (Fig. 1). The CO_2 levels required to elicit cardiac responses in larvae far exceeded those needed to promote hyperventilation in adults (1 mmHg) (Vulesevic et al., 2006). Without directly comparing the response characteristics of larval and adult zebrafish NECs to CO_2 , it is not possible to determine whether these differences in responsiveness reflect age-dependent differences in NEC sensitivity thresholds, lack of maturation of the efferent limb of the CO_2 -mediated reflex in larvae (see below) or simply intrinsic variability in the CO_2 thresholds required to elicit cardiac versus ventilation responses.

Given that gross motor responses to hypoxia were observed in zebrafish as early as 2 d.p.f. (Jonz and Nurse, 2005) and ventilatory responses to hypoxia were apparent at 3 d.p.f. (Coccimiglio and Jonz, 2012), it is tempting to speculate that the skin NECs may be less responsive to changes in ambient CO_2 (cardiac responses observed only at 4 d.p.f.) in comparison with O_2 . However, for similar reasons as discussed above, these differences may represent varying rates of maturation of the efferent branches of the cardiac and ventilation control systems as well as the severe levels of hypoxia used to elicit responses in larvae [e.g. 25 mmHg P_{O_2} in larvae (Coccimiglio and Jonz, 2012) compared with 100 mmHg in adults (Vulesevic et al., 2006)]. The delayed onset of sensitivity to CO_2 (relative to O_2) may reflect the relatively benign effects of hypercapnia on zebrafish development compared with the marked negative effects on growth in fish exposed to hypoxia (Pelster, 2002; Vulesevic and Perry, 2006).

Typically adult fish exhibit similar cardiorespiratory responses to hypoxia and hypercapnia, the conserved responses being hyperventilation and bradycardia (see Introduction). There are insufficient data to draw similar conclusions for larvae and those few studies that have been performed on hypoxic zebrafish have yielded widely conflicting results. For example, although hypoxic bradycardia was observed in 4 d.p.f. larvae in at least two studies (Bagatto, 2005; Steele et al., 2009), there are also reports of the zebrafish heart being unresponsive to hypoxia until 30 d.p.f. at which time bradycardia develops (Barrionuevo and Burggren, 1999), or actually exhibiting tachycardia (Jacob et al., 2002). The underlying basis for this variability in the cardiac responses to hypoxia in larval zebrafish is unclear but may reflect, at least in part, the different levels of hypoxia used [e.g. Jacob et al. (Jacob et al., 2002) exposed fish to a much milder hypoxia (75 mmHg)] or the different regimes of exposure [e.g. Barrionuevo and Burggren (Barrionuevo and Burggren, 1999) exposed fish acutely to hypoxia whereas in the other studies, embryos/larvae were reared in hypoxic water]. Thus until additional experiments are performed that directly compare hypoxic and hypercapnic responses under similar conditions, it is not yet possible to conclude that the tachycardia observed in the present study in response to hypercapnia truly differs from the larval hypoxic response.




Fig. 10. Representative immunocytochemistry micrographs. These depict (A) zCac-expressing cells (ionocytes) on the yolk sac epithelium and (B) their absence in larvae at 4 d.p.f. after zCac knockdown. Panels C–E illustrate a subset of 5-HT-positive NECs expressing zCac (arrows). Panel A is reproduced from fig. 2 in the review paper of Kwong et al. (Kwong et al., 2014), with permission.

CO₂ versus pH

As CO₂ concentrations in water increase, the external pH is decreased, a condition known as acidic hypercapnia. To assess the effects of decreasing water pH in the absence of elevated CO₂, the pH of the water was decreased from 7.2 to 6.6 (representing the change in pH accompanying the exposure to 0.75% CO₂). This isocapnic acidosis resulted in a drop in heart rate, showing that the increase in heart rate caused by acidic hypercapnia was related specifically to CO₂ (Fig. 2). Similar results were obtained in previous studies using adult fish (Burlison and Smatresk, 2000; McKendry et al., 2001; McKendry and Perry, 2001; Perry and McKendry, 2001; Perry and Reid, 2002; Gilmour et al., 2005; Abdallah et al., 2014). Thus the bulk of available evidence now suggests (although see Introduction) that the cardiorespiratory reflexes associated with acidic hypercapnia reflect the increase in P_{CO_2} rather than the decrease in pH (Gilmour and Perry, 2007). Given the apparent opposite effects of external isocapnic acidosis and acidic hypercapnia, it is conceivable that exposure to isohydric hypercapnia (increase in CO₂ with no change in pH) would lead to a greater increase in heart rate than with acidic hypercapnia, but that was not measured in this study.

Hypercapnic tachycardia is an adrenergic reflex

In the present study, the increase in heart rate associated with hypercapnia was blocked by the application of hexamethonium, showing that nicotinic acetylcholine receptors in the sympathetic and/or parasympathetic ganglia are involved. Therefore, it can be concluded that the tachycardia induced by CO₂ is a neural reflex and that the increase in heart rate is not caused by a local effect of CO₂ on the heart. Because hexamethonium acts on both the parasympathetic and sympathetic divisions of the autonomic nervous system, the CO₂-mediated increase in heart rate could potentially reflect decreased activity of the inhibitory parasympathetic pathways or increased activity of stimulatory adrenergic pathways, assuming that both of these pathways are tonically active (see below). The muscarinic receptor blocker, atropine, had no effect on resting heart rate or the cardiac response to CO₂ (results not shown). Therefore, it would appear that the parasympathetic pathway is not involved. Previous studies have also

reported a lack of an effect of atropine on resting heart rate until about 12 d.p.f. (Schwerte et al., 2006; Steele et al., 2009), suggesting that parasympathetic cholinergic tone is absent until ~12 d.p.f. However, Hsieh and Liao (Hsieh and Liao, 2002) reported a significant reduction in heart rate in 3 d.p.f. larvae experiencing muscarinic (M₂) receptor knockdown and Steele et al. (Steele et al., 2009) demonstrated that hypoxic bradycardia at 3 d.p.f. was prevented by M₂ receptor knockdown. The latter two findings provide evidence for parasympathetic tone in larvae of the age used in the present study but nevertheless it would appear to play no role in regulating heart rate during hypercapnia. The present study, like others that have assessed cardiac function in zebrafish larvae, utilized animals lightly anaesthetized with MS-222. Thus it is conceivable that the anaesthesia may have somehow interfered with the normal parasympathetic control of the heart.

Blockade of β -adrenergic receptors using the non-specific β -AR antagonist propranolol, or the β_1 -AR antagonist atenolol, prevented the increase in heart rate during hypercapnia. To resolve potential problems associated with non-specific effects of pharmacological blockade of β -receptors, a gene knockdown approach was employed to specifically target the β_1 receptor (Steele et al., 2011). In contrast to the results of Steele et al. (Steele et al., 2011), β_1 -AR knockdown did not lower resting heart rate. However, similar to β -AR receptor pharmacological blockade, the tachycardia induced by hypercapnia was prevented by β_1 -AR knockdown. Therefore, the increase in heart rate during hypercapnia is certainly related to adrenergic activation of β_1 -adrenergic receptors.

Although the present results and those of previous studies provide evidence for adrenergic tone in larvae at 4–7 d.p.f. (Bagatto, 2005; Schwerte et al., 2006; Wang et al., 2009; Steele et al., 2011), it is not straightforward to distinguish between tone arising from a functional sympathetic innervation of the heart and tone associated with circulating catecholamines. Control of catecholamine secretion, although well characterized in adults (Reid et al., 1998; Perry and Bernier, 1999; Perry and Capaldo, 2011), has not been studied in larval fish. Thus it remains unknown whether the larvae investigated in the present study had yet developed an acute humoral adrenergic stress response capable of raising circulating catecholamine levels to stimulate cardiac β_1 -ARs. Although it has been suggested

(Schwerte et al., 2006) that the zebrafish heart is not yet innervated at 4–7 d.p.f., the prevention of the cardiac response to hypercapnia by hexamethonium in the current study (Fig. 3) does support a role for sympathetic cardiac innervation. Alternatively, it is conceivable that hexamethonium exerted its effects by interacting with nicotinic receptors (if innervated in larvae at 5 d.p.f.) on the catecholamine-secreting chromaffin cells (Montpetit and Perry, 1999). Thus the results of the present study can be interpreted in at least two ways: activation of cardiac sympathetic nerve fibres or activation of pre-ganglionic sympathetic nerve fibres innervating chromaffin tissue. In both cases, elevated catecholamines produced locally via sympathetic nerves or arriving via the circulatory system would activate cardiac adrenergic receptors. We propose that regardless of the precise mechanism(s) underlying cardiac β -AR activation, the response is initiated by the CO_2 -sensing NECs of the skin or gill (in older larvae). Future experiments presumably incorporating new techniques will be required to firmly establish a direct link between the NECs and downstream cardiorespiratory responses.

A role for carbonic anhydrase

Qin et al. (Qin et al., 2010) demonstrated that CA was present in NECs isolated from adult zebrafish gills where it played an important role in setting the magnitude and speed of membrane depolarization during exposure of these cells to hypercapnia. In the current study, CA was localized to the NECs of the skin of larvae, and its role in CO_2 transduction was assessed using pharmacological inhibition (ACTZ) and selective gene knockdowns. When ACTZ was added to the water, the magnitude of the tachycardia response to elevated CO_2 was reduced. In carotid body type I cells, inhibition of CA delayed the onset of, and reduced the magnitude of chemosensory responses to CO_2 (Iturriaga, 1993). Similarly, CA inhibition reduced the discharge rate from laryngeal receptors (Coates et al., 1996) and blunted central chemosensory responses to elevated CO_2 in the cat (Coates et al., 1991).

Acetazolamide inhibits all isoforms of CA including the red cell-specific isoform zCAB and the more general cytosolic form zCAC. It is the zCAC paralog that is believed to be present in the zebrafish NEC, a notion that was confirmed in the present study by using a zCAC-specific antibody. Given the previous results of Qin et al. (Qin et al., 2010) demonstrating the involvement of CA in promoting NEC membrane depolarization with elevated CO_2 , and the generally held view that CA in the type I cells of the carotid body plays a role in CO_2 sensing (Iturriaga et al., 1991), the most parsimonious explanation for the effects of ACTZ observed in the current study is that they reflect inhibition of NEC CA activity. To ensure that fish were still able to respond to adrenergic stimulation with an increase in cardiac frequency (especially in light of the increased basal frequency in ACTZ-treated larvae at 5 d.p.f.), noradrenaline was added to the bathing water; at 5 and 7 d.p.f., zebrafish were still able to increase their heart rates significantly in response to noradrenaline. Thus ACTZ in itself does not appear to interfere with the efferent arm of the CO_2 -mediated cardiac reflex.

To more selectively inhibit specific CA isoforms, a gene knockdown strategy was employed to specifically lower zCAC activity. The results of a previous study (Gilmour et al., 2009) demonstrated the specificity of the zCAC knockdown and most importantly showed that zCAB was not a target for the zCAC morpholino. Thus we are confident that the similar knockdown approach used in this study also specifically targeted zCAC. While a decrease in whole body CA activity was anticipated after zCAC knockdown (Fig. 9), the large magnitude of the response was unexpected given that zCAB activity should have remained. The

most likely explanation for this finding is that red blood cell CA activity accounts for only a small fraction of total CA activity in larvae at 4 d.p.f.

Although CA activity was previously implicated in the speed and magnitude of membrane depolarization in isolated zebrafish NECs exposed to CO_2 (Qin et al., 2010) and the results of the current study clearly identify CA as a key component of the CO_2 transduction pathway *in vivo*, Abdallah et al. (Abdallah et al., 2014) demonstrated that the CO_2 -mediated rise in NEC intracellular $[\text{Ca}^{2+}]$ was unaffected by CA inhibition although it slowed the rate of intracellular acidification. The uncoupling of intracellular acidification and intracellular Ca^{2+} levels ultimately was attributed to the exclusive dependency of Ca^{2+} mobilization in zebrafish NECs on extracellular rather than intracellular acidification (Abdallah et al., 2014), an observation which is consistent with the presence of extracellular pH-sensing TASK channels in zebrafish (Pena-Munzenmayer et al., 2014). Clearly, further research should be directed at elucidating how CA inhibition impedes NEC membrane depolarization (Qin et al., 2010) and blunts CO_2 -mediated cardiac reflexes (present study) without influencing intracellular Ca^{2+} levels.

MATERIALS AND METHODS

Animals

Adult zebrafish, *Danio rerio*, were purchased from a commercial supplier (Big Al's, Ottawa, Canada) and maintained at the University of Ottawa Aquatic Care Facility. They were kept in 10-litre tanks that were supplied with well aerated, dechloraminated tap water at 28°C. The fish were kept at a constant photoperiod of 14 h:10 h light:dark and fed daily until satiation with Zeigler No. 1 crumble™ (Aquatic Habitats, Apopka, FL, USA). Embryos were collected and reared in 50 ml Petri dishes with E3 medium (5 mmol l⁻¹ NaCl, 0.17 mmol l⁻¹ KCl, 0.33 mmol l⁻¹ CaCl₂, 0.33 mmol l⁻¹ MgSO₄ and 0.1% Methylene Blue). The Petri dishes were kept in incubators set at 28.5°C (in the dark) until they were removed for experimentation (4–7 d.p.f.). Dead embryos were removed and E3 medium was changed daily.

To obtain larvae, 1-litre breeder traps were placed on the bottom of the tanks the night before breeding, and fish were allowed to spawn for 3 h. To obtain embryos for micro-injections (see below), 2-litre breeding traps were set up with two females and one male that were allowed to breed for 30 min the following morning. The experiments and handling of the animals were carried out in accordance with institutional guidelines (University of Ottawa Animal Care and Veterinary Service protocol BL-226) that conform to the guidelines of the Canadian Council on Animal Care (CCAC).

Heart rate measurements

Larvae were anaesthetized in a small volume of 80 mg l⁻¹ of Tris-buffered MS-222 (ethyl 3-aminobenzoate methanesulphonate salt, Sigma-Aldrich Inc., St Louis, MO, USA). After 5 min, larvae ($N=6$ for all trials unless noted otherwise) were transferred to a flow through system for heart rate observations. Individual larvae at 4–7 d.p.f. were placed in a rectangular depression well (14 mm) inside a superfusion chamber made from a glass-bottomed Petri dish (MatTek, Ashland, MA, USA) and Sylgard (Dow Corning, Midland, MI, USA). Two holes were carved out of the Sylgard on opposite sides of the depression well to enable inflow and outflow of water. These holes were covered with nylon mesh to retain the fish in the well. Using a gravity-fed system, dechloraminated water containing 0.05 mg ml⁻¹ MS-222 was continuously provided to the chamber at 4 ml min⁻¹ through gas-impermeable tubing (Tygon, Saint-Gobain Performance Plastics, Pittsburgh, PA, USA). Solutions were removed from the chamber using a peristaltic pump (ThermoFisher Scientific, Waltham, MA, USA). To visualize heart rate, the chamber was placed under a stereomicroscope fitted with a CCD video camera with output to a computer.

Series 1 – the effect of CO_2 on heart rate

Baseline measurements were taken on fish for 1 min; CO_2 levels were increased from air-saturated water to 0.75% CO_2 in 0.25% increments with

heart rate measurements taken after 30 min for each level of CO₂. CO₂ levels in the perfusate reservoirs were controlled using a Cameron 2- or 3-channel gas mixer, which provided mixed gas to the reservoirs at a rate of 2000 ml min⁻¹.

To distinguish between the effects of CO₂ and H⁺ on cardiac function during hypercapnia, fish were exposed to isocapnic acidosis where the pH of the water was lowered to 6.6 (matching the acidosis associated with 0.75% CO₂) using HCl.

A separate group of fish was exposed to the CA inhibitor acetazolamide (10⁻⁴ mol l⁻¹; Sigma-Aldrich), dissolved in water for 20 min prior to measurements and for the course of the treatment. To dissolve the acetazolamide, the pH of the water was increased to ~11.0 using NaOH and then titrated back down to 7.2 using HCl (Gilmour et al., 2009).

Series 2 – the effects of ganglionic and β-adrenergic receptor blockade on the cardiac responses to hypercapnia

Fish were anaesthetized as described above and placed in the flow-through system for baseline measurements. However, instead of first increasing the CO₂, one of the following drugs (prepared in water; 10⁻⁴ mol l⁻¹) was added to the air-equilibrated water: hexamethonium (nicotinic receptor antagonist and ganglionic blocker), propranolol (non-specific β-receptor antagonist), or atenolol (β₁-adrenergic receptor antagonist). Another measurement was taken after 30 min of exposure to one of the three drugs. The CO₂ was then increased to 0.75%, and a final measurement was taken.

Series 3 – translational gene knockdown using morpholinos

Morpholino injection

Morpholinos (tagged with 3'-carboxyfluorescein) for zebrafish cytosolic CA (zCAc; NM_199215, also referred to as CA2-like a) (Gilmour et al., 2009), the β₁-adrenergic receptor (β₁-AR; AB294717) (Steele et al., 2011) and a control morpholino were purchased from Gene Tools (Philomath, OR, USA) with the following sequences, respectively: 5'-AGTGGTCAGCCAT-TCCGCCAGCTGT-3', 5'-ACGGTAGCCCGTCTCCATTG-3' and 5'-CCTCTTACCTCAGTTACAATTTATA-3'. All morpholinos were prepared to a final concentration of 4 ng nl⁻¹ in Danieau buffer (58 mmol l⁻¹ NaCl, 0.7 mmol l⁻¹ KCl, 0.4 mmol l⁻¹ MgSO₄, 0.6 mmol l⁻¹ Ca(NO₃)₂, and 5 mmol l⁻¹ Hepes with a final pH of 7.6), and Phenol Red was used as an indicator of positive injection. Injections into one- or two-cell embryos were performed using a Narishige IM300 microinjector (Narishige International USA Inc., Long Island, NY, USA) with needles made from 1.0 mm borosilicate glass (Sutter Instrument, Novato, CA, USA). Dosage and sequence of zCA isoforms are based on Gilmour et al. (Gilmour et al., 2009) and β₁-AR from Steele et al. (Steele et al., 2011). Neither study reported adverse effects when using a dosage of 4 ng per embryo. After injections, embryos were placed in Petri dishes containing E3 medium and placed in an incubator at 28°C. On the following day, embryos were screened for positive expression of fluorescein using a Nikon SMZ 150 stereomicroscope (Nikon Instruments Inc., Melville, NY, USA).

Series 4 – carbonic anhydrase rescue

For rescue experiments, zCAc mRNA was synthesized *in vitro* with five neutral mismatches in the morpholino-binding 5' region and included in the injection solution at a concentration of 100 or 200 pg nl⁻¹, or 100–200 pg per embryo. To control for toxic effects of the morpholino, a separate group of embryos were injected with 4 ng per embryo of nonsense morpholino (standard control morpholino oligonucleotide; Gene Tools LLC). Because the effects of rescue on CA activity were declining by 5 d.p.f., the effects of hypercapnia on f_{H_1} were assessed in larvae at 4 d.p.f. using a higher level of CO₂ (1%; 7.5 mmHg).

Zebrafish cDNA was made using RevertAid primers (Fermentas Canada Inc., Burlington, ON, Canada), according to the manufacturer's instructions and primed with random hexamers. Primers used to make the mRNA for the zCAc rescue were: forward: 5'-ACGGCAGGGCATGGCTGACCACT-3'; reverse: 5'-TTAAAAGATGCACGCACCAC-3'. The product was inserted into a P-drive (Qiagen Inc., Toronto, ON, Canada) for sequence confirmation and sub-cloned into a pCS2+ vector. The mRNA was then synthesized using a mMessage-mMachine kit (Ambion, Streetsville, ON, Canada), according to the manufacturer's instructions. The product was purified by phenol:chloroform extraction following the transcription and

then run on a denaturing gel to confirm the size and integrity of the mRNA. mRNA (100 or 200 pg) was injected into the zebrafish embryos at the 1- or 2-cell stage, either alone or in conjunction with zCAc morpholino.

Western blotting for zCAc and CA activity measurements

To confirm knockdown and exogenous rescue of cytosolic zCAc, western blotting was carried out on larvae at 4 d.p.f. and CA activity measurements were performed at 3–5 d.p.f. For western blotting, larvae were collected, pooled into groups of 20 (*N*=1) and homogenized on ice with a protease inhibitor cocktail (Roche, USA) and radio-immunoprecipitation assay (RIPA) buffer (150 mmol l⁻¹ NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 50 mmol l⁻¹ Tris-HCl, 1 mmol l⁻¹ EDTA, 1 mmol l⁻¹ phenylmethanesulfonyl fluoride, 0.1% SDS). The samples were incubated for 30 min and centrifuged at 15,000 *g* for 20 min at 4°C. The protein-containing supernatant was stored at -20°C until use. Protein samples were size-separated using 10% SDS-PAGE and then transferred to a polyvinylidene difluoride (PVDF) membrane (Bio-Rad, USA). The membrane was blocked for 1 h in Tris-buffered saline with Tween 20 (TBST) (137 mmol l⁻¹ NaCl, 19 mmol l⁻¹ Tris base, 2.7 mmol l⁻¹ KCl, 0.01% Tween 20) with 5% milk, and then incubated overnight on a shaker at 4°C with a 1:3000 dilution of zCAc antibody in TBST containing 2% milk (peptide sequence: WGYDKHNGPDKWGC; Genscript, NJ, USA) in 2% milk (TBST). The membrane was washed three times for 15 min in TBST and subsequently incubated for 2 h with 1:15,000 goat anti-rabbit secondary antibody (Pierce, USA). Luminata Western HRP Substrates (Millipore, USA) was used to visualize immunoreactive bands. To verify equal protein loading, the membrane was stripped with Re-Blot Plus solution (Millipore, USA) and incubated with 1:4000 β-actin antibody (A2066, Sigma-Aldrich) at room temperature for 2 h. Band intensity was estimated using ImageJ (<http://imagej.nih.gov/ij/>) and compared with that of β-actin.

Carbonic anhydrase activity was measured using the electrometric delta pH method as described by Henry (Henry, 1991). The assay used a temperature-controlled pH electrode (Radiometer GK2401C combination electrode) attached to a pH meter (Radiometer PHM64 research pH meter) to measure the change in pH of a Tris-buffered media (10 mmol l⁻¹ Tris base, 225 mmol l⁻¹ mannitol, 75 mmol l⁻¹ sucrose; Sigma-Aldrich, pH adjusted to 7.4 with 30% phosphoric acid) in the presence of CO₂-saturated H₂O when a source of CA was added. To assess effectiveness of CA knockdown and efficiency of rescue, zebrafish were raised to 3, 4 and 5 d.p.f. Larvae were euthanized, pooled (~40 embryos to yield *N*=1), weighed and homogenized in Tris reaction medium (as described above). To calculate the CA activity (pmol min⁻¹ μg⁻¹ wet weight) in the larvae, the buffer capacity of the medium was first measured before each assay using 70 μl of 0.1 mol l⁻¹ HCl to ensure a sufficiently high buffering capacity (between 4 and 5.5 μmol H⁺ per pH unit). The buffer capacity was used to calculate the uncatalysed rate of CO₂ hydration (without CA source added). The uncatalysed rate was subtracted from the measured rate in the reaction chamber after the addition of sample to determine the rate of CA activity in the larvae.

Series 5 – localization of cytosolic CA in zebrafish NECs by immunocytochemistry

To determine whether larval zebrafish NECs are enriched with cytosolic CA, 4 d.p.f. larvae were immunostained with anti-5-HT rabbit polyclonal antibody (Sigma) and a custom-synthesized, homologous rabbit polyclonal antibody against zCAc (see above) conjugated with biotin. The same 5-HT antibody was used previously to label NECs in zebrafish larvae (e.g. Coccimiglio and Jonz, 2012). Larvae were killed by MS-222 overdose, fixed with 4% paraformaldehyde solution (prepared in phosphate-buffered saline, PBS) for 2 h at room temperature (RT), briefly rinsed in PBS and stored in 100% ethanol at -20°C until use. For whole mount ICC, larvae were rehydrated in PBS and subjected to heat-induced antigen retrieval as described previously (Inoue and Wittbrodt, 2011). After antigen retrieval, larvae were blocked with 10% natural donkey goat serum in 5% Triton-X/PBS (PBST) for 2 h at RT. Larvae were first incubated overnight with 5-HT antibody (1:300 in PBS), rinsed with PBS and stained with a secondary antibody (Alexa-546 conjugated anti-rabbit IgG; Invitrogen; 1:400) in PBS for 2 h at RT. Subsequently, larvae were blocked for 2 h at RT with unconjugated streptavidin (30 μg ml⁻¹ in PBS) to block endogenous biotin, incubated overnight at RT with anti-CA antibody (1:200 in PBS). After rinsing in PBS,

samples were stained with Alexa-488 conjugated streptavidin (2.5 $\mu\text{g ml}^{-1}$ in PBS) for 2 h at RT and washed in PBS. Samples were then mounted onto microscope slides and observed with an Olympus FV1000 BX61 confocal microscope with Olympus Fluoview software. To test whether the block of endogenous biotin was successful, experiments were performed in which the primary antibody was omitted; no staining was observed.

Statistical analyses

All data are presented as means \pm s.e.m. Data were analysed either by two-way repeated measures ANOVA, one-way ANOVA or paired Student's *t*-tests. When appropriate, ANOVA was followed by a Holm–Sidak *post hoc* analysis to determine statistical differences between data points within a series. All statistical analyses were completed using commercial software (SigmaPlot 9, SPSS Inc.). A *P*-value of <0.05 was set to determine statistical significance.

Acknowledgements

We are grateful to Vishal Saxena and Bill Fletcher for their support in caring for the zebrafish used in this study.

Competing interests

The authors declare no competing financial interests.

Author contributions

S.F.P. conceived of and designed the experimental approach. S.M., J.P., J.B. and Y.K. performed the experiments. The manuscript was written by S.F.P. using the MSc thesis of S.M. as an initial draft. All authors contributed to revising the original manuscript.

Funding

The research was supported by National Sciences and Engineering Council of Canada (NSERC) Discovery and Research Tools and Instruments (RTI) grants to S.F.P.

References

- Abdallah, S. J., Perry, S. F. and Jonz, M. G. (2012). CO₂ signaling in chemosensory neuroepithelial cells of the zebrafish gill filaments: role of intracellular Ca²⁺ and pH. *Adv. Exp. Med. Biol.* **758**, 143–148.
- Abdallah, S. J., Jonz, M. G. and Perry, S. F. (2014). Extracellular H⁺ induces Ca²⁺ signals in respiratory chemoreceptors of zebrafish. *Pflügers Arch.* [Epub ahead of print] doi: 10.1007/s00424-014-1514-2
- Bagatto, B. (2005). Ontogeny of cardiovascular control in zebrafish (*Danio rerio*): effects of developmental environment. *Comp. Biochem. Physiol.* **141A**, 391–400.
- Barrionuevo, W. R. and Burggren, W. W. (1999). O₂ consumption and heart rate in developing zebrafish (*Danio rerio*): influence of temperature and ambient O₂. *Am. J. Physiol.* **276**, R505–R513.
- Burleson, M. L. and Smatresk, N. J. (2000). Branchial chemoreceptors mediate ventilatory responses to hypercapnic acidosis in channel catfish. *Comp. Biochem. Physiol.* **125A**, 403–414.
- Burleson, M. L., Mercer, S. E. and Wilk-Blaszczak, M. A. (2006). Isolation and characterization of putative O₂ chemoreceptor cells from the gills of channel catfish (*Ictalurus punctatus*). *Brain Res.* **1092**, 100–107.
- Coates, E. L., Li, A. H. and Nattie, E. E. (1991). Acetazolamide on the ventral medulla of the cat increases phrenic output and delays the ventilatory response to CO₂. *J. Physiol.* **441**, 433–451.
- Coates, E. L., Knuth, S. L. and Bartlett, D., Jr (1996). Laryngeal CO₂ receptors: influence of systemic PCO₂ and carbonic anhydrase inhibition. *Respir. Physiol.* **104**, 53–61.
- Coccimiglio, M. L. and Jonz, M. G. (2012). Serotonergic neuroepithelial cells of the skin in developing zebrafish: morphology, innervation and oxygen-sensitive properties. *J. Exp. Biol.* **215**, 3881–3894.
- Crocker, C. E., Farrell, A. P., Gamperl, A. K. and Cech, J. J., Jr (2000). Cardiorespiratory responses of white sturgeon to environmental hypercapnia. *Am. J. Physiol.* **279**, R617–R628.
- Dejours, P. (1973). Problems of control of breathing in fishes. In *Comparative Physiology: Locomotion, Respiration, Transport and Blood* (ed. L. Bolis, K. Schmidt-Nielsen and S. H. P. Maddrell), pp. 117–133. Amsterdam; New York, NY: North Holland/American Elsevier.
- Dunel-Erb, S., Bailly, Y. and Laurent, P. (1982). Neuroepithelial cells in fish gill primary lamellae. *J. Appl. Physiol.* **53**, 1342–1353.
- Gilmour, K. M. (2001). The CO₂/pH ventilatory drive in fish. *Comp. Biochem. Physiol.* **130A**, 219–240.
- Gilmour, K. M. and Perry, S. F. (2007). Branchial chemoreceptor regulation of cardiorespiratory function. In *Sensory Systems Neuroscience* (ed. T. J. Hara and B. Zielinski), pp. 97–151. San Diego, CA: Academic Press.
- Gilmour, K. M., Milsom, W. K., Rantin, F. T., Reid, S. G. and Perry, S. F. (2005). Cardiorespiratory responses to hypercapnia in tambaqui *Colossoma macropomum*: chemoreceptor orientation and specificity. *J. Exp. Biol.* **208**, 1095–1107.
- Gilmour, K. M., Thomas, K., Esbaugh, A. J. and Perry, S. F. (2009). Carbonic anhydrase expression and CO₂ excretion during early development in zebrafish *Danio rerio*. *J. Exp. Biol.* **212**, 3837–3845.
- Heisler, N., Toews, D. P. and Holetton, G. F. (1988). Regulation of ventilation and acid-base status in the elasmobranch *Scyliorhinus stellaris* during hyperoxia-induced hypercapnia. *Respir. Physiol.* **71**, 227–246.
- Henry, R. P. (1991). Techniques for measuring carbonic anhydrase activities *in vitro*: the electrometric delta pH and pH stat assays. In *The Carbonic Anhydrases: Cellular Physiology and Molecular Genetics* (ed. S. J. Dodgson, R. E. Tashian, G. Gros, and N. D. Carter), pp. 119–126. New York, NY: Plenum.
- Hsieh, D. J. and Liao, C. F. (2002). Zebrafish M₂ muscarinic acetylcholine receptor: cloning, pharmacological characterization, expression patterns and roles in embryonic bradycardia. *Br. J. Pharmacol.* **137**, 782–792.
- Inoue, D. and Wittbrodt, J. (2011). One for all – a highly efficient and versatile method for fluorescent immunostaining in fish embryos. *PLoS ONE* **6**, e19713.
- Iturriaga, R. (1993). Carotid body chemoreception: the importance of CO₂-HCO₃⁻ and carbonic anhydrase. *Biol. Res.* **26**, 319–329.
- Iturriaga, R., Lahiri, S. and Mokashi, A. (1991). Carbonic anhydrase and chemoreception in the cat carotid body. *Am. J. Physiol.* **261**, C565–C573.
- Jacob, E., Drexel, M., Schwerte, T. and Pelster, B. (2002). Influence of hypoxia and of hypoxemia on the development of cardiac activity in zebrafish larvae. *Am. J. Physiol.* **283**, R911–R917.
- Janssen, R. G. and Randall, D. J. (1975). The effects of changes in pH and PCO₂ in blood and water on breathing in rainbow trout, *Salmo gairdneri*. *Respir. Physiol.* **25**, 235–245.
- Jonz, M. G. and Nurse, C. A. (2005). Development of oxygen sensing in the gills of zebrafish. *J. Exp. Biol.* **208**, 1537–1549.
- Jonz, M. G., Fearon, I. M. and Nurse, C. A. (2004). Neuroepithelial oxygen chemoreceptors of the zebrafish gill. *J. Physiol.* **560**, 737–752.
- Kinkead, R. and Perry, S. F. (1991). The effects of catecholamines on ventilation in rainbow trout during hypoxia or hypercapnia. *Respir. Physiol.* **84**, 77–92.
- Kwong, R. W. M., Kumai, Y. and Perry, S. F. (2014). The physiology of fish at low pH: the zebrafish as a model system. *J. Exp. Biol.* **217**, 651–662.
- Lahiri, S. and Forster, R. E., II (2003). CO₂/H⁺ sensing: peripheral and central chemoreception. *Int. J. Biochem. Cell Biol.* **35**, 1413–1435.
- McKendry, J. E. and Perry, S. F. (2001). Cardiovascular effects of hypercapnia in rainbow trout (*Oncorhynchus mykiss*): a role for externally oriented chemoreceptors. *J. Exp. Biol.* **204**, 115–125.
- McKendry, J. E., Milsom, W. K. and Perry, S. F. (2001). Branchial CO₂ receptors and cardiorespiratory adjustments during hypercapnia in Pacific spiny dogfish (*Squalus acanthias*). *J. Exp. Biol.* **204**, 1519–1527.
- Milsom, W. K. (1995). The role of CO₂/pH chemoreceptors in ventilatory control. *Braz. J. Med. Biol. Res.* **28**, 1147–1160.
- Milsom, W. K. and Burleson, M. L. (2007). Peripheral arterial chemoreceptors and the evolution of the carotid body. *Respir. Physiol. Neurobiol.* **157**, 4–11.
- Montpetit, C. J. and Perry, S. F. (1999). Neuronal control of catecholamine secretion from chromaffin cells in the rainbow trout (*Oncorhynchus mykiss*). *J. Exp. Biol.* **202**, 2059–2069.
- Olson, K. R., Healy, M. J., Qin, Z., Skovgaard, N., Vulesevic, B., Duff, D. W., Whitfield, N. L., Yang, G., Wang, R. and Perry, S. F. (2008). Hydrogen sulfide as an oxygen sensor in trout gill chemoreceptors. *Am. J. Physiol.* **295**, R669–R680.
- Pelster, B. (2002). Developmental plasticity in the cardiovascular system of fish, with special reference to the zebrafish. *Comp. Biochem. Physiol.* **133A**, 547–553.
- Pena-Munzenmayer, G., Niemeier, M. I., Sepulveda, F. V. and Cid, L. P. (2014). Zebrafish and mouse TASK-2 K⁺ channels are inhibited by increased CO₂ and intracellular acidification. *Pflügers Arch.* **466**, 1317–1327.
- Perry, S. F. and Abdallah, S. (2012). Mechanisms and consequences of carbon dioxide sensing in fish. *Respir. Physiol. Neurobiol.* **184**, 309–315.
- Perry, S. F. and Bernier, N. J. (1999). The acute humoral adrenergic stress response in fish: facts and fiction. *Aquaculture* **177**, 285–295.
- Perry, S. F. and Capaldo, A. (2011). The autonomic nervous system and chromaffin tissue: neuroendocrine regulation of catecholamine secretion in non-mammalian vertebrates. *Auton. Neurosci.* **165**, 54–66.
- Perry, S. F., Esbaugh, A., Braun, M. and Gilmour, K. M. (2009a). Gas transport and gill function in water-breathing fish. In *Cardio-Respiratory Control in Vertebrates: Comparative and Evolutionary Aspects* (ed. M. L. Glass and S. C. Wood), pp. 5–42. Berlin; Heidelberg: Springer.
- Perry, S. F., Fritsche, R., Hoagland, T. M., Duff, D. W. and Olson, K. R. (1999). The control of blood pressure during external hypercapnia in the rainbow trout (*Oncorhynchus mykiss*). *J. Exp. Biol.* **202**, 2177–2190.
- Perry, S. and Gilmour, K. (1996). Consequences of catecholamine release on ventilation and blood oxygen transport during hypoxia and hypercapnia in an elasmobranch *Squalus acanthias* and a teleost *Oncorhynchus mykiss*. *J. Exp. Biol.* **199**, 2105–2118.
- Perry, S. F. and Gilmour, K. M. (2002). Sensing and transfer of respiratory gases at the fish gill. *J. Exp. Zool.* **293**, 249–263.
- Perry, S. F. and McKendry, J. E. (2001). The relative roles of external and internal CO₂ versus H⁺ in eliciting the cardiorespiratory responses of *Salmo salar* and *Squalus acanthias* to hypercapnia. *J. Exp. Biol.* **204**, 3963–3971.
- Perry, S. F. and Reid, S. G. (2002). Cardiorespiratory adjustments during hypercapnia in rainbow trout *Oncorhynchus mykiss* are initiated by external CO₂ receptors on the first gill arch. *J. Exp. Biol.* **205**, 3357–3365.
- Perry, S. F., Vulesevic, B., Braun, M. and Gilmour, K. M. (2009b). Ventilation in Pacific hagfish (*Eptatretus stoutii*) during exposure to acute hypoxia or hypercapnia. *Respir. Physiol. Neurobiol.* **167**, 227–234.

- Perry, S. F. and Wood, C. M. (1989). Control and coordination of gas transfer in fishes. *Can. J. Zool.* **67**, 2961-2970.
- Qin, Z., Lewis, J. E. and Perry, S. F. (2010). Zebrafish (*Danio rerio*) gill neuroepithelial cells are sensitive chemoreceptors for environmental CO₂. *J. Physiol.* **588**, 861-872.
- Randall, D. J. (1982). The control of respiration and circulation in fish during exercise and hypoxia. *J. Exp. Biol.* **100**, 275-288.
- Randall, D. J., Heisler, N. and Drees, F. (1976). Ventilatory response to hypercapnia in the larger spotted dogfish *Scyliorhinus stellaris*. *Am. J. Physiol.* **230**, 590-594.
- Randall, D. J. and Jones, D. R. (1973). The effect of deafferentation of the pseudobranch on the respiratory response to hypoxia and hyperoxia in the trout (*Salmo gairdneri*). *Respir. Physiol.* **17**, 291-301.
- Randall, D. J. and Shelton, G. (1963). The effects of changes in environmental gas concentrations on the breathing and heart rate of a teleost fish. *Comp. Biochem. Physiol.* **9**, 229-239.
- Reid, S. G., Bernier, N. J. and Perry, S. F. (1998). The adrenergic stress response in fish: control of catecholamine storage and release. *Comp. Biochem. Physiol.* **120C**, 1-27.
- Reid, S. G., Sundin, L., Kalinin, A. L., Rantin, F. T. and Milsom, W. K. (2000). Cardiovascular and respiratory reflexes in the tropical fish, traira (*Hoplias malabaricus*): CO₂/pH chemoresponses. *Respir. Physiol.* **120**, 47-59.
- Schwerte, T., Prem, C., Mairösl, A. and Pelster, B. (2006). Development of the sympatho-vagal balance in the cardiovascular system in zebrafish (*Danio rerio*) characterized by power spectrum and classical signal analysis. *J. Exp. Biol.* **209**, 1093-1100.
- Shelton, G., Jones, D. R. and Milsom, W. K. (1986). Control of breathing in ectothermic vertebrates. In *Handbook of Physiology, Section 3. The Respiratory System: Control of Breathing*, Vol. 2 (ed. N. S. Cherniak and J. G. Widdicombe), pp. 857-909. Bethesda, MD: American Physiological Society.
- Smatresk, N. J. and Cameron, J. N. (1982). Respiration and acid-base physiology of the spotted gar, a bimodal breather. II. Responses to temperature change and hypercapnia. *J. Exp. Biol.* **96**, 281-293.
- Smith, F. M. and Jones, D. R. (1982). The effect of changes in blood oxygen-carrying capacity on ventilation volume in the rainbow trout (*Salmo gairdneri*). *J. Exp. Biol.* **97**, 325-334.
- Steele, S. L., Lo, K. H., Li, V. W., Cheng, S. H., Ekker, M. and Perry, S. F. (2009). Loss of M₂ muscarinic receptor function inhibits development of hypoxic bradycardia and alters cardiac β-adrenergic sensitivity in larval zebrafish (*Danio rerio*). *Am. J. Physiol.* **297**, R412-R420.
- Steele, S. L., Yang, X., Debais-Thibaud, M., Schwerte, T., Pelster, B., Ekker, M., Tiberi, M. and Perry, S. F. (2011). *In vivo* and *in vitro* assessment of cardiac β-adrenergic receptors in larval zebrafish (*Danio rerio*). *J. Exp. Biol.* **214**, 1445-1457.
- Sundin, L., Reid, S. G., Rantin, F. T. and Milsom, W. K. (2000). Branchial receptors and cardiorespiratory reflexes in a neotropical fish, the tambaqui (*Colossoma macropomum*). *J. Exp. Biol.* **203**, 1225-1239.
- Vulesevic, B., McNeill, B. and Perry, S. F. (2006). Chemoreceptor plasticity and respiratory acclimation in the zebrafish *Danio rerio*. *J. Exp. Biol.* **209**, 1261-1273.
- Vulesevic, B. and Perry, S. F. (2006). Developmental plasticity of ventilatory control in zebrafish, *Danio rerio*. *Respir. Physiol. Neurobiol.* **154**, 396-405.
- Wang, Z., Nishimura, Y., Shimada, Y., Umemoto, N., Hirano, M., Zang, L., Oka, T., Sakamoto, C., Kuroyanagi, J. and Tanaka, T. (2009). Zebrafish beta-adrenergic receptor mRNA expression and control of pigmentation. *Gene* **446**, 18-27.
- Wood, C. and Munger, R. (1994). Carbonic anhydrase injection provides evidence for the role of blood acid-base status in stimulating ventilation after exhaustive exercise in rainbow trout. *J. Exp. Biol.* **194**, 225-253.
- Wood, C. M., Turner, J. D., Munger, R. S. and Graham, M. S. (1990). Control of ventilation in the hypercapnic skate *Raja ocellata*. II. Cerebrospinal fluid and intracellular pH in the brain and other tissues. *Respir. Physiol.* **80**, 279-297.